

The power of slicing in noncommutative geometry

Raf Bocklandt

Geert Van de Weyer*

Abstract

In this note we will show how one can describe the representation varieties of certain Fuchsian groups using the Luna Slice theorem and the technique of local quivers. We will illustrate with one specific example: the Fuchsian group $\mathbb{Z}_2 * \mathbb{Z}_4$.

1 Introduction

Fuchsian groups are the crystallographic groups of the hyperbolic plane, i.e. the discrete subgroups of $\mathrm{PSL}_2(\mathbb{R})$. The orientation-preserving Fuchsian groups can be presented in the following form

- generators

hyperbolic: $a_1, b_1, \dots, a_g, b_g$

elliptic: x_1, \dots, x_e

parabolic: y_1, \dots, y_p

- relations

$$x_1^{m_1} = \dots x_e^{m_e} = 1$$

$$x_1 \cdots x_e y_1 \cdots y_p [a_1, b_1] \cdots [a_g, b_g] = 1$$

*The authors are Postdoctoral Fellows of the Fund for Scientific Research - Flanders (Belgium)(F.W.O. - Vlaanderen)


Received by the editors May 2005.

Communicated by H. Van Maldeghem.

If the group has parabolic generators we can use one to get rid of the last relation, so that it can also be written as a free product of cyclic groups.

$$\mathbb{Z}_{m_1} * \cdots * \mathbb{Z}_{m_e} * \mathbb{Z}^{*(p+2g-1)}$$

For instance the orientation-preserving symmetry group of the Escher-like picture below is $G = \mathbb{Z}_2 * \mathbb{Z}_4$.



2 The representation spaces of G

To G we can assign the variety of n -dimensional representations

$$\begin{aligned} \mathbf{rep}(G, n) &:= \{\rho : G \rightarrow \mathrm{GL}_n : \rho(gh) = \rho(g)\rho(h)\} \\ &\cong \{(X, Y) \in \mathrm{GL}_n \times \mathrm{GL}_n \mid X^2 = Y^4 = 1\} \end{aligned}$$

On this space we have a GL_n -action by conjugation: $g \cdot (X, Y) = (gXg^{-1}, gYg^{-1})$. Orbits under this action correspond to isomorphism classes of representations and if one takes the algebraic quotient of this action one obtains a new variety $\mathbf{iss}(G, n)$ which classifies the semisimple representations of G up to isomorphism (see [9]). The projection map $\pi : \mathbf{rep}(G, n) \rightarrow \mathbf{iss}(G, n)$ maps every representation to the isomorphism class of its semisimplification.

The main goal of geometric representation theory is now the study this map and find the answer to the following questions:

- what is the geometry of $\mathbf{rep}(G, n)$, does it contain singularities and if so what is their nature?
- what is the geometry of $\mathbf{iss}(G, n)$, does it contain singularities and if so what is their nature?
- what do the fibers of π look like?

In this case the first question can be answered quickly using only the theory of representations of finite groups, because $\mathbf{rep}(G, n) = \mathbf{rep}(\mathbb{Z}_2, n) \times \mathbf{rep}(\mathbb{Z}_4, n)$. The two factors contain each a finite number of disjoint conjugation classes of diagonal matrices with eigenvalues ± 1 resp. $\pm 1, \pm i$.

This fact can be used to split $\mathbf{rep}(G, n)$ into disjoint components corresponding to a dimension vector, which is an element in \mathbb{N}^{2+4} consisting of the dimensions of

the eigenspaces of X (with eigenvalues ± 1) and Y (with eigenvalues $\pm 1, \pm i$). Each of these components is smooth because it is the product of 2 conjugation classes of matrices and we will denote these varieties by $\mathbf{rep}(\mathbf{G}, \delta)$ with $\delta \in \mathbb{N}^{2+4}$. We can also split the quotient map π and its target to obtain quotient varieties denoted as $\mathbf{iss}(\mathbf{G}, \delta)$.

3 The Luna Slice Theorem

The fact that the representation spaces $\mathbf{rep}(\mathbf{G}, \delta)$ are smooth provides us with a very powerful tool to study both the representation spaces and their quotient spaces $\mathbf{iss}(\mathbf{G}, \delta)$, the *Luna Slice Theorem*. This theorem is an algebraic version of the classical slice theorem in differential geometry, using étale morphisms instead of diffeomorphisms.

Recall that a morphism $\varphi : X \rightarrow Y$ between algebraic varieties is called *étale* if it is a *smooth morphism with finite fibers*. If both X and Y are smooth, this is equivalent to the fact that $d_x\varphi : T_xX \rightarrow T_{\varphi(x)}Y$ is an isomorphism. The Luna Slice Theorem now states

Theorem 1 (Luna, [12]). *Let $x \in \mathbf{rep}(\mathbf{G}, \delta)$ be a semisimple representation and let H be the stabilizer of x in $GL_n(\mathbb{C})$. Let N be the normal space to the small tangent space $T_x GL_n(\mathbb{C}).x$ in the big tangent space $T_x \mathbf{rep}(\mathbf{G}, \delta)$. Then there exists a locally closed affine subset $S \subset \mathbf{rep}(\mathbf{G}, \delta)$, called a slice, such that*

1. $x \in S$;
2. S is stable under the action of H ;
3. the action map $GL_n(\mathbb{C}) \times S \rightarrow \mathbf{rep}_\alpha \mathbf{G} : (g, s) \mapsto g.s$ induces an étale and $GL_n(\mathbb{C})$ -equivariant morphism

$$\psi : GL_n(\mathbb{C}) \times^H S \rightarrow \mathbf{rep}(\mathbf{G}, \delta)$$

for which the induced morphism

$$\psi // GL_n(\mathbb{C}) : S // H \rightarrow \mathbf{iss}(\mathbf{G}, \delta)$$

is étale;

4. there exists a H -equivariant map $\varphi : S \rightarrow N = T_x S$ with affine image and $\varphi(x) = 0$ such that the induced map

$$\varphi // H : S // H \rightarrow N // H$$

is étale;

5. we have a commutative diagram

$$\begin{array}{ccccc}
 & & GL_n(\mathbb{C}) \times^H S & & \\
 & \swarrow^{GL_n(\mathbb{C}) \times^H \varphi} & \downarrow // GL_n(\mathbb{C}) & \searrow^\psi & \\
 GL_n(\mathbb{C}) \times^H N & & S // H & & \mathbf{rep}(\mathbf{G}, \delta) \\
 \downarrow // GL_n(\mathbb{C})_{\varphi // H} & & \searrow^{\psi // GL_n(\mathbb{C})} & & \downarrow // GL_n(\mathbb{C}) \\
 N // H & & & & \mathbf{iss}(\mathbf{G}, \delta).
 \end{array}$$

4 Local Quivers

To get a combinatorial description of the Luna Slice Theorem we need quivers.

A *quiver* $Q = (V, A, h, t)$ consists of a set of vertices V , a set of arrows A between those vertices and maps $h, t : A \rightarrow V$ which assign to each arrow its head and tail.

A *dimension vector* of a quiver is a map $\alpha : V \rightarrow \mathbb{N}$ and a couple (Q, α) consisting of a quiver and a dimension vector is called a *quiver setting* and for every vertex $v \in V$, α_v is referred to as the dimension of v . In the examples the dimensions are written inside the vertices.

An α -dimensional complex representation W of Q assigns to each vertex v a linear space \mathbb{C}^{α_v} and to each arrow a a matrix

$$W_a \in \text{Mat}_{\alpha_{h(a)} \times \alpha_{t(a)}}(\mathbb{C})$$

The space of all α -dimensional representations is denoted by $\text{rep}_\alpha Q$.

$$\text{rep}(Q, \alpha) := \bigoplus_{a \in A} \text{Mat}_{\alpha_{h(a)} \times \alpha_{t(a)}}(\mathbb{C})$$

To the dimension vector α we can also assign a reductive group

$$\text{GL}_\alpha := \prod_{v \in V} \text{GL}_{\alpha_v}(\mathbb{C}).$$

An element of this group, g , has a natural action on $\text{rep}(Q, \alpha)$:

$$W := (W_a)_{a \in Q_1}, \quad W^g := (g_{h(a)} W_a g_{t(a)}^{-1})_{a \in Q_1}$$

Using this definitions the normal space N from the previous section can be described as follows:

Theorem 2. *Let x be a semisimple representation in $\text{rep}(\mathbf{G}, n)$ with decomposition in simples*

$$x = s_1^{\oplus e_1} \oplus \cdots \oplus s_k^{\oplus e_k},$$

and let Q_x be the quiver with

- vertex set $\{1, \dots, k\}$;
- the number of arrows between vertex i and vertex j given by $\dim \text{Ext}_A^1(s_i, s_j)$.

Let $\alpha_x = (e_1, \dots, e_k)$, then

$$N \cong \text{rep}(Q_x, \alpha_x) \quad \text{and} \quad H \cong \text{GL}_{\alpha_x}(\mathbb{C}),$$

so

$$N//H \cong \text{iss}(Q_x, \alpha_x).$$

Combining this description with the Luna Slice Theorem, we are able to study étale invariant properties such as the existence of simple representations and smoothness by passing from the representation spaces of \mathbf{G} to the representation space of a quiver, where the latter has as a great advantage that a lot of properties can be computed combinatorially.

5 Applications

5.1 Which $\text{rep}(\mathbf{G}, \delta)$ contain simple representations?

To solve this problem we start with the set of 8 one-dimensional simples. These elementary simples $S_{\xi\eta}$ let X and Y act as scalars ξ, η with $\xi^2 = \eta^4 = 1$ and have dimension vectors $(1, 0|1, 0, 0, 0), \dots, (0, 1|0, 0, 0, 1)$.

As every dimension vector can be seen as a sum of these basic dimension vectors, we can find in every $\text{iss}(\mathbf{G}, \delta)$ a semisimple representation whose factors are these one-dimensional simples. If we fix such a representation R , we can look at its local quiver setting (Q_R, α_R) . The fact that simplicity is a Zariski-open condition and $\text{rep}(\mathbf{G}, \delta)$ is irreducible implies that either every (étale) neighborhood of R will contain simples or $\text{rep}(\mathbf{G}, \delta)$ does not contain simples. Therefore $\text{rep}(\mathbf{G}, \delta)$ will contain simple representations if and only if $\text{rep}(Q_R, \alpha_R)$ does.

For instance, the representation space with dimensionvector $\alpha = (2, 1|1, 1, 1, 0)$ contains the representation $S_{11} \oplus S_{1-1} \oplus S_{-1i}$. The local quiver looks like



because one can easily check that the space of extensions between two elementary simples is zero if they have an eigenvalue in common and one-dimensional otherwise.

The local quiver setting has a simple representation that assigns to all arrows a nonzero scalar, hence $\text{rep}(\mathbf{G}, \delta)$ contains simple representations.

Using the classification of quiver settings with simple representations by Le Bruyn and Procesi in [7], one can obtain a criterium that determines whether $\text{rep}_\alpha G$ contains simples


Theorem 3. [2] Let $\delta = (a_1, a_2, b_1, b_2, b_3, b_4)$ with $a_1 + a_2 = b_1 + \dots + b_4 = n$, then $\text{rep}(\mathbf{G}, \delta)$ contains simples if and only if $\forall i, j : a_i + b_j \leq n$.

5.2 Which points in $\text{iss}(\mathbf{G}, \delta)$ are smooth?


To check whether a given point is smooth, one only has to check whether the corresponding local quiver setting has a smooth quotient space. This can be done using the technique of reduction moves.

To a local quiver one can apply the following reduction moves which keep the quotient space intact up to a product with an affine space.


\mathcal{R}_I If $\sum_{j=1}^k i_j \leq \alpha_v$ or $\sum_{j=1}^l u_j \leq \alpha_v$ we delete the vertex v and connect all the arrows.



\mathcal{R}_{II} Remove the loops on a vertex with dimension 1.




\mathcal{R}_{III} Remove the only loop on a vertex with dimension $k > 1$ which has a neighborhood like the picture below or its dual with the arrows reversed.




After applying all possible reduction step one obtains a *reduced local quiver*. To check whether its quotient space is smooth one can use the following theorem

Theorem 4. [3] $\text{rep}(Q, \alpha)$ is smooth if and only if the strongly connected components of the corresponding reduced setting are listed below:



As an example let us again take the representation $R = S_{11} \oplus S_{1-1} \oplus S_{-1i}$. The local quiver can be reduced as follows:



so the image of R in $\text{rep}(\mathbf{G}, \delta)$ is a smooth point.

In general there are only a finite number of local quiver settings that can occur in $\text{rep}(\mathbf{G}, \delta)$. By studying all of these one can also obtain a list of dimension vectors whose quotient spaces are smooth.

Theorem 5. [1] Let $\delta = (a_1, a_2, b_1, b_2, b_3, b_4)$ with $a_1 + a_2 = b_1 + \dots + b_4 = n$, then $\text{iss}(\mathbf{G}, \delta)$ is a smooth space, if up to permutation of the a 's and the b 's, the dimension vector is one of the following forms

$$(1, 0|1, 0, 0, 0) \quad (1, 1|1, 1, 0, 0) \quad (2, 1|1, 1, 1, 0) \\ (3, 1|1, 1, 1, 1) \quad (2, 2|2, 1, 1, 0) \quad (4, 2|2, 2, 2, 0).$$

5.3 Which singularities can occur?

The notion of reduced quiver settings can also be used in principle to give a list of all possible singularities that can occur in a given dimension. This follows from the fact the for a fixed n there are only a finite number of reduced quivers whose $\text{iss}(Q, \alpha)$ has dimension n . A classification up to dimension 6 has been given in [4].

5.4 What fibers can occur?

Using the Luna Slice Theorem the question of the fibers can again be tracked back to calculating the nullcone of the local quiver $\text{Null}(Q, \alpha) = \{W \in \text{rep}(Q, \alpha) | 0 \in \text{GL}_n^- \cdot W\}$. In [6], the authors obtained a list of the local quivers whose nullcone has minimal dimension, which allows one to calculate the fibers with generic dimension.

6 The Future

The methods explained above work not only for the specific group we chose as example but more in general for formally smooth algebras (see [11]). Other examples of these algebras include group algebras of trees of groups, universal localizations of path algebra and free products of coordinate rings of curves.

The framework however breaks down if the representation space is not smooth. To conclude this note, we propose two possible pathways to extend the results to these algebras.

- The first path is to use the Luna Slice theorem for singular points to extend the local description to a broader class of algebras. A first step in this approach was made by W. Crawley-Boevey in [8] in which he obtained a local description of representation spaces of deformed preprojective algebras.
- A second approach is to first blow up the singularities in the representation space and then to take the quotient. The formalism of quivers then extends to quivers with an automorphism action on it. First steps in this direction have been explored by L. Le Bruyn, S. Symens and the first author in [5] and [10].

References

- [1] Jan Adriaenssens, Raf Bocklandt, and Geert Van de Weyer, *Smooth character varieties for torus knot groups*, *Comm. Algebra* **30** (2002), no. 6, 3045–3061. MR MR1908257 (2004c:16015)
- [2] Jan Adriaenssens and Lieven Le Bruyn, *Local quivers and stable representations*, *Comm. Algebra* **31** (2003), no. 4, 1777–1797. MR MR1972892 (2004c:16016)
- [3] Raf Bocklandt, *Smooth quiver representation spaces*, *J. Algebra* **253** (2002), no. 2, 296–313. MR MR1929191 (2003i:16020)
- [4] Raf Bocklandt, Lieven Le Bruyn, and Geert Van de Weyer, *Smooth order singularities*, *J. Algebra Appl.* **2** (2003), no. 4, 365–395. MR MR2020934 (2005e:16029)
- [5] Raf Bocklandt and Stijn Symens, *The local structure of graded representations*, 2005.
- [6] Raf Bocklandt and Geert Van de Weyer, *Cofree quiver representations*, 2004.
- [7] Lieven Le Bruyn and Claudio Procesi, *Semisimple representations of quivers*, **317** (1990), 585–598.
- [8] William Crawley-Boevey, *Decomposition of Marsden-Weinstein reductions for representations of quivers*, *Compositio Math.* **130** (2002), no. 2, 225–239. MR MR1883820 (2003j:16019)
- [9] Hanspeter Kraft, *Geometrische methoden in der invariantentheorie*, *Aspekte der Mathematik*, vol. D1, Vieweg-Verlag, Braunschweig, 1984.

- [10] Lieven Le Bruyn, *Etale cohomology in non-commutative geometry*, 1997, UIA-preprint, available from the author's website.
- [11] Lieven Le Bruyn, *One quiver to rule them all*, 2003, arXiv math.RA/0304196.
- [12] Domingo Luna, *Slices etales*, **33** (1973), 81–105.

University of Antwerp
Middelheimlaan 1
B-2020 Antwerpen (Belgium)
email: rafael.bocklandt@ua.ac.be, geert.vandeweyer@ua.ac.be