

## Research Article

# Products of Composition and Differentiation Operators from $Q_K(p, q)$ Spaces to Bloch-Type Spaces

**Weifeng Yang**

*Department of Mathematics and Physics, Hunan Institute of Engineering, Xiangtan 411104, China*

Correspondence should be addressed to Weifeng Yang, yangweifeng09@163.com

Received 25 March 2009; Accepted 16 April 2009

Recommended by Stevo Stevic

We study the boundedness and compactness of the products of composition and differentiation operators from  $Q_K(p, q)$  spaces to Bloch-type spaces and little Bloch-type spaces.

Copyright © 2009 Weifeng Yang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

Let  $\mathbb{D}$  be the open unit disk in the complex plane and  $H(\mathbb{D})$  the class of all analytic functions on  $\mathbb{D}$ . The  $\alpha$ -Bloch space  $\mathcal{B}^\alpha$  ( $\alpha > 0$ ) on  $\mathbb{D}$  is the space of all analytic functions  $f$  on  $\mathbb{D}$  such that

$$\|f\|_{\mathcal{B}^\alpha} = |f(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2)^\alpha |f'(z)| < \infty. \quad (1.1)$$

Under the above norm,  $\mathcal{B}^\alpha$  is a Banach space. When  $\alpha = 1$ ,  $\mathcal{B}^1 = \mathcal{B}$  is the well-known Bloch space. Let  $\mathcal{B}_0^\alpha$  denote the subspace of  $\mathcal{B}^\alpha$  consisting of those  $f \in \mathcal{B}^\alpha$  for which  $(1 - |z|^2)^\alpha |f'(z)| \rightarrow 0$  as  $|z| \rightarrow 1$ . This space is called the little  $\alpha$ -Bloch space.

Assume that  $\mu$  is a positive continuous function on  $[0, 1)$ , and there exist positive numbers  $s$  and  $t$ ,  $0 < s < t$ , and  $\delta \in [0, 1)$  such that

$$\begin{aligned} \frac{\mu(r)}{(1-r)^s} \text{ is decreasing on } [\delta, 1), \lim_{r \rightarrow 1} \frac{\mu(r)}{(1-r)^s} = 0, \\ \frac{\mu(r)}{(1-r)^t} \text{ is increasing on } [\delta, 1), \lim_{r \rightarrow 1} \frac{\mu(r)}{(1-r)^t} = \infty, \end{aligned} \quad (1.2)$$

then  $\mu$  is called a normal function (see [1]).

An  $f \in H(\mathbb{D})$  is said to belong to the Bloch-type space  $\mathcal{B}_\mu = \mathcal{B}_\mu(\mathbb{D})$ , if (see, e.g, [2–5])

$$\|f\|_{\mathcal{B}_\mu} = |f(0)| + \sup_{z \in \mathbb{D}} \mu(|z|) |f'(z)| < \infty. \quad (1.3)$$

$\mathcal{B}_\mu$  is a Banach space with the norm  $\|\cdot\|_{\mathcal{B}_\mu}$  (see [3]). When  $\mu(r) = (1 - r^2)^\alpha$ , the induced space  $\mathcal{B}_\mu$  becomes the  $\alpha$ -Bloch space  $\mathcal{B}^\alpha$ .

Throughout this paper, we assume that  $K : [0, \infty) \rightarrow [0, \infty)$  is a nondecreasing continuous function. Assume that  $p > 0$ ,  $q > -2$ . A function  $f \in H(\mathbb{D})$  is said to belong to  $\mathcal{Q}_K(p, q)$  (see [6]) if

$$\|f\| = \left( \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |f'(z)|^p (1 - |z|^2)^q K(g(z, a)) dA(z) \right)^{1/p} < \infty, \quad (1.4)$$

where  $dA$  denotes the normalized Lebesgue area measure in  $\mathbb{D}$  (i.e.,  $A(\mathbb{D}) = 1$ ) and  $g(z, a)$  is the Green function with logarithmic singularity at  $a$ , that is,  $g(z, a) = \log(1/|\varphi_a(z)|)$  ( $\varphi_a$  is a conformal automorphism defined by  $\varphi_a(z) = (a - z)/(1 - \bar{a}z)$  for  $a \in \mathbb{D}$ ). If  $K(x) = x^s$ ,  $s \geq 0$ , the space  $\mathcal{Q}_K(p, q)$  equals to  $F(p, q, s)$ , which is introduced by Zhao in [7]. Moreover (see [7]) we have that,  $F(p, q, s) = \mathcal{B}^{(q+2)/p}$ , and  $F_0(p, q, s) = \mathcal{B}_0^{(q+2)/p}$  for  $s > 1$ ,  $F(p, q, s) \subseteq \mathcal{B}^{(q+2)/p}$ , and  $F_0(p, q, s) \subseteq \mathcal{B}_0^{(q+2)/p}$  for  $0 < s \leq 1$ ,  $F(2, 0, s) = \mathcal{Q}_s$ , and  $F_0(2, 0, s) = \mathcal{Q}_{s,0}$ ,  $F(2, 1, 0) = H^2$ ,  $F(2, 0, 1) = BMOA$ , and  $F_0(2, 0, 1) = VMOA$ . When  $p \geq 1$ ,  $\mathcal{Q}_K(p, q)$  is a Banach space under the norm

$$\|f\|_{\mathcal{Q}_K(p, q)} = |f(0)| + \|f\|. \quad (1.5)$$

From [6], we know that  $\mathcal{Q}_K(p, q) \subseteq \mathcal{B}^{(q+2)/p}$ ,  $\mathcal{Q}_K(p, q) = \mathcal{B}^{(q+2)/p}$  if and only if

$$\int_0^1 (1 - r^2)^{-2} K(-\log r) r dr < \infty. \quad (1.6)$$

Moreover,  $\|f\|_{\mathcal{B}^{(q+2)/p}} \leq C \|f\|_{\mathcal{Q}_K(p, q)}$  (see in [6, Theorem 2.1] or [8, Lemma 2.1]). Throughout the paper we assume that (see [6])

$$\int_0^1 (1 - r^2)^q K(-\log r) r dr < \infty, \quad (1.7)$$

since otherwise  $\mathcal{Q}_K(p, q)$  consists only of constant functions.

Let  $\varphi$  denote a nonconstant analytic self-map of  $\mathbb{D}$ . Associated with  $\varphi$  is the composition operator  $C_\varphi$  defined by  $C_\varphi f = f \circ \varphi$  for  $f \in H(\mathbb{D})$ . The problem of characterizing the boundedness and compactness of composition operators on many Banach spaces of analytic functions has attracted lots of attention recently, see, for example, [9, 10] and the reference therein.

Let  $D$  be the differentiation operator on  $H(\mathbb{D})$ , that is,  $Df(z) = f'(z)$ . For  $f \in H(\mathbb{D})$ , the products of composition and differentiation operators  $DC_\varphi$  and  $C_\varphi D$  are defined, respectively, by

$$\begin{aligned} DC_\varphi(f) &= (f \circ \varphi)' = f'(\varphi) \varphi', \\ C_\varphi D(f) &= f'(\varphi), \quad f \in H(\mathbb{D}). \end{aligned} \tag{1.8}$$

The boundedness and compactness of  $DC_\varphi$  on the Hardy space were investigated by Hirschweiler and Portnoy in [11] and by Ohno in [12]. The case of the Bergman spaces was studied in [11], while the case of the Hilbert-Bergman space was studied by Stević in [13]. In [14], Li and Stević studied the boundedness and compactness of the operator  $DC_\varphi$  on  $\alpha$ -Bloch spaces, while in [15] they studied these operators between  $H^\infty$  and  $\alpha$ -Bloch spaces. The boundedness and compactness of the operator  $DC_\varphi$  from mixed-norm spaces to  $\alpha$ -Bloch spaces was studied by Li and Stević in [16]. Norm and essential norm of the operator  $DC_\varphi$  from  $\alpha$ -Bloch spaces to weighted-type spaces were studied by Stević in [17]. Some related operators can be also found in [18–21]. For some other papers on products of linear operators on spaces of holomorphic functions, mostly integral-type and composition operators, see, for example, the following papers by Li and Stević: [5, 22–30].

Motivated basically by papers [14, 15], in this paper, we study the operators  $DC_\varphi$  and  $C_\varphi D$  from  $Q_K(p, q)$  space to  $\mathcal{B}_\mu$  and  $\mathcal{B}_{\mu,0}$  spaces. Some sufficient and necessary conditions for the boundedness and compactness of these operators are given.

Throughout this paper, constants are denoted by  $C$ , they are positive and may differ from one occurrence to the other. The notation  $A \asymp B$  means that there is a positive constant  $C$  such that  $B/C \leq A \leq CB$ .

## 2. Main Results and Proofs

In this section we give our main results and proofs. For this purpose, we need some auxiliary results. The following lemma can be proved in a standard way (see, e.g. in [9, Proposition 3.11]). A detailed proof, can be found, for example, in [31].

**Lemma 2.1.** *Let  $\varphi$  be an analytic self-map of  $\mathbb{D}$ . Suppose that  $\mu$  is normal,  $p > 0$ ,  $q > -2$ . Then  $DC_\varphi$  (or  $C_\varphi D$ ) :  $Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is compact if and only if  $DC_\varphi$  (or  $C_\varphi D$ ) :  $Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is bounded and for any bounded sequence  $(f_n)_{n \in \mathbb{N}}$  in  $Q_K(p, q)$  which converges to zero uniformly on compact subsets of  $\mathbb{D}$ , one has  $\|DC_\varphi f_n\|_{\mathcal{B}_\mu} \rightarrow 0$  (or  $\|C_\varphi D f_n\|_{\mathcal{B}_\mu} \rightarrow 0$ ) as  $n \rightarrow \infty$ .*

The following lemma can be proved similarly as [32], one omits the details (see also [2, 4]).

**Lemma 2.2.** *A closed set  $K$  in  $\mathcal{B}_{\mu,0}$  is compact if and only if it is bounded and satisfies*

$$\lim_{|z| \rightarrow 1^-} \sup_{f \in K} \mu(|z|) |f'(z)| = 0. \tag{2.1}$$

Now one is in a position to state and prove the main results of this paper.

**Theorem 2.3.** Let  $\varphi$  be an analytic self-map of  $\mathbb{D}$ . Suppose that  $\mu$  is normal,  $p > 0$ ,  $q > -2$ , and  $K$  is a nonnegative nondecreasing function on  $[0, \infty)$  such that

$$\int_0^1 K(-\log r)(1-r)^{\min\{-1,q\}} \left(\log \frac{1}{1-r}\right)^{X_1(q)} r dr < \infty, \quad (2.2)$$

where  $\chi_O(x)$  denote the characteristic function of the set  $O$ . Then  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is bounded if and only if

$$\sup_{z \in \mathbb{D}} \frac{\mu(|z|)|\varphi'(z)|^2}{(1-|\varphi(z)|^2)^{(2+q+p)/p}} < \infty, \quad \sup_{z \in \mathbb{D}} \frac{\mu(|z|)|\varphi''(z)|}{(1-|\varphi(z)|^2)^{(2+q)/p}} < \infty. \quad (2.3)$$

*Proof.* Suppose that the conditions in (2.3) hold. Then for any  $z \in \mathbb{D}$  and  $f \in Q_K(p, q)$ ,

$$\begin{aligned} \mu(|z|) \left| (DC_\varphi f)'(z) \right| &= \mu(|z|) \left| (f'(\varphi)\varphi')'(z) \right| \\ &\leq \mu(|z|) |\varphi'(z)|^2 |f''(\varphi(z))| + \mu(|z|) |\varphi''(z)| |f'(\varphi(z))| \\ &\leq \frac{\mu(|z|) |\varphi'(z)|^2}{(1-|\varphi(z)|^2)^{(2+q+p)/p}} \|f\|_{\mathcal{B}^{(q+2)/p}} + \frac{\mu(|z|) |\varphi''(z)|}{(1-|\varphi(z)|^2)^{(2+q)/p}} \|f\|_{\mathcal{B}^{(q+2)/p}} \quad (2.4) \\ &\leq \frac{C\mu(|z|) |\varphi'(z)|^2}{(1-|\varphi(z)|^2)^{(2+q+p)/p}} \|f\|_{Q_K(p,q)} + \frac{C\mu(|z|) |\varphi''(z)|}{(1-|\varphi(z)|^2)^{(2+q)/p}} \|f\|_{Q_K(p,q)}, \end{aligned}$$

where we have used the fact that  $\|f\|_{\mathcal{B}^{(q+2)/p}} \leq \|f\|_{Q_K(p,q)}$ , as well as the following well-known characterization for  $\alpha$ -Bloch functions (see, e.g., [33])

$$\sup_{z \in \mathbb{D}} (1-|z|^2)^\beta |\varphi'(z)| \asymp |\varphi'(0)| + \sup_{z \in \mathbb{D}} (1-|z|^2)^{(1+\beta)} |\varphi''(z)|. \quad (2.5)$$

Taking the supremum in (2.4) for  $z \in \mathbb{D}$ , then employing (2.3) we obtain that  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is bounded.

Conversely, suppose that  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is bounded, that is, there exists a constant  $C$  such that  $\|DC_\varphi f\|_{\mathcal{B}_\mu} \leq C \|f\|_{Q_K(p,q)}$  for all  $f \in Q_K(p, q)$ . Taking the functions  $f(z) \equiv z$ , and  $f(z) \equiv z^2$ , which belong to  $Q_K(p, q)$ , we get

$$\sup_{z \in \mathbb{D}} \mu(|z|) |\varphi''(z)| < \infty, \quad (2.6)$$

$$\sup_{z \in \mathbb{D}} \mu(|z|) \left| (\varphi'(z))^2 + \varphi''(z)\varphi(z) \right| < \infty. \quad (2.7)$$

From (2.6), (2.7), and the boundedness of the function  $\varphi(z)$ , it follows that

$$\sup_{z \in \mathbb{D}} \mu(|z|) |\varphi'(z)|^2 < \infty. \tag{2.8}$$

For  $w \in \mathbb{D}$ , let

$$f_w(z) = \frac{1 - |w|^2}{(1 - z\bar{w})^{(q+2)/p}}. \tag{2.9}$$

By some direct calculation we have that

$$\begin{aligned} f'_w(w) &= \frac{q+2}{p} \frac{\bar{w}}{(1 - |w|^2)^{(q+2)/p}}, \\ f''_w(w) &= \left(\frac{q+2}{p}\right) \left(\frac{q+2}{p} + 1\right) \frac{\bar{w}^2}{(1 - |w|^2)^{(q+2)/p+1}}. \end{aligned} \tag{2.10}$$

From [8], we know that  $f_w \in \mathcal{Q}_K(p, q)$ , for each  $w \in \mathbb{D}$ , moreover there is a positive constant  $C$  such that  $\sup_{w \in \mathbb{D}} \|f_w\|_{\mathcal{Q}_K(p, q)} \leq C$ . Hence, we have

$$\begin{aligned} C \|DC_\varphi\|_{\mathcal{Q}_K(p, q) \rightarrow \mathcal{B}_\mu} &\geq \|DC_\varphi f_{\varphi(\lambda)}\|_{\mathcal{B}_\mu} \\ &\geq \frac{q+2}{p} \frac{q+2+p}{p} \frac{\mu(|\lambda|) |\varphi'(\lambda)|^2 |\varphi(\lambda)|^2}{(1 - |\varphi(\lambda)|^2)^{(2+q+p)/p}} \\ &\quad + \frac{q+2}{p} \frac{\mu(|\lambda|) |\varphi''(\lambda)| |\varphi(\lambda)|}{(1 - |\varphi(\lambda)|^2)^{(2+q)/p}}, \end{aligned} \tag{2.11}$$

for  $\lambda \in \mathbb{D}$ . Therefore, we obtain

$$\frac{\mu(|\lambda|) |\varphi''(\lambda)| |\varphi(\lambda)|}{(1 - |\varphi(\lambda)|^2)^{(2+q)/p}} \leq C \|DC_\varphi\|_{\mathcal{Q}_K(p, q) \rightarrow \mathcal{B}_\mu} + \frac{q+2+p}{p} \frac{\mu(|\lambda|) |\varphi'(\lambda)|^2 |\varphi(\lambda)|^2}{(1 - |\varphi(\lambda)|^2)^{(2+q+p)/p}}. \tag{2.12}$$

Next, for  $w \in \mathbb{D}$ , let

$$g_w(z) = \frac{(1 - |w|^2)^2}{(1 - z\bar{w})^{(q+2)/p+1}} - \frac{(q+2)/p + 1}{(q+2)/p} \frac{1 - |w|^2}{(1 - z\bar{w})^{(q+2)/p}}. \tag{2.13}$$

Then from [8], we see that  $g_w \in Q_K(p, q)$  and  $\sup_{w \in \mathbb{D}} \|g_w\|_{Q_K(p, q)} < \infty$ . Since

$$g'_{\varphi(\lambda)}(\varphi(\lambda)) = 0, \quad \left| g''_{\varphi(\lambda)}(\varphi(\lambda)) \right| = \frac{q+2+p}{p} \frac{|\varphi(\lambda)|^2}{(1-|\varphi(\lambda)|^2)^{(2+q+p)/p}}, \quad (2.14)$$

we have

$$\infty > C \|DC_\varphi\|_{Q_K(p, q) \rightarrow \mathcal{B}_\mu} \geq \|DC_\varphi g_{\varphi(\lambda)}\|_{\mathcal{B}_\mu} \geq \frac{q+2+p}{p} \frac{\mu(|\lambda|) |\varphi'(\lambda)|^2 |\varphi(\lambda)|^2}{(1-|\varphi(\lambda)|^2)^{(2+q+p)/p}}. \quad (2.15)$$

Thus

$$\begin{aligned} \sup_{|\varphi(\lambda)| > 1/2} \frac{\mu(|\lambda|) |\varphi'(\lambda)|^2}{(1-|\varphi(\lambda)|^2)^{(2+q+p)/p}} &\leq \sup_{|\varphi(\lambda)| > 1/2} 4 \frac{\mu(|\lambda|) |\varphi'(\lambda)|^2 |\varphi(\lambda)|^2}{(1-|\varphi(\lambda)|^2)^{(2+q+p)/p}} \\ &\leq C \|DC_\varphi\|_{Q_K(p, q) \rightarrow \mathcal{B}_\mu} < \infty. \end{aligned} \quad (2.16)$$

Inequality (2.8) gives

$$\sup_{|\varphi(\lambda)| \leq 1/2} \frac{\mu(|\lambda|) |\varphi'(\lambda)|^2}{(1-|\varphi(\lambda)|^2)^{(2+q+p)/p}} \leq \frac{4^{(2+q+p)/p}}{3^{(2+q+p)/p}} \sup_{|\varphi(\lambda)| \leq 1/2} \mu(|\lambda|) |\varphi'(\lambda)|^2 < \infty. \quad (2.17)$$

Therefore, the first inequality in (2.3) follows from (2.16) and (2.17). From (2.12) and (2.15), we obtain

$$\sup_{\lambda \in \mathbb{D}} \frac{\mu(|\lambda|) |\varphi''(\lambda)| |\varphi(\lambda)|}{(1-|\varphi(\lambda)|^2)^{(2+q)/p}} < \infty. \quad (2.18)$$

Equations (2.6) and (2.18) imply

$$\sup_{|\varphi(\lambda)| > 1/2} \frac{\mu(|\lambda|) |\varphi''(\lambda)|}{(1-|\varphi(\lambda)|^2)^{(2+q)/p}} \leq 2 \sup_{|\varphi(\lambda)| > 1/2} \frac{\mu(|\lambda|) |\varphi''(\lambda)| |\varphi(\lambda)|}{(1-|\varphi(\lambda)|^2)^{(2+q)/p}} < \infty, \quad (2.19)$$

$$\sup_{|\varphi(\lambda)| \leq 1/2} \frac{\mu(|\lambda|) |\varphi''(\lambda)|}{(1-|\varphi(\lambda)|^2)^{(2+q)/p}} \leq \frac{4^{(2+q)/p}}{3^{(2+q)/p}} \sup_{|\varphi(\lambda)| \leq 1/2} \mu(|\lambda|) |\varphi''(\lambda)| < \infty. \quad (2.20)$$

Inequality (2.19) together with (2.20) implies the second inequality of (2.3). This completes the proof of Theorem 2.3.  $\square$

**Theorem 2.4.** Let  $\varphi$  be an analytic self-map of  $\mathbb{D}$ . Suppose that  $\mu$  is normal,  $p > 0$ ,  $q > -2$  and  $K$  is a nonnegative nondecreasing function on  $[0, \infty)$  such that (2.2) holds. Then  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is compact if and only if  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is bounded,

$$\lim_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|\varphi'(z)|^2}{(1 - |\varphi(z)|^2)^{(2+q+p)/p}} = 0, \quad \lim_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|\varphi''(z)|}{(1 - |\varphi(z)|^2)^{(2+q)/p}} = 0. \quad (2.21)$$

*Proof.* Suppose that  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is bounded and (2.21) holds. Let  $(f_k)_{k \in \mathbb{N}}$  be a sequence in  $Q_K(p, q)$  such that  $\sup_{k \in \mathbb{N}} \|f_k\|_{Q_K(p, q)} < \infty$  and  $f_k$  converges to 0 uniformly on compact subsets of  $\mathbb{D}$  as  $k \rightarrow \infty$ . By the assumption, for any  $\varepsilon > 0$ , there exists a  $\delta \in (0, 1)$  such that

$$\frac{\mu(|z|)|\varphi'(z)|^2}{(1 - |\varphi(z)|^2)^{(2+p+q)/p}} < \varepsilon, \quad \frac{\mu(|z|)|\varphi''(z)|}{(1 - |\varphi(z)|^2)^{(2+q)/p}} < \varepsilon, \quad (2.22)$$

when  $\delta < |\varphi(z)| < 1$ . Since  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is bounded, then from the proof of Theorem 2.3 we have

$$M_1 := \sup_{z \in \mathbb{D}} \mu(|z|)|\varphi''(z)| < \infty, \quad M_2 := \sup_{z \in \mathbb{D}} \mu(|z|)|\varphi'(z)|^2 < \infty. \quad (2.23)$$

Let  $K = \{z \in \mathbb{D} : |\varphi(z)| \leq \delta\}$ . Then, we have

$$\begin{aligned} \|DC_\varphi f_k\|_{\mathcal{B}_\mu} &= \sup_{z \in \mathbb{D}} \mu(|z|) \left| (DC_\varphi f_k)'(z) \right| + |f_k'(\varphi(0))| |\varphi'(0)| \\ &= \sup_{z \in \mathbb{D}} \mu(|z|) \left| (\varphi' f_k'(\varphi))'(z) \right| + |f_k'(\varphi(0))| |\varphi'(0)| \\ &\leq \sup_{z \in \mathbb{D}} \mu(|z|) |\varphi'(z)|^2 |f_k''(\varphi(z))| + \sup_{z \in \mathbb{D}} \mu(|z|) |\varphi''(z)| |f_k'(\varphi(z))| \\ &\quad + |f_k'(\varphi(0))| |\varphi'(0)| \\ &\leq \sup_K \mu(|z|) |\varphi'(z)|^2 |f_k''(\varphi(z))| + \sup_K \mu(|z|) |\varphi''(z)| |f_k'(\varphi(z))| \\ &\quad + \sup_{\mathbb{D} \setminus K} \mu(|z|) |\varphi''(z)|^2 |f_k''(\varphi(z))| + \sup_{\mathbb{D} \setminus K} \mu(|z|) |\varphi''(z)| |f_k'(\varphi(z))| \\ &\quad + |f_k'(\varphi(0))| |\varphi'(0)| \\ &\leq \sup_K \mu(|z|) |\varphi'(z)|^2 |f_k''(\varphi(z))| + \sup_K \mu(|z|) |\varphi''(z)| |f_k'(\varphi(z))| \\ &\quad + |f_k'(\varphi(0))| |\varphi'(0)| + C \sup_{\mathbb{D} \setminus K} \frac{\mu(|z|)|\varphi'(z)|^2}{(1 - |\varphi(z)|^2)^{(2+p+q)/p}} \|f_k\|_{Q_K(p, q)} \end{aligned}$$

$$\begin{aligned}
& + \sup_{\mathbb{D} \setminus K} \frac{\mu(|z|)|\varphi''(z)|}{(1-|\varphi(z)|^2)^{(2+p)/p}} \|f_k\|_{Q_K(p,q)} \\
& \leq M_2 \sup_K |f_k''(\varphi(z))| + M_1 \sup_K |f_k'(\varphi(z))| + 2C\varepsilon \|f_k\|_{Q_K(p,q)} \\
& \quad + |f_k'(\varphi(0))||\varphi'(0)|.
\end{aligned} \tag{2.24}$$

The assumption that  $f_k \rightarrow 0$  as  $k \rightarrow \infty$  on compact subsets of  $\mathbb{D}$  along with Cauchy's estimate give that  $f_k' \rightarrow 0$  and  $f_k'' \rightarrow 0$  as  $k \rightarrow \infty$  on compact subsets of  $\mathbb{D}$ . Letting  $k \rightarrow \infty$  in (2.24) and using the fact that  $\varepsilon$  is an arbitrary positive number, we obtain  $\lim_{k \rightarrow \infty} \|DC_\varphi f_k\|_{\mathcal{B}_\mu} = 0$ . Applying Lemma 2.1, the result follows.

Now, suppose that  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is compact. Then it is clear that  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is bounded. Let  $(z_k)_{k \in \mathbb{N}}$  be a sequence in  $\mathbb{D}$  such that  $|\varphi(z_k)| \rightarrow 1$  as  $k \rightarrow \infty$  (if such a sequence does not exist then condition (2.21) is vacuously satisfied). Let

$$f_k(z) = \frac{1 - |\varphi(z_k)|^2}{(1 - \overline{\varphi(z_k)}z)^{(q+2)/p}}. \tag{2.25}$$

Then,  $\sup_{k \in \mathbb{N}} \|f_k\|_{Q_K(p,q)} < \infty$  and  $f_k$  converges to 0 uniformly on compact subsets of  $\mathbb{D}$  as  $k \rightarrow \infty$ . Since  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is compact, by Lemma 2.1 we have  $\lim_{k \rightarrow \infty} \|DC_\varphi f_k\|_{\mathcal{B}_\mu} = 0$ . On the other hand, from (2.11) we have

$$C \|DC_\varphi f_k\|_{\mathcal{B}_\mu} \geq \left| -\frac{2+q}{p} \frac{2+p+q}{p} \frac{\mu(|z_k|)|\varphi'(z_k)|^2 |\varphi(z_k)|^2}{(1-|\varphi(z_k)|^2)^{(2+q+p)/p}} + \frac{2+q}{p} \frac{\mu(|z_k|)|\varphi''(z_k)| |\varphi(z_k)|}{(1-|\varphi(z_k)|^2)^{(2+q)/p}} \right|, \tag{2.26}$$

which implies that

$$\lim_{|\varphi(z_k)| \rightarrow 1} \frac{2+q+p}{p} \frac{\mu(|z_k|)|\varphi'(z_k)|^2 |\varphi(z_k)|^2}{(1-|\varphi(z_k)|^2)^{(2+q+p)/p}} = \lim_{|\varphi(z_k)| \rightarrow 1} \frac{\mu(|z_k|)|\varphi''(z_k)| |\varphi(z_k)|}{(1-|\varphi(z_k)|^2)^{(2+q)/p}}, \tag{2.27}$$

if one of these two limits exists.

Next, for  $k \in \mathbb{N}$ , set

$$g_k(z) = \frac{(1-|\varphi(z_k)|^2)^2}{(1-\overline{\varphi(z_k)}z)^{(q+2)/p+1}} - \frac{q+2+p}{q+2} \frac{1-|\varphi(z_k)|^2}{(1-\overline{\varphi(z_k)}z)^{(q+2)/p}}. \tag{2.28}$$

Then  $(g_k)_{k \in \mathbb{N}}$  is a sequence in  $Q_K(p, q)$ . Notice that  $g'_k(\varphi(z_k)) = 0$ ,

$$|g''_k(\varphi(z_k))| = \frac{2 + q + p}{p} \frac{|\varphi(z_k)|^2}{(1 - |\varphi(z_k)|^2)^{(2+q+p)/p}}, \quad (2.29)$$

and  $g_k$  converges to 0 uniformly on compact subsets of  $\mathbb{D}$  as  $k \rightarrow \infty$ . Since  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is compact, we have  $\lim_{k \rightarrow \infty} \|DC_\varphi g_k\|_{\mathcal{B}_\mu} = 0$ . On the other hand, we have

$$\frac{2 + q + p}{p} \frac{\mu(|z_k|) |\varphi'(z_k)|^2 |\varphi(z_k)|^2}{(1 - |\varphi(z_k)|^2)^{(2+q+p)/p}} \leq \|DC_\varphi g_k\|_{\mathcal{B}_\mu}. \quad (2.30)$$

Therefore

$$\lim_{|\varphi(z_k)| \rightarrow 1} \frac{\mu(|z_k|) |\varphi'(z_k)|^2}{(1 - |\varphi(z_k)|^2)^{(2+q+p)/p}} = \lim_{|\varphi(z_k)| \rightarrow 1} \frac{\mu(|z_k|) |\varphi'(z_k)|^2 |\varphi(z_k)|^2}{(1 - |\varphi(z_k)|^2)^{(2+q+p)/p}} = 0. \quad (2.31)$$

This along with (2.27) implies

$$\lim_{|\varphi(z_k)| \rightarrow 1} \frac{\mu(|z_k|) |\varphi''(z_k)|}{(1 - |\varphi(z_k)|^2)^{(2+p)/p}} = 0. \quad (2.32)$$

From the last two equalities, the desired result follows.  $\square$

**Theorem 2.5.** *Let  $\varphi$  be an analytic self-map of  $\mathbb{D}$ . Suppose that  $\mu$  is normal,  $p > 0$ ,  $q > -2$  and  $K$  is a nonnegative nondecreasing function on  $[0, \infty)$  such that (2.2) holds. Then  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_{\mu,0}$  is compact if and only if*

$$\lim_{|z| \rightarrow 1} \frac{\mu(|z|) |\varphi'(z)|^2}{(1 - |\varphi(z)|^2)^{(2+q+p)/p}} = 0, \quad \lim_{|z| \rightarrow 1} \frac{\mu(|z|) |\varphi''(z)|}{(1 - |\varphi(z)|^2)^{(2+q)/p}} = 0. \quad (2.33)$$

*Proof.* Sufficiency. Let  $f \in Q_K(p, q)$ . By the proof of Theorem 2.3 we have

$$\mu(|z|) \left| (DC_\varphi f)'(z) \right| \leq C \left( \frac{\mu(|z|) |\varphi'(z)|^2}{(1 - |\varphi(z)|^2)^{(2+q+p)/p}} + \frac{\mu(|z|) |\varphi''(z)|}{(1 - |\varphi(z)|^2)^{(2+q)/p}} \right) \|f\|_{Q_K(p,q)}. \quad (2.34)$$

Taking the supremum in (2.34) over all  $f \in Q_K(p, q)$  such that  $\|f\|_{Q_K(p, q)} \leq 1$ , then letting  $|z| \rightarrow 1$ , we get

$$\lim_{|z| \rightarrow 1} \sup_{\|f\|_{Q_K(p, q)} \leq 1} \mu(|z|) \left| (DC_\varphi f)'(z) \right| = 0. \quad (2.35)$$

From which by Lemma 2.2 we see that the operator  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_{\mu, 0}$  is compact. Necessity. Assume that  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_{\mu, 0}$  is compact. By taking the function given by  $f(z) \equiv z$  and using the boundedness of  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_{\mu, 0}$ , we get

$$\lim_{|z| \rightarrow 1} \mu(|z|) |\varphi''(z)| = 0. \quad (2.36)$$

From this, by taking the test function  $f(z) \equiv z^2$  and using the boundedness of  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_{\mu, 0}$  it follows that

$$\lim_{|z| \rightarrow 1} \mu(|z|) |\varphi'(z)|^2 = 0. \quad (2.37)$$

If  $\|\varphi\|_\infty < 1$ , from (2.36) and (2.37), we obtain that

$$\begin{aligned} \lim_{|z| \rightarrow 1} \frac{\mu(|z|) |\varphi'(z)|^2}{(1 - |\varphi(z)|^2)^{(2+q+p)/p}} &\leq \frac{1}{(1 - \|\varphi\|_\infty^2)^{(2+q+p)/p}} \lim_{|z| \rightarrow 1} \mu(|z|) |\varphi'(z)|^2 = 0, \\ \lim_{|z| \rightarrow 1} \frac{\mu(|z|) |\varphi''(z)|}{(1 - |\varphi(z)|^2)^{(2+q)/p}} &\leq \frac{1}{(1 - \|\varphi\|_\infty^2)^{(2+q)/p}} \lim_{|z| \rightarrow 1} \mu(|z|) |\varphi''(z)| = 0, \end{aligned} \quad (2.38)$$

from which the result follows in this case.

Assume that  $\|\varphi\|_\infty = 1$ . Let  $(\varphi(z_k))_{k \in \mathbb{N}}$  be a sequence such that  $\lim_{k \rightarrow \infty} |\varphi(z_k)| = 1$ . From the compactness of  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_{\mu, 0}$  we see that  $DC_\varphi : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is compact. From Theorem 2.4 we get

$$\lim_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|) |\varphi'(z)|^2}{(1 - |\varphi(z)|^2)^{(2+p+q)/p}} = 0, \quad (2.39)$$

$$\lim_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|) |\varphi''(z)|}{(1 - |\varphi(z)|^2)^{(2+q)/p}} = 0. \quad (2.40)$$

From (2.36) and (2.40), we have that for every  $\varepsilon > 0$ , there exists an  $r \in (0, 1)$  such that

$$\frac{\mu(|z|)|\varphi''(z)|}{(1 - |\varphi(z)|^2)^{(2+q)/p}} < \varepsilon, \tag{2.41}$$

when  $r < |\varphi(z)| < 1$ , and there exists a  $\sigma \in (0, 1)$  such that  $\mu(|z|)|\varphi''(z)| \leq \varepsilon(1 - r^2)^{(2+q)/p}$  when  $\sigma < |z| < 1$ . Therefore, when  $\sigma < |z| < 1$ , and  $r < |\varphi(z)| < 1$ , we have

$$\frac{\mu(|z|)|\varphi''(z)|}{(1 - |\varphi(z)|^2)^{(2+q)/p}} < \varepsilon. \tag{2.42}$$

On the other hand, if  $\sigma < |z| < 1$ , and  $|\varphi(z)| \leq r$ , we obtain

$$\frac{\mu(|z|)|\varphi''(z)|}{(1 - |\varphi(z)|^2)^{(2+q)/p}} < \frac{1}{(1 - r^2)^{(2+q)/p}} \mu(|z|)|\varphi''(z)| < \varepsilon. \tag{2.43}$$

Inequality (2.42) together with (2.43) gives the second equality of (2.33). Similarly to the above arguments, by (2.37) and (2.39) we get the first equality of (2.33). The proof is completed.  $\square$

From the above three theorems, we get the following corollary (see [14]).

**Corollary 2.6.** *Let  $\varphi$  be an analytic self-map of  $\mathbb{D}$ . Then the following statements hold.*

(i)  $DC_\varphi : \mathcal{B} \rightarrow \mathcal{B}$  is bounded if and only if

$$\sup_{z \in \mathbb{D}} \frac{(1 - |z|^2)|\varphi'(z)|^2}{(1 - |\varphi(z)|^2)^2} < \infty, \quad \sup_{z \in \mathbb{D}} \frac{(1 - |z|^2)|\varphi''(z)|}{1 - |\varphi(z)|^2} < \infty; \tag{2.44}$$

(ii)  $DC_\varphi : \mathcal{B} \rightarrow \mathcal{B}$  is compact if and only if  $DC_\varphi : \mathcal{B} \rightarrow \mathcal{B}$  is bounded,

$$\lim_{|\varphi(z)| \rightarrow 1} \frac{(1 - |z|^2)|\varphi'(z)|^2}{(1 - |\varphi(z)|^2)^2} = 0, \quad \lim_{|\varphi(z)| \rightarrow 1} \frac{(1 - |z|^2)|\varphi''(z)|}{1 - |\varphi(z)|^2} = 0; \tag{2.45}$$

(iii)  $DC_\varphi : \mathcal{B} \rightarrow \mathcal{B}_0$  is compact if and only if

$$\lim_{|z| \rightarrow 1} \frac{(1 - |z|^2)|\varphi'(z)|^2}{(1 - |\varphi(z)|^2)^2} = 0, \quad \lim_{|z| \rightarrow 1} \frac{(1 - |z|^2)|\varphi''(z)|}{1 - |\varphi(z)|^2} = 0. \tag{2.46}$$

Similarly to the proofs of Theorems 2.3–2.5, we can get the following result. We omit the proof.

**Theorem 2.7.** *Let  $\varphi$  be an analytic self-map of  $\mathbb{D}$ . Suppose that  $\mu$  is normal,  $p > 0$ ,  $q > -2$  and  $K$  is a nonnegative nondecreasing function on  $[0, \infty)$  such that (2.2) holds. Then the following statements hold.*

(i)  $C_\varphi D : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is bounded if and only if

$$\sup_{z \in \mathbb{D}} \frac{\mu(|z|)|\varphi'(z)|}{(1 - |\varphi(z)|^2)^{(2+p+q)/p}} < \infty; \quad (2.47)$$

(ii)  $C_\varphi D : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is compact if and only if  $C_\varphi D : Q_K(p, q) \rightarrow \mathcal{B}_\mu$  is bounded and

$$\lim_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|\varphi'(z)|}{(1 - |\varphi(z)|^2)^{(2+p+q)/p}} = 0; \quad (2.48)$$

(iii)  $C_\varphi D : Q_K(p, q) \rightarrow \mathcal{B}_{\mu,0}$  is compact if and only if

$$\lim_{|z| \rightarrow 1} \frac{\mu(|z|)|\varphi'(z)|}{(1 - |\varphi(z)|^2)^{(2+p+q)/p}} = 0. \quad (2.49)$$

From Theorem 2.7 we get the following corollary.

**Corollary 2.8.** *Let  $\varphi$  be an analytic self-map of  $\mathbb{D}$ . Then the following statements hold.*

(i)  $C_\varphi D : \mathcal{B} \rightarrow \mathcal{B}$  is bounded if and only if

$$\sup_{z \in \mathbb{D}} \frac{(1 - |z|^2)|\varphi'(z)|}{(1 - |\varphi(z)|^2)^2} < \infty; \quad (2.50)$$

(ii)  $C_\varphi D : \mathcal{B} \rightarrow \mathcal{B}$  is compact if and only if  $C_\varphi D : \mathcal{B} \rightarrow \mathcal{B}$  is bounded and

$$\lim_{|\varphi(z)| \rightarrow 1} \frac{(1 - |z|^2)|\varphi'(z)|}{(1 - |\varphi(z)|^2)^2} = 0; \quad (2.51)$$

(iii)  $C_\varphi D : \mathcal{B} \rightarrow \mathcal{B}_0$  is compact if and only if

$$\lim_{|z| \rightarrow 1} \frac{(1 - |z|^2)|\varphi'(z)|}{(1 - |\varphi(z)|^2)^2} = 0. \quad (2.52)$$

## References

- [1] A. L. Shields and D. L. Williams, "Bonded projections, duality, and multipliers in spaces of analytic functions," *Transactions of the American Mathematical Society*, vol. 162, pp. 287–302, 1971.
- [2] X. Fu and X. Zhu, "Weighted composition operators on some weighted spaces in the unit ball," *Abstract and Applied Analysis*, vol. 2008, Article ID 605807, 8 pages, 2008.
- [3] Z. Hu and S. Wang, "Composition operators on Bloch-type spaces," *Proceedings of the Royal Society of Edinburgh A*, vol. 135, no. 6, pp. 1229–1239, 2005.
- [4] S. Stević, "Norm of weighted composition operators from Bloch space to  $H_\mu^\infty$  on the unit ball," *Ars Combinatoria*, vol. 88, pp. 125–127, 2008.
- [5] S. Stević, "On a new operator from  $H^\infty$  to the Bloch-type space on the unit ball," *Utilitas Mathematica*, vol. 77, pp. 257–263, 2008.
- [6] H. Wulan and J. Zhou, " $Q_K$  type spaces of analytic functions," *Journal of Function Spaces and Applications*, vol. 4, no. 1, pp. 73–84, 2006.
- [7] R. Zhao, "On a general family of function spaces," *Annales Academiæ Scientiarum Fennicæ. Mathematica Dissertationes*, no. 105, pp. 1–56, 1996.
- [8] M. Kotilainen, "On composition operators in  $Q_K$  type spaces," *Journal of Function Spaces and Applications*, vol. 5, no. 2, pp. 103–122, 2007.
- [9] C. C. Cowen and B. D. MacCluer, *Composition Operators on Spaces of Analytic Functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, Fla, USA, 1995.
- [10] S. Stević, "Weighted composition operators from weighted Bergman spaces to weighted-type spaces on the unit ball," *Applied Mathematics and Computation*, vol. 212, no. 2, pp. 499–504, 2009.
- [11] R. A. Hibschweiler and N. Portnoy, "Composition followed by differentiation between Bergman and Hardy spaces," *The Rocky Mountain Journal of Mathematics*, vol. 35, no. 3, pp. 843–855, 2005.
- [12] S. Ohno, "Products of composition and differentiation between Hardy spaces," *Bulletin of the Australian Mathematical Society*, vol. 73, no. 2, pp. 235–243, 2006.
- [13] S. Stević, "Products of composition and differentiation operators on the weighted Bergman space," to appear in *Bulletin of the Belgian Mathematical Society—Simon Stevin*.
- [14] S. Li and S. Stević, "Composition followed by differentiation between Bloch type spaces," *Journal of Computational Analysis and Applications*, vol. 9, no. 2, pp. 195–205, 2007.
- [15] S. Li and S. Stević, "Composition followed by differentiation between  $H^\infty$  and  $\alpha$ -Bloch spaces," *Houston Journal of Mathematics*, vol. 35, no. 1, pp. 327–340, 2009.
- [16] S. Li and S. Stević, "Composition followed by differentiation from mixed-norm spaces to  $\alpha$ -Bloch spaces," *Sbornik Mathematics*, vol. 199, no. 12, pp. 1847–1857, 2008.
- [17] S. Stević, "Norm and essential norm of composition followed by differentiation from  $\alpha$ -Bloch spaces to  $H_\mu^\infty$ ," *Applied Mathematics and Computation*, vol. 207, no. 1, pp. 225–229, 2009.
- [18] S. Stević, "Generalized composition operators between mixed-norm and some weighted spaces," *Numerical Functional Analysis and Optimization*, vol. 29, no. 7-8, pp. 959–978, 2008.
- [19] S. Stević, "Weighted differentiation composition operators from mixed-norm spaces to weighted-type spaces," *Applied Mathematics and Computation*, vol. 211, no. 1, pp. 222–233, 2009.
- [20] X. Zhu, "Generalized weighted composition operators from Bloch type spaces to weighted Bergman spaces," *Indian Journal of Mathematics*, vol. 49, no. 2, pp. 139–150, 2007.
- [21] X. Zhu, "Products of differentiation, composition and multiplication from Bergman type spaces to Bers type spaces," *Integral Transforms and Special Functions*, vol. 18, no. 3, pp. 223–231, 2007.
- [22] S. Li and S. Stević, "Generalized composition operators on Zygmund spaces and Bloch type spaces," *Journal of Mathematical Analysis and Applications*, vol. 338, no. 2, pp. 1282–1295, 2008.
- [23] S. Li and S. Stević, "Products of composition and integral type operators from  $H^\infty$  to the Bloch space," *Complex Variables and Elliptic Equations*, vol. 53, no. 5, pp. 463–474, 2008.
- [24] S. Li and S. Stević, "Products of Volterra type operator and composition operator from  $H^\infty$  and Bloch spaces to Zygmund spaces," *Journal of Mathematical Analysis and Applications*, vol. 345, no. 1, pp. 40–52, 2008.
- [25] S. Li and S. Stević, "Products of integral-type operators and composition operators between Bloch-type spaces," *Journal of Mathematical Analysis and Applications*, vol. 349, no. 2, pp. 596–610, 2009.

- [26] S. Stević, "Generalized composition operators from logarithmic Bloch spaces to mixed-norm spaces," *Utilitas Mathematica*, vol. 77, pp. 167–172, 2008.
- [27] S. Stević, "On a new integral-type operator from the weighted Bergman space to the Bloch-type space on the unit ball," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 154263, 14 pages, 2008.
- [28] S. Stević, "On a new operator from the logarithmic Bloch space to the Bloch-type space on the unit ball," *Applied Mathematics and Computation*, vol. 206, no. 1, pp. 313–320, 2008.
- [29] S. Stević, "On a new integral-type operator from the Bloch space to Bloch-type spaces on the unit ball," *Journal of Mathematical Analysis and Applications*, vol. 354, no. 2, pp. 426–434, 2009.
- [30] S. Stević, "On an integral operator between Bloch-type spaces on the unit ball," *Bulletin des Sciences Mathématiques*. In press.
- [31] S. Stević, "Composition operators between  $H^\infty$  and  $a$ -Bloch spaces on the polydisc," *Zeitschrift für Analysis und ihre Anwendungen*, vol. 25, no. 4, pp. 457–466, 2006.
- [32] K. Madigan and A. Matheson, "Compact composition operators on the Bloch space," *Transactions of the American Mathematical Society*, vol. 347, no. 7, pp. 2679–2687, 1995.
- [33] K. H. Zhu, "Bloch type spaces of analytic functions," *The Rocky Mountain Journal of Mathematics*, vol. 23, no. 3, pp. 1143–1177, 1993.