

Algebraicity of holomorphic mappings between real algebraic sets in $\mathbf{C}^n$

by

M. S. BAOUENDI

*University of California, San Diego
La Jolla, CA, U.S.A.*

P. EBENFELT

*Royal Institute of Technology
Stockholm, Sweden*

and

L. P. ROTHCHILD

*University of California, San Diego
La Jolla, CA, U.S.A.*

Contents

- 0. Introduction
- 1. Holomorphic nondegeneracy of real-analytic manifolds
 - 1.1. Preliminaries on real submanifolds of $\mathbf{C}^N$
 - 1.2. Holomorphic nondegeneracy and its propagation
 - 1.3. The Levi number and essential finiteness
 - 1.4. Holomorphic nondegeneracy of real algebraic sets
- 2. The Segre sets of a real-analytic CR submanifold
 - 2.1. Complexification of M , involution, and projections
 - 2.2. Definition of the Segre sets of M at p_0
 - 2.3. Homogeneous submanifolds of CR dimension 1
 - 2.4. Homogeneous submanifolds of arbitrary CR dimension
 - 2.5. Proof of Theorem 2.2.1
- 3. Algebraic properties of holomorphic mappings between real algebraic sets
 - 3.1. A generalization of Theorems 1 and 4
 - 3.2. Propagation of algebraicity
 - 3.3. Proof of Theorem 3.1.2
 - 3.4. Proof of Theorem 3.1.8
 - 3.5. An example
 - 3.6. Proofs of Theorems 1 through 4

The first and third authors were partially supported by National Science Foundation Grant DMS 95-01516. The second author was supported by a grant from the Swedish Natural Science Research Council and would like to thank the University of California, San Diego, for its hospitality during the preparation of this paper.