

# CORRECTION TO “PAIR CORRELATION DENSITIES OF INHOMOGENEOUS QUADRATIC FORMS, II”

---

JENS MARKLOF

The set  $C \subset \mathbb{T}^k$  in [1, Th. 1.7] is wrongly characterized as a set of second Baire category since we explicitly restrict  $C$  to Diophantine vectors of type  $(k-1)/(k-2)$ .  $C$  is instead only a dense subset in  $\mathbb{T}^k$ . The correct statement of [1, Th. 1.7] is as follows.

## THEOREM 1.7

Let  $k > 2$ . For any  $a > 0$ , there exists a dense set  $C \subset \mathbb{T}^k$  for which the following hold.

- (i) All  $\alpha \in C$  are Diophantine of type  $\kappa = (k-1)/(k-2)$ , and the components of the vector  $(\alpha, 1) \in \mathbb{R}^{k+1}$  are linearly independent over  $\mathbb{Q}$ .
- (ii) For  $\alpha \in C$ , we find arbitrarily large  $X$  such that

$$R_2[-a, a](X) \geq \frac{\log X}{\log \log \log X}.$$

- (iii) For  $\alpha \in C$ , there exists an infinite sequence  $L_1 < L_2 < \dots \rightarrow \infty$  such that

$$\lim_{j \rightarrow \infty} R_2[-a, a](L_j) = 2\pi a.$$

## Proof

One follows the argument in [1, pp. 432–433] (cf. also [2, Sec. 9]) to show that for each fixed badly approximable  $(k-2)$ -tuple  $(\alpha_1, \dots, \alpha_{k-2})$  there is a set of second Baire category of  $(\alpha_{k-1}, \alpha_k) \in \mathbb{T}^2$  such that conditions (i), (ii), and (iii) of Theorem 1.7 hold for  $\alpha = (\alpha_1, \dots, \alpha_k)$ . Because the set of badly approximable  $(k-2)$ -tuples is dense in  $\mathbb{T}^{k-2}$ , and the set of second Baire category of  $(\alpha_{k-1}, \alpha_k)$  is dense in  $\mathbb{T}^2$ , the set  $C$  of  $\alpha$  satisfying conditions (i), (ii), and (iii) of Theorem 1.7 is dense in  $\mathbb{T}^k$ .  $\square$

DUKE MATHEMATICAL JOURNAL

Vol. 120, No. 1, © 2003

Received 19 June 2003.

2000 *Mathematics Subject Classification*. Primary 11P21; Secondary 11F27, 22E40, 58F11.