

88. On Non-prolongable Riemann Surfaces.

By Masatsugu TSUJI.

Mathematical Institute, Tokyo Imperial University.

(Comm. by T. YOSIE, M.I.A., Oct. 12, 1943.)

Let F be a Riemann surface spread over the w -plane. T. Radó¹⁾ called F prolongable, if we can map F on a proper part \mathfrak{F}_0 of another Riemann surface \mathfrak{F} spread over the z -plane and non-prolongable, if otherwise. A closed Riemann surface is evidently non-prolongable, but Radó proved that there exists an open non-prolongable Riemann surface by an example of a Riemann surface, which consists of two sheets, whose branch points lie at points $w=n$ ($n=0, \pm 1, \pm 2, \dots$). We will give a class of open non-prolongable Riemann surfaces, which contains the above Radó's example as a special case.

Theorem. Let F be a Riemann surface spread over the w -plane, which consists of n sheets and whose every boundary point is a cluster point of branch points and such that the set E of projections of boundary points of F on the w -plane is a closed set of capacity²⁾ zero. Then F is non-prolongable.

Proof. Suppose that F is prolongable and that we can map F by $w=f(z)$ on a proper part \mathfrak{F}_0 of another Riemann surface \mathfrak{F} spread over the z -plane. Then there exists an inner point z_0 of \mathfrak{F} , which is a boundary point of \mathfrak{F}_0 . We may assume that z_0 is not a branch point of \mathfrak{F} , since otherwise, we can map \mathfrak{F} by $(z-z_0)^{1/n}$ on another Riemann surface, such that z_0 corresponds to an inner point differing from the branch point.

From the definition of E , there exists at least one boundary point of F above any point P of E , but there may exist inner points of F above P .

We take off all points from \mathfrak{F}_0 , which are the images of inner points of F lying above E and the remaining part of \mathfrak{F}_0 be denoted by \mathfrak{F}'_0 . We take ρ so small that all points of a disc: $|z-z_0| \leq \rho$ are inner points of \mathfrak{F} differing from branch points.

Let $\mathfrak{F}_0(\rho)$, $\mathfrak{F}'_0(\rho)$ be the part of \mathfrak{F}_0 , \mathfrak{F}'_0 inside a circle $C: |z-z_0| = \rho$ and $e_0(\rho)$, $e'_0(\rho)$ be the sets of boundary points of $\mathfrak{F}_0(\rho)$, $\mathfrak{F}'_0(\rho)$ inside C respectively.

Since $\text{cap. } E=0$, we see that $e'_0(\rho)$ differs from $e_0(\rho)$ only by a set of capacity zero. Now in $\mathfrak{F}'_0(\rho)$, $f(z)$ does not take values belonging to E and since F consists of only n sheets, if z tends to $e'_0(\rho)$, then $w=f(z)$ has cluster points belonging to E . Hence by a lemma³⁾ proved before, we have $\text{cap. } e'_0(\rho)=0$ and hence $\text{cap. } e_0(\rho)=0$. $w=f(z)$ is one-

1) T. Radó: Über eine nicht fortsetzbare Riemannsche Mannigfaltigkeit. Math. Zeits. **20** (1924).

2) In this paper, "capacity" means "logarithmic capacity".

3) M. Tsuji: On the Riemann surface of an inverse function of a meromorphic function in the neighbourhood of a closed set of capacity zero. Proc. **19** (1943).