

## 122. Note on Mr. Tsuji's Theorem.

By Satoru TAKENAKA.

Shiomi Institute, Osaka.

(Rec. Sep. 5, 1926. Comm. by M. FUJIWARA, M.I.A., Oct. 12, 1926.)

In these Proceedings, **2** (1926), 245, Mr. TSUJI proved an interesting theorem concerning the zero points of a bounded analytic function. Analogous theorems can be established for certain classes of non-bounded functions by similar method.

1. First let  $f(z)$  be regular and analytic for  $|z| < 1$ , and suppose that

$$f(0) = 1, \quad \text{and} \quad \left\{ \frac{1}{2\pi} \int_0^{2\pi} |f(e^{i\theta})|^p d\theta \right\}^{\frac{1}{p}} \leq M_p, \quad (p: \text{real}).$$

We call such a function  $f(z)$  a function of the class  $M_p$ .

If we put

$$r_n^{(p)} = \frac{1}{\sqrt[n]{M_p}},$$

we can prove that

(i) *Every function of the class  $M_p$  has at most  $n-1$  roots in the circle  $|z| < r_n^{(p)}$ .*

(ii) *Among the functions of the class  $M_p$  there exists a function which has just  $n$  roots in the circle  $|z| \leq r_n^{(p)}$ . This function must be of the form*

$$f(z) = \frac{1}{a_1 a_2 \cdots a_n} \cdot \frac{a_1 - z}{1 - \bar{a}_1 z} \cdot \frac{a_2 - z}{1 - \bar{a}_2 z} \cdots \frac{a_n - z}{1 - \bar{a}_n z},$$

where

$$|a_\nu| = \frac{1}{\sqrt[n]{M_p}}, \quad (\nu = 1, 2, \dots, n).$$

These properties can be proved if we use the inequality<sup>1)</sup>

$$|f(z)| \leq M_p \frac{1}{\{1 - |z|^2\}^{\frac{1}{p}}} \prod_{\nu=1}^n \left| \frac{a_\nu - z}{1 - \bar{a}_\nu z} \right|$$

instead of JENJEN's in TSUJI's paper, where  $a_\nu$  ( $\nu = 1, 2, \dots, n$ ) are the roots of  $f(z)$  in  $|z| < 1$  in ascending order of absolute values.

1) S. TAKENAKA, On the power series whose values are given at given points, Japanese Journal of Mathematics, **2** (1925), 81.