

## 16. Note on the Archimedean Property in an Ordered Semigroup

By Tôru SAITÔ

Tokyo Gakugei University

(Comm. by Kinjirô KUNUGI, M. J. A., Jan. 12, 1970)

By an *ordered semigroup* we mean a semigroup with a simple order which is compatible with the semigroup operation. In this note we denote by  $S$  an ordered semigroup. An element  $x$  of  $S$  is called *positive* if  $x < x^2$ , and is called *negative* if  $x^2 < x$ . For an element  $x$  of  $S$ , the number of distinct natural powers of  $x$  is called the *order* of  $x$ .

In [3], we studied some properties of the archimedean equivalence in an ordered semigroup in which every element is non-negative. In this note, we define the archimedean equivalence  $\mathcal{A}$  in a general ordered semigroup and show that similar results hold in this general case.

**Definition.** *The archimedean equivalence  $\mathcal{A}$  on  $S$  is defined by: for  $x, y \in S$ ,  $x \mathcal{A} y$  if and only if there exist natural numbers  $p, q, r$  and  $s$  such that  $x^p \leq y^q$  and  $y^r \leq x^s$ .*

**Theorem 1.** *The archimedean equivalence  $\mathcal{A}$  on  $S$  is an equivalence relation on  $S$ . Each  $\mathcal{A}$ -class is a convex subsemigroup of  $S$ .*

**Lemma 2.** *Each  $\mathcal{A}$ -class contains at most one idempotent.*

**Theorem 3.** *For an  $\mathcal{A}$ -class  $C$ , the following conditions are equivalent:*

- (1)  $C$  contains an idempotent;
- (2) the set of all nonnegative elements of  $C$  is nonempty and has the greatest element;
- (3) the set of all nonpositive elements of  $C$  is nonempty and has the least element;
- (4)  $C$  has the zero element;
- (5) every element of  $C$  is an element of finite order;
- (6)  $C$  contains an element of finite order;
- (7)  $C$  contains at least one nonnegative and at least one nonpositive element.

Moreover, under these conditions, an idempotent of  $C$  is the greatest nonnegative element, the least nonpositive element and also the zero element of  $C$ .

**Corollary 4.** *Let  $x$  be a nonnegative element and  $y$  be an element of an  $\mathcal{A}$ -class  $C$  of  $S$ . Then*

- (1)  $y \leq xy$  if and only if  $y$  is nonnegative;
- (2)  $y \leq yx$  if and only if  $y$  is nonnegative.