

## 167. Monotone Sequence of 0-dimensional Subsets of Metric Spaces

By Keiô NAGAMI

(Comm. by Kinjirô KUNUGI, M.J.A., Nov. 12, 1965)

Let  $\omega_1$  be the first uncountable ordinal and  $\omega(c)$  the first ordinal whose power is  $c$ . This paper proves the following two theorems.

**Theorem 1.** *Let  $X$  be a metric space which is the countable sum of 0-dimensional subsets. Then there exists a sequence  $\{J_i : i < \omega\}$  of subsets of  $X$  such that i)  $J_i \subset J_{i+1}$  for every  $i$ , ii)  $\dim J_i \leq 0$  for every  $i$ , and iii)  $\cup J_i = X$ .*

**Theorem 2.** *Let  $X$  be a non-empty metric space. Then there exists a transfinite sequence  $\{J_\alpha : \alpha < \omega_1\}$  of subsets of  $X$  such that i)  $J_\alpha \subset J_\beta$  whenever  $\alpha < \beta$ , ii)  $\dim J_\alpha \leq 0$  for every  $\alpha$ , and iii)  $\cup J_\alpha = X$ .*

In both cases we use the following notations, where  $\rho$  is the preassigned metric on  $X$ . Take two sequences  $\mathfrak{U}_{ij} = \{U_\lambda : \lambda \in A_{ij}\}$  and  $\mathfrak{F}_{ij} = \{F_\lambda : \lambda \in A_{ij}\}$ , where  $i, j = 1, 2, \dots$ , which satisfy the following conditions (cf. Bing [1]):

- (1)  $\mathfrak{U}_{ij}$  is a discrete collection of open sets of  $X$ .
- (2)  $\mathfrak{F}_{ij}$  is a collection of non-empty closed sets of  $X$ .
- (3)  $F_\lambda \subset U_\lambda$  for every  $\lambda \in A$ , where  $A = \cup A_{ij}$ .
- (4)  $\mathfrak{F}_i = \{F_\lambda : \lambda \in A_i\}$  covers  $X$  for every  $i$ , where  $A_i = \cup_j A_{ij}$ .
- (5)  $\mathfrak{U}_i = \{U_\lambda : \lambda \in A_i\}$  is locally finite.
- (6)  $\rho(\mathfrak{U}_i) < 1/i$ .

Set  $U_{ij} = \cup \{U_\lambda : \lambda \in A_{ij}\}$  and  $F_{ij} = \cup \{F_\lambda : \lambda \in A_{ij}\}$ .

*Proof of Theorem 1.* Let  $I'$  be the set of all rational numbers  $r$  with  $0 < r < 1$ . By Nagata [4, Lemma 4.1] there exists a collection  $\{U_{ijr} : i, j = 1, 2, \dots, r \in I'\}$  of open sets of  $X$  which satisfies the following conditions:

- (7)  $F_{ij} \subset U_{ijr} \subset \bar{U}_{ijr} \subset U_{ijs} \subset \bar{U}_{ijs} \subset U_{ij}$  for  $r < s$ .
- (8)  $\{B(U_{ijr}) = \bar{U}_{ijr} - U_{ijr} : i, j = 1, 2, \dots, r \in I'\}$  is pointfinite.

Let  $I' = \{r_1, r_2, \dots\}$  and  $J_i = X - \cup \{B(U_{jkr}) : j, k = 1, 2, \dots, r \in I' - \{r_1, \dots, r_i\}\}$ . Then by Morita [3, Lemma 3.3]  $\dim J_i \leq 0$ . It is evident that  $J_1 \subset J_2 \subset \dots$ . To see  $\cup J_i = X$  let  $x$  be an arbitrary point of  $X$ . By (8) there exists  $i$  such that  $x \notin B(U_{ijr})$  for any  $i, j$  and any  $r \in I' - \{r_1, \dots, r_i\}$ . Hence  $x \in J_i$  and the proof is completed.

*Proof of Theorem 2.* Let  $I$  be the unit interval  $[0, 1]$  and  $\{I_\alpha : \alpha < \omega(c)\}$  the family of all residue classes of  $I$  modulo the rational numbers. Set

$$L_\alpha = \cup \{I_\beta : \beta \leq \alpha \text{ or } \omega_1 \leq \beta < \omega(c)\}, \quad \alpha < \omega_1.$$