

44. A Counter Example of Ampleness of Positive Line Bundles

By Takeo OHSAWA

Research Institute for Mathematical Sciences, Kyoto University

(Communicated by Kunihiko KODAIRA, M. J. A., May 12, 1979)

0. Let X be a weakly 1-complete manifold and B be a positive line bundle over X . If X is compact or Stein it is well known that B is ample, i.e., there exist an integer ν and sections $s_0, \dots, s_N \in \Gamma(X, B^\nu)$ such that $(s_0 : \dots : s_N)$ gives a holomorphic embedding of X into P^N . The purpose of this note is to show that B is not ample in general. For the definitions of weakly 1-complete manifold and positive line bundles, see [2]. The author would like to thank Dr. S. Iitaka for valuable criticisms.

1. Let p and q be relatively prime positive integers. Let $V_{p,q}$ be a surface in C^3 defined by $z_1^p + z_2^q + z_3^{pq} = 0$. It is shown by Ono and Watanabe [3], that the singularity of $V_{p,q}$ is resolved by a manifold $M_{p,q}$ with exceptional set $S_{p,q}$ which is a non-singular curve of genus $\frac{1}{2}(p-1)(q-1)$ and self-intersection number -1 .

We use the following elementary

Lemma. *For any $c > 0$ there exists $a_0 > c$ such that if $a > a_0$ and $x > a$,*

$$(x-a)^a < (a-c)^c \exp(6(a-c)x - 3(a^2 - c^2)).$$

Let p_i ($i=1, 2, \dots$) be an increasing sequence of prime integers greater than 3 such that every p_{i+1} satisfies the property of a_0 if we take p_i for c in the lemma. This condition implies that any two of the functions

$$x_1^3 \exp(-3(x_2 - p_i)^2) + (x_2 - p_i)^{p_i}, \quad i=1, 2, \dots$$

have no common zeroes on R^2 . Let V' be a real analytic subvariety of R^3 defined by

$$x_3 = f(x_1, x_2) := \sum_{p \in A} \exp(-x_1^2 - (x_2 - p)^2) \\ \times (x_1^3 \exp(-3(x_2 - p)^2) + (x_2 - p)^{1/3p})$$

where $A = \{p_i; i=1, 2, \dots\}$. Note that at $(u_1, u_2) \in R^2$ either $f(x_1, x_2)$ is real analytic or $f(x_1, x_2) - \exp(-x_1^2 - (x_2 - p)^2)(x_1^3 \exp(-3(x_2 - p)^2) + (x_2 - p)^{1/3p})$ is real analytic where $u_1^3 \exp(-3(u_2 - p)^2) + (u_2 - p)^{1/3p} = 0$. Hence V' is well defined.

We apply a theorem of Grauert ([1, Proposition 7]) and get a connected complex hypersurface V of a Stein neighbourhood of R^3 in C^3