

## 32. On Certain Cubic Fields. II

By Mutsuo WATABE

Department of Mathematics, Keio University

(Communicated by Shokichi IYANAGA, M. J. A., March 12, 1983)

1. For an algebraic number field  $F$ , we denote the class number and the regulator of  $F$  by  $h_F$  and  $R_F$  respectively. The notations  $E_F$  and  $D_F$  have the same meanings as in [4].

In [2], C. L. Siegel proved the following :

**Theorem.** *Let  $F$  be imaginary quadratic field with discriminant  $D_F$ . If  $|D_F| \rightarrow \infty$ , then  $h_F \rightarrow \infty$ .*

To prove this, Siegel used the formula

$$(*) \quad \lim_{D_F \rightarrow \infty} \frac{\log h_F R_F}{\log \sqrt{|D_F|}} = 1,$$

which was established first by Siegel [2] for quadratic fields  $F$ , then by Brauer [1] for general algebraic number fields  $F$ .

The purpose of this note is to show that an analogous result holds for the class of cubic fields treated in [4].

We shall prove :

**Theorem.** *Let  $K$  be cyclic cubic field  $K = \mathbf{Q}(\theta)$ ,  $\text{Irr}(\theta : \mathbf{Q}) = f(x) = x^3 - mx^2 - (m+3)x - 1$ ,  $m \in \mathbf{Z}$ , with square free  $m^2 + 3m + 9$ . Then  $h_K \rightarrow \infty$  as  $D_K \rightarrow \infty$ .*

**Remark.** There are infinitely many rational integers  $m$  such that  $m^2 + 3m + 9$  is square free (cf. [3]).

2. *Proof of Theorem.* We see easily that the roots of  $f(x)$  can be denoted by  $\theta, \theta', \theta''$  so that they satisfy

$$-\frac{m+1}{m} < \theta < -\frac{m^2+1}{m^2}, \quad -\frac{1}{m} < \theta' < -\frac{1}{m^2} \quad \text{and} \quad m+1 < \theta'' < m+2$$

when  $m \geq 1$ .

In [4], we have proved that  $E_K = \langle \pm 1 \rangle \times \langle \theta, \theta' \rangle$ . So we get  $0 < R_K = \text{abs} \{ \log |\theta| \log |\theta''| - (\log |\theta'|)^2 \} < \log(m+1) \log(m+2)$  because  $\log |\theta| < \log(m+1)$ ,  $\log |\theta''| < \log(m+2)$ . This yields

$$R_K = o(m^2) = o(\sqrt{D_K})$$

as we have  $\sqrt{D_K} = m^2 + 3m + 9$  because  $m^2 + 3m + 9$  is square free.

Now, the formula (\*) holds for the set of all algebraic number fields  $F$ . So it holds also for our class of cubic fields  $K$ , where  $\sqrt{D_K} = m^2 + 3m + 9$ , and  $D_K \rightarrow \infty$  means the same meaning as  $m \rightarrow \infty$ .

Therefore we obtain