

102. On Strongly Regular Rings. II

By Katsuo CHIBA and Hisao TOMINAGA

Hokkaido University and Okayama University

(Comm. by Kenjiro SHODA, M. J. A., Sept. 12, 1974)

This is a natural sequel to [1] as well as to [2]. The notation and terminology employed there will be used here. In [2], R. Yue Chi Ming proved the following: *Let A be a ring with identity. If A is left non-singular and every finitely generated left ideal of A is the annihilator of a finitely generated left ideal then A is strongly regular, and conversely.* In this note, we shall prove an analogue without assuming the existence of identity. To this end, we shall introduce the following definition: A left ideal I of a ring A is called *quasi-finitely generated* if either I is finitely generated or $I = I' \oplus l(f)$ with a finitely generated left ideal I' and an idempotent f . Needless to say, if A is a ring with identity then every quasi-finitely generated left ideal is finitely generated.

Theorem. *The following conditions are equivalent:*

- (i) *A ring A is strongly regular.*
- (ii) *A is left non-singular and every quasi-finitely generated left ideal of A is the left annihilator of a quasi-finitely generated left ideal.*

In advance of the proof of our theorem, we state a couple of lemmas whose proofs are obvious by those of Lemmas 1 and 3 in [2].

Lemma 1. *The following conditions are equivalent:*

- (i) *A ring A is left non-singular.*
- (ii) *A is faithful as a right A -module and every left annihilator is closed in A .*

Lemma 2. *The following conditions are equivalent:*

- (i) *A ring A is left non-singular and every closed left ideal of A is two-sided.*
- (ii) *A contains no non-zero nilpotent element and $I + l(I) (= I \oplus l(I))$ is essential in A for every left ideal I of A .*
- (iii) *A contains no non-zero nilpotent element and every closed left ideal of A is the left annihilator of a left ideal.*

Proof of Theorem. Assume first A is strongly regular. Obviously, A is then left non-singular. If I is an arbitrary quasi-finitely generated (left) ideal then one will easily see that $I = l(l(e))$ or $I = l(f - e)$ according as $I = Ae$ or $I = Ae \oplus l(f)$ with some central idempotents e and f . Conversely, assume (ii). Then, it is obvious that A is a left duo ring,