

162. On Mappings of Type l^p

By Masatoshi NAKAMURA

(Comm. by Kinjirô KUNUGI, M.J.A., Oct. 12, 1967)

In this note, we shall improve a result in A. Pietsch [1], and we shall consider the problem 8.4.4 mentioned in [1]. We mainly follow notations in [1].

Let E and F be normed spaces. Let $\mathcal{L}(E, F)$ be all continuous linear mapping of E into F , let $\mathcal{A}(E, F)$ be all mappings $t \in \mathcal{L}(E, F)$ such that $t(E)$ is finite dimensional subspace of F , and let $\mathcal{A}_r(E, F)$ be all mappings $t \in \mathcal{L}(E, F)$ such that $t(E)$ is at most r -dimensional subspace of F .

The norm $\| \cdot \|$ of t in $\mathcal{L}(E, F)$ is defined by $\|t\| = \sup \{\|t(x)\|; \|x\| \leq 1, x \in E\}$ and the r -th approximation number of t in $\mathcal{L}(E, F)$ is defined by

$$\alpha_r(t) = \inf \{\|t - u\|; u \in \mathcal{A}_r(E, F)\}.$$

By $l^p(E, F)$ we denote all $t \in \mathcal{L}(E, F)$ such that $\sum_{r=0}^{\infty} \alpha_r(t)^p < \infty$ holds for all positive number p , and we put

$$\rho_p(t) = \left\{ \sum_{r=0}^{\infty} \alpha_r(t)^p \right\}^{1/p}$$

Lemma. Each mapping $t \in \mathcal{A}(E, F)$ can be represented in following form

$$t(x) = \sum_{i=1}^r \lambda_i \langle x, a_i \rangle y_i, \quad |\lambda_i| \leq \|t\| \quad \text{for } i=1, 2, \dots, r,$$

where $a_i \in U^0$, $y_i \in V$, and λ_i are real or complex numbers.

Proof of this lemma is found in ([1], p. 121).

Proposition. Each mapping $t \in l^p(E, F)$ (where p is any positive number) can be represented in following form, for any positive number δ ,

$$t(x) = \sum_{r=0}^{\infty} \lambda_r \langle x, a_r \rangle y_r$$

and

$$\left\{ \sum_{r=0}^{\infty} |\lambda_r|^p \right\}^{1/p} \leq 2^{1+3/p} (1+\delta) \rho_p(t),$$

where $a_r \in U^0$, $y_r \in V$, and λ_r are real or complex numbers.

Proof. By definition of $\alpha_r(t)$, for all natural number n , there exist u_n in $\mathcal{A}_{2^{n-2}}(E, F)$ such that $\|t - u_n\| \leq (1+\delta) \alpha_{2^{n-2}}(t)$.

Let $v_n = u_{n+1} - u_n$ then we have $d_n \equiv \text{dimensional of } v_n(E) \leq 2^{n+2}$ and

$$\|v_n\| \leq \|t - u_n\| + \|t - u_{n+1}\| \leq 2(1+\delta) \alpha_{2^{n-2}}(t).$$