

## 129. Contributions to the Theory of Semi-groups. V

By Kiyoshi ISÉKI

Kobe University

(Comm. by K. KUNUGI, M.J.A., Oct. 12, 1956)

S. Schwarz [3] considered a disjoint conjugate class decomposition of a semi-group. He proved any finite commutative semi-group has such a decomposition and any character on it takes the same value on each conjugate class. The present author [1, (IV)] proved that, in any commutative periodic semi-group  $S$ , there is a character  $\chi(x)$  of  $S$  such that  $\chi(a) \neq \chi(b)$  if  $a, b$  are two elements of distinct conjugate classes.

Let  $S$  be a strongly reversible (see G. Thierrin [4]) periodic semi-group, and let  $G^{(\omega)}$  be the maximal subgroup of  $K^{(\omega)}$  for an idempotent  $e_a$  (see K. Iséki [1, (I)]). Then, following S. Schwarz [3], for  $a$  of  $G^{(\omega)}$ , the set  $T_a$  of all elements  $x$  of  $K^{(\omega)}$  satisfying  $xe_a = a$  is called a *conjugate class* of  $S$ . By Theorem 2 of K. Iséki [1, (I), p. 174], *the semi-group  $S$  is the set-sum of disjoint conjugate classes  $T_a$ .*

In any strongly reversible compact semi-group<sup>1)</sup>  $S$ , if  $G^{(\omega)}$  is the maximal subgroup of  $K^{(\omega)}$  containing an idempotent  $e_a$ ,  $S$  is the set-sum of  $K^{(\omega)}$  (see K. Iséki [2]) and  $K^{(\omega)}e_a = e_aK^{(\omega)} = G^{(\omega)}$ . Therefore we can define conjugate classes of  $S$ , and  $S$  is the set-sum of disjoint conjugate classes  $T_a$ . Hence if a given semi-group is compact commutative, it is decomposed into disjoint conjugate classes  $T_a$ .

Now, let  $S$  be a strongly reversible periodic homogroup (see G. Thierrin [5]), then  $S$  has only one smallest idempotent  $e$  (see K. Iséki [1, (II)]).

Let  $\chi$  be a character of  $S$ , i.e. a homomorphism into the multiplicative group of complex numbers of absolute value one. Let  $G^{(e)}$  be a maximal group relative to the smallest idempotent  $e$ . From  $T_a e_a = a \in G^{(\omega)}$  and  $e_a e = e = e e_a$ , we have

$$T_a e = T_a e_a e = a e.$$

Hence

$$\chi(T_a) = \chi(T_a e) = \chi(a e).$$

Therefore we have the following

*Theorem 1. In a strongly reversible periodic homogroup  $S$ , any character of  $S$  takes the same value on each conjugate class.*

Further, we have

*Theorem 2. Any character of a compact homogroup takes the same*

---

1) For topological semi-groups, see A. D. Wallace [6] or [7].