

130. On a Radical in a Semiring

By Kiyoshi ISÉKI and Yasue MIYANAGA

(Comm. by K. KUNUGI, M.J.A., Oct. 12, 1956)

In our previous paper [2], we considered the structure space of maximal ideals of a commutative semiring, and K. Iséki [3], one of the present authors, considered some relations of two structure spaces of it. In this paper, we shall consider a new kind of ideals of a semiring A with 0 (for the definition, see our paper [2]). A similar theory of an associative ring was treated by L. Fuchs [1].

An element a of A is said to be a *left zerodivisor* if there is a $b (\neq 0)$ of A such that $ab=0$. Let \mathfrak{A} be a two sided ideal, if every element of \mathfrak{A} is a left zerodivisor, then \mathfrak{A} is said to be *left zerodivisor*. In the sequel, by the term *ideal*, we mean a *two sided ideal*. An ideal is *maximal left zerodivisor* if there is no left zerodivisor ideal containing properly it. By Zorn's lemma, any left zerodivisor ideal is contained in a maximal left zerodivisor. Following L. Fuchs [1], we shall define a left zeroid ideal. If \mathfrak{A} is an ideal and $\mathfrak{A} + \mathfrak{B}$ for every left zerodivisor ideal \mathfrak{B} is a left zerodivisor, then \mathfrak{A} is said to be a *left zeroid ideal*. Therefore we have the following propositions which are proved easily.

Proposition 1. The sum of two left zeroid ideals is a left zeroid ideal.

Proposition 2. The join of all left zeroid ideals is also a left zeroid ideal.

The left zeroid ideal stated in Proposition 2 is said to be the *left radical* of A which is denoted by $\mathfrak{R}^{(l)}$. Similarly, we can define right zerodivisor ideals, right zeroid ideals and the right radical $\mathfrak{R}^{(r)}$ of A . We shall define the radical \mathfrak{R} of A as $\mathfrak{R}^{(l)} \cap \mathfrak{R}^{(r)}$.

If every element of an ideal \mathfrak{A} is nilpotent, \mathfrak{A} is said to be a *nil ideal*. Then any nil ideal \mathfrak{N} is left zeroid and right zeroid.

Let b be an element of \mathfrak{N} , and let \mathfrak{A} be a left zerodivisor ideal. Then there is some positive integer n such that $b^n=0$. For an element a of \mathfrak{A} , $(a+b)^n$ is in \mathfrak{A} . Hence there is an element $c (\neq 0)$ such that $(a+b)^n c=0$. Let m be the least positive integer such that $(a+b)^m c=0$, then we have $(a+b)^{m-1} c \neq 0$. Hence $(a+b) \times (a+b)^{m-1} c = 0$ implies that $a+b$ is a left zerodivisor. Hence \mathfrak{N} is a left zeroid ideal. Similarly we can prove that \mathfrak{N} is a right zeroid ideal.

Therefore the nil radical defined as the join of all nil ideals of A is contained in the radical \mathfrak{R} .

On the left radical $\mathfrak{R}^{(l)}$, we have the following