

## 28. A Note on a Conjecture of K. Harada

By Masao KIYOTA<sup>\*)</sup> and Tetsuro OKUYAMA<sup>\*\*)</sup>

(Communicated by Shokichi IYANAGA, M. J. A., Feb. 12, 1981)

Let  $G$  be a finite group and  $p$  be prime number. Let  $\{\chi_1, \dots, \chi_s\}$  be the set of all irreducible complex characters of  $G$ . For a subset  $J$  of the index set  $\{1, \dots, s\}$ , we put  $\{\chi_J\} = \{\chi_j; j \in J\}$  and  $\rho_J = \sum_{j \in J} \chi_j(1)\chi_j$ .

In [1], K. Harada stated the following;

**Conjecture A.** *If  $\rho_J(x) = 0$  for any  $p$ -singular element  $x$  of  $G$ , then  $\{\chi_J\}$  is a union of  $p$ -blocks of  $G$ .*

And he proved that if a Sylow-subgroup of  $G$  is cyclic, then Conjecture A holds. In this note we prove the conjecture in the following another case.

**Theorem.** *If  $G$  is  $p$ -solvable, then Conjecture A holds.*

**Proof.** Assume that  $\{\chi_J\}$  satisfies the condition of Conjecture A. As in [1], we may assume that  $\{\chi_J\} \subseteq B$ , for some  $p$ -block  $B$  of  $G$ . So we need to show  $\{\chi_J\} = B$  or  $\{\chi_J\} = \phi$ .

By rearranging the index set if necessary, we may assume that  $B = \{\chi_1, \dots, \chi_k\}$ . Let  $\{\varphi_1, \dots, \varphi_l\}$  be the set of all irreducible Brauer characters in  $B$  and  $\{\Phi_1, \dots, \Phi_l\}$  be the set of all principal indecomposable characters in  $B$ . For  $x \in G$ , we define  $\chi_B(x)$  to be the column vector of size  $k$  whose  $i$ -th component is  $\chi_i(x)$ . For  $1 \leq m \leq l$ , let  $d_m$  be the column of size  $k$  whose  $i$ -th component  $d_{im}$  is the decomposition number of  $\chi_i$  with respect to  $\varphi_m$ . Then we have

$$\chi_B(x) = \sum_{m=1}^l d_m \varphi_m(x) \quad \text{for any } p\text{-regular element } x.$$

In particular

$$\chi_B(1) = \sum_{m=1}^l d_m \varphi_m(1).$$

Let  $\chi_J$  be the column of size  $k$  whose  $i$ -th component  $a_i$  is defined as follows. If  $i \in J$ , then  $a_i = \chi_i(1)$  and  $a_i = 0$  otherwise. At first we show that  $\chi_J$  is a linear combination of  $d_m$ ,  $m = 1, \dots, l$ . Since  $\rho_J$  vanishes on all  $p$ -singular elements of  $G$ ,  $\rho_J$  is an integral linear combination of  $\Phi_m$ ,  $m = 1, \dots, l$ ;

$$\rho_J = \sum_m \alpha_m \Phi_m = \sum_m \alpha_m \sum_i d_{im} \chi_i = \sum_i \left( \sum_m \alpha_m d_{im} \right) \chi_i.$$

By the linear independence of  $\{\chi_i\}$ , we obtain

$$\chi_J = \sum_m \alpha_m d_m.$$

---

<sup>\*)</sup> Department of Mathematics, University of Tokyo.

<sup>\*\*)</sup> Department of Mathematics, Osaka City University.