

A Form of Classical Picard Principle

By Yujiro ISHIKAWA,^{*)} Mitsuru NAKAI,^{**)} and Toshimasa TADA^{***)}

(Communicated by Kiyosi ITÔ, M. J. A., Jan. 12, 1996)

Let H_d be the fundamental harmonic function on the Euclidean space \mathbf{R}^d of dimension $d \geq 2$, i.e. $H_2(x) = \log(1/|x|)$ and $H_d(x) = 1/|x|^{d-2}$ ($d \geq 3$), where $|x| = (\sum_{i=1}^d |x^i|^2)^{1/2}$ is the length of a vector $x = (x^1, \dots, x^d) \in \mathbf{R}^d$. We denote by B^d the unit ball $|x| < 1$ in \mathbf{R}^d and by B_0^d the punctured unit ball $0 < |x| < 1$ in \mathbf{R}^d . Then we have the following

Theorem A. *If $u \geq 0$ is harmonic in B_0^d ($d \geq 2$), then $u = cH_d + v$, where $c \geq 0$ is a constant and v is harmonic on B^d .*

This result has been called the *Picard principle* by many authors since Bouligand [4] and then Brelot [5] first used the term because of the papers of Picard [10,11] (see also Stôzek [13]); it is also referred to as the *Bôcher theorem* by Helms [6], Wermer [14], and Axler *et al.* [2], etc. since the result is proved by Bôcher [3] 20 years earlier than Picard. Anyway this is one of the results in the potential theory much talked about from various view points: thousands of different proofs have been given to the result; the result is also discussed in the frame of wider degenerate harmonicity such as one given by the Schrödinger equations with potentials having singularities at the origin (cf. e.g. Pinsky [12]); the Martin theory is of course another extension. Recently the following result of Anandam and Damlaakhi [1] called our attention:

Theorem B. *Suppose u is harmonic on B_0^2 such that $u(x) \geq o(|x|^{-s})$ as $|x| \rightarrow 0$ with*

$s \leq 1$. Then $u = cH_2 + v$, where c is a constant and v is harmonic on B^2 .

This is of course superficially a generalization of Theorem A for $d = 2$. Anandam and Damlaakhi [1] proved this in complex analytic way and thus the restriction to the dimension $d = 2$ seems to be essential in their proof. The purpose of this note is to remark that the Fourier expansion method, one of the most standard ways of proving Theorem A, also instantly proves not only Theorem B but also its generalization to higher dimensions. Namely, we will prove the following

1. Theorem. *Suppose u is harmonic in B_0^d such that $u(x) \geq o(|x|^{-s})$ as $|x| \rightarrow 0$ with $s \leq d - 1$. Then $u = cH_d + v$, where c is a constant and v is harmonic on B^d .*

Proof. We use the polar coordinates $x = r\xi$ for points $x \in \mathbf{R}^d \setminus \{0\}$, where $r = |x| > 0$ and $\xi = x/|x| \in S^{d-1} = \partial B^d$. We choose and then fix an orthonormal basis $\{S_{nj} : j = 1, \dots, N(n)\}$ of the subspace of all spherical harmonics of degree n of $L_2(S^{d-1}, d\sigma)$, where $d\sigma$ is the area element on S^{d-1} . Then $\{S_{nj} : j = 1, \dots, N(n) ; n = 0, 1, \dots\}$ is a complete orthonormal system of $L_2(S^{d-1}, d\sigma)$. We have, as the special case of the addition theorem, $\sum_{j=1}^{N(n)} S_{nj}(\xi)^2 = N(n)/\sigma_d$ on S^{d-1} ($n = 0, 1, \dots$), where σ_d is the surface area $\sigma(S^{d-1})$ of S^{d-1} . Here $N(0) = 1$ and $N(n) = (2n + d - 2)\Gamma(n + d - 2)/\Gamma(n + 1)\Gamma(d - 1)$ for $n = 1, 2, \dots$. Then we have the following Fourier expansion of $u(r\xi)$ in terms of spherical harmonics $\{S_{nj}\}$:

$$(2) \quad u(r\xi) = \sum_{n=0}^{\infty} \left(\sum_{j=1}^{N(n)} a_{nj} S_{nj}(\xi) \right) r^n + b_{01} H_d(r) + \sum_{n=1}^{\infty} \left(\sum_{j=1}^{N(n)} b_{nj} S_{nj}(\xi) \right) r^{-n-d+2},$$

where $H_d(r) = H_d(x)$ with $|x| = r$ and a_{nj} and b_{nj} ($j = 1, \dots, N(n) ; n = 0, 1, \dots$) are constants. Here the series on the right hand side of (2) converges uniformly in $\xi \in S^{d-1}$ for any fixed $0 < r < 1$. Multiply $\sqrt{N(n)/\sigma_d} \pm S_{nj}(\xi) \geq 0$ ($n \geq 1$) to both sides of $u(r\xi) \geq o(r^{-s})$ and

^{*)} Department of Electrical and Computer Engineering, Nagoya Institute of Technology.

^{**)} Department of Mathematics, Nagoya Institute of Technology.

^{***)} Department of Mathematics, Daido Institute of Technology.

1991 *Mathematics Subject Classification*. Primary 31B25; Secondary 31C35.

This work was partly supported by Grant-in-Aid for Scientific Research, Nos. 06640227, 07640196, Ministry of Education, Science, Sports and Culture, Japan.