

A note on Terai's conjecture concerning Pythagorean numbers^{*)}

By Xigeng CHEN^{**)} and Maohua LE^{***)}

(Communicated by Shokichi IYANAGA, M. J. A., May 12, 1998)

Abstract: Let (a, b, c) be a primitive Pythagorean triple with $2 \mid a$. In this note we prove that if $b \not\equiv 1 \pmod{16}$, $b^2 + 1 = 2c$, b and c are both odd primes, then the equation $x^2 + b^y = c^z$ has only the positive integer solutions $(x, y, z) = (a, 2, 2)$.

1. Introduction. Let $\mathbf{Z}, \mathbf{N}, \mathbf{Q}$ be the sets of integers, positive integers and rational numbers respectively. Let (a, b, c) be a primitive Pythagorean triple such that

$$(1) \quad a^2 + b^2 = c^2, a, b, c \in \mathbf{N}, \\ \gcd(a, b, c) = 1, 2 \mid a.$$

Then we have

$$(2) \quad a = 2st, b = s^2 - t^2, c = s^2 + t^2, \\ \text{where } s, t \text{ are positive integers satisfying } s > t, \\ \gcd(s, t) = 1 \text{ and } 2 \mid st. \text{ In 1993, Terai [4] conjectured that the equation}$$

$$(3) \quad x^2 + b^y = c^z, x, y, z \in \mathbf{N}, \\ \text{has only the solution } (x, y, z) = (a, 2, 2). \\ \text{This conjecture is not solved as yet. In [4], Terai proved that if } b \equiv 1 \pmod{4}, b^2 + 1 = 2c, b, c \\ \text{are odd primes, } c \text{ splits in the imaginary quadratic field } K = \mathbf{Q}(\sqrt{-b}) \text{ and the order } d \text{ of a prime ideal divisor of } [c] \text{ in } K \text{ satisfies either } d = 1 \text{ or } 2 \mid d, \text{ then (3) has only the solution } (x, y, z) = (a, 2, 2). \text{ In this note we prove the following general result.}$$

Theorem. If $b \not\equiv 1 \pmod{16}$, $b^2 + 1 = 2c$, b, c are both odd primes, then (3) has only the solution $(x, y, z) = (a, 2, 2)$.

2. Preliminaries. **Lemma 1** ([2] and [3]). The equation

$$X^2 + 1 = 2Y^n, X, Y, n \in \mathbf{N}, Y > 1, n > 2, \\ \text{has only the solution } (X, Y, n) = (239, 13, 4).$$

Lemma 2 ([1, Lemma 2]). Let k be a positive integer. All solutions (X, Y, Z) of the equation

$$X^2 + Y^2 = k^2, X, Y, Z \in \mathbf{Z}, \\ \gcd(X, Y) = 1, Z > 0$$

are given by

$$Z = n, X + Y\sqrt{-1} = \lambda_1(X_1 + \lambda_2 Y_1 \sqrt{-1})^n \\ \text{or } \lambda_1(Y_1 + \lambda_2 X_1 \sqrt{-1})^n,$$

$$n \in \mathbf{N}, \lambda_1, \lambda_2 \in \{-1, 1\},$$

where X_1, Y_1 run through all positive integers satisfying

$$X_1^2 + Y_1^2 = k, \gcd(X_1, Y_1) = 1.$$

3. Proof of theorem. Since $b^2 + 1 = 2c$ and $2 \nmid b$, we have

$$(4) \quad \left(\frac{b+1}{2}\right)^2 + \left(\frac{b-1}{2}\right)^2 = c.$$

Notice that c is an odd prime. We see from (4) that

$$(5) \quad (X_1, Y_1) = \left(\frac{b+1}{2}, \frac{b-1}{2}\right), \left(\frac{b-1}{2}, \frac{b+1}{2}\right)$$

are all positive integers X_1, Y_1 satisfying

$$(6) \quad X_1^2 + Y_1^2 = c, \gcd(X_1, Y_1) = 1.$$

Hence, by (2) and (4), we get $s = (b+1)/2$, $t = (b-1)/2$, $s = t+1$,

$$(7) \quad a = 2t(t+1), b = 2t+1, c = 2t^2 + 2t + 1.$$

Let (x, y, z) be a solution of (3). Since b is an odd prime, if $2 \mid z$, then from (3) we get $c^{z/2} + x = b^y$, and $c^{z/2} - x = 1$. It implies that

$$(8) \quad b^y + 1 = 2c^{z/2}.$$

Since $b+1 = 2t+2$ and $c \equiv 1 \pmod{2t+2}$ by (7), we find from (8) that $2 \mid y$. Since $b^2 + 1 = 2c$, if $z/2 = 1$, then from (1) and (8) we get the solution $(x, y, z) = (a, 2, 2)$. If $z/2 = 2$, then we have $b^y + 1 = 2c^2 = 2((b^2 + 1)/2)^2$. It follows that $2 \equiv 1 \pmod{b}$, a contradiction. If $z/2 > 2$, by Lemma 1, then we get $(b, y, c, z) = (239, 2, 13, 8)$. It is impossible, by (3). Thus, (3) has only the solution $(x, y, z) = (a, 2, 2)$ with $2 \mid z$.

If $2 \nmid y$ and $2 \nmid z$, then the equation $X^2 + Y^2 = c^z, X, Y, Z \in \mathbf{Z}$,

1991 Mathematics Subject Classification. 11D61.

^{*)} Supported by the National Natural Science Foundation of China and the Guangdong Provincial Natural Science Foundation.

^{**)} Department of Mathematics, Maoming Education College, P. R. China.

^{***)} Department of Mathematics, Zhanjiang Teachers College, P. R. China.