

ABSTRACTS OF PAPERS

SUBMITTED FOR PRESENTATION TO THE SOCIETY

The following papers have been submitted to the Secretary and the Associate Secretaries of the Society for presentation at meetings of the Society. They are numbered serially throughout this volume. Cross references to them in the reports of the meetings will give the number of this volume, the number of this issue, and the serial number of the abstract.

ALGEBRA AND THEORY OF NUMBERS

230. Reinhold Baer: *Rings with duals.*

The ring R' is said to be a right-dual of the ring R , if there exists a duality between the partially ordered set of all the right-ideals in R and the partially ordered set of all the right-ideals in R' . If the ring R possesses a right-identity element, and if there exists to every quotient-ring of R a right-dual possessing a left-identity element, then both the maximum and the minimum condition are satisfied by the right-ideals in R , and R meets a generalized uni-seriality requirement. A complete theory of the primary rings with right-duals may be developed, determining their structure and their dualities. (Received May 13, 1942.)

231. Nathan Jacobson: *Classes of restricted Lie algebras of characteristic p . II.*

Let $\mathfrak{A} = \Phi(x_1, \dots, x_m)$ be an associative commutative algebra having the basis $x_1^{k_1} x_2^{k_2} \dots x_m^{k_m}$, $0 \leq k_i \leq p-1$ such that $x_i^p = \xi_i$ is in the field Φ of characteristic p . The restricted Lie algebras considered in this paper are the derivation algebras θ of \mathfrak{A} 's of the above type. The case where \mathfrak{A} is a field was defined in a previous paper of the author and the case where $m=1$ and $\xi_1=1$ has been considered by Witt, by Zassenhaus and by Ho-Jui Chang. The author proves that θ is normal simple except when $p=2$ and $m=1$. The derivations of θ are all inner and if $p \geq 5$ the automorphisms of θ have the form $D \rightarrow S^{-1}DS$ where S is an automorphism of \mathfrak{A} . This implies that if $p \geq 5$, two algebras $\theta(\mathfrak{A}_1)$ and $\theta(\mathfrak{A}_2)$ are isomorphic if and only if the associative algebras \mathfrak{A}_1 and \mathfrak{A}_2 are isomorphic. (Received April 3, 1942.)

232. A. E. Ross: *On a theorem of Kloosterman.*

Kloosterman (Acta Mathematica, vol. 49 (1926)) determined conditions under which the form (1) $f = ax^2 + by^2 + cz^2 + du^2$ would represent all large integers and pointed out that there was a finite (and actually small) number of such forms f for which his method failed to yield conclusive results. Upon close examination of these exceptional cases one notices that should one replace the original form f by a suitably chosen equivalent form $f_1 = x'Ax$ (this Bulletin, abstract 45-9-369) then one can reduce the problem to the study of a related form of type (1) to which Kloosterman's sufficient conditions apply. Although the need for choosing the quotients of the upper left-hand corner principal minors of A by the order invariants as primes introduces an element of tentativeness, the actual computation is carried out without difficulty. (Received April 17, 1942.)