

AN ARITHMETICAL IDENTITY FOR THE FORM $ab - c^2$

WALTER H. GAGE

1. **Introduction.** The number of solutions in positive integers of the equation $n = xy + yz + zx$, n a positive integer, has been investigated Liouville,¹ Bell,² and Mordell.³ Mordell, who was the first to obtain complete results, gave a strictly arithmetical treatment, while Bell made use of formulae which he obtained by paraphrasing theta-function identities. Although the latter considered only the case of n prime, his methods were extended later to the general case.⁴

Making use of other formulae derived by the method of paraphrase, Bell⁵ has also solved the problem of representations in the forms $xy + yz + 2zx$, $xy + 2yz + 2zx$. As he has pointed out, a feature of the method is the handling of the two forms simultaneously.

In this paper we derive by elementary methods a simple identity which on specialization not only yields complete results for representations of n in the forms

$$xy + yz + zx, \quad xy + 2yz + 2zx, \quad xy + yz + 2zx,$$

but as in Bell's paper,⁵ handles the latter two forms at the same time.

2. **Fundamental identity.** Let $f(a, b, c)$ be a function, uniform and finite for all integer triples (a, b, c) , but otherwise (so far) completely arbitrary. If the summation sign refers to the sum over all those integer solutions (a, b, c) of $n = ab - c^2$ subject to the restrictions indicated under it, we then have

$$(1) \quad \sum_{a, b > c > 0} f(a, b, c) = \sum_{a > b > c > 0} f(a, b, c) + \sum_{b > a > c > 0} f(a, b, c) + \sum_{a = b > c > 0} f(a, b, c).$$

Imposing on $f(a, b, c)$ the condition

$$(2) \quad f(a, b, c) = f(b, a, c)$$

Received by the editors February 13, 1942.

¹ Journal de Mathématiques, (2), vol. 7 (1862), p. 44.

² E. T. Bell, *Class numbers and the form $yz + zx + xy$* , Tôhoku Mathematical Journal, vol. 19 (1921), pp. 105-116.

³ L. J. Mordell, *On the number of solutions in positive integers of the equation $yz + zx + xy = n$* , American Journal of Mathematics, vol. 45 (1923), pp. 1-4.

⁴ W. H. Gage, *Representations in the form $xy + yz + zx$* , American Journal of Mathematics, vol. 51 (1929), pp. 345-348.

⁵ E. T. Bell, *Numbers of representations of integers in a certain triad of ternary quadratic forms*, Transactions of this Society, vol. 32 (1930), pp. 1-5.