

We can of course prove the following theorem: The necessary and sufficient condition for the continuum to be of power \aleph_{x+1} is that R shall be the sum of \aleph_x sets consisting of rationally independent numbers, and that R shall not be the sum of less than \aleph_x such sets. The proof is the same as that of Theorem 2.

UNIVERSITY OF PENNSYLVANIA AND
THE INSTITUTE FOR ADVANCED STUDY

**ADDENDUM TO THE PAPER "GENERALIZED FISCHER
GROUPS AND ALGEBRAS"**

R. H. BRUCK

The author regrets the omission, in a paper which recently appeared,¹ of an important reference to a paper by N. Jacobson.² Indeed Lemma IIIa of the author's paper, and its immediate consequence, Theorem I, are rather special cases (albeit independently obtained) of Theorem I of the latter paper. Accordingly Professor Jacobson's name should have appeared in the introduction along with those of M. Schiffer and W. Specht, and chronologically before that of Specht.

UNIVERSITY OF WISCONSIN

Received by the editors December 9, 1942.

¹ R. H. Bruck, *Generalized Fischer groups and algebras*, Bull. Amer. Math. Soc. vol. 48 (1942) pp. 618-626, in particular p. 623.

² N. Jacobson, *Normal semi-linear transformations*, Amer. J. Math. vol. 61 (1939) pp. 45-58, in particular p. 49.