

RESEARCH ANNOUNCEMENTS

The purpose of this department is to provide early announcement of significant new results, with some indications of proof. Although ordinarily a research announcement should be a brief summary of a paper to be published in full elsewhere, papers giving complete proofs of results of exceptional interest are also solicited. Manuscripts more than eight typewritten double spaced pages long will not be considered as acceptable.

SOME ARITHMETIC PROPERTIES OF THE BELL POLYNOMIALS¹

BY L. CARLITZ

Communicated by M. L. Curtis, September 14, 1964

Let $\alpha_1, \alpha_2, \alpha_3, \dots$ denote indeterminates. The Bell polynomials $\phi_n(\alpha_1, \alpha_2, \alpha_3, \dots)$ may be defined by $\phi_0 = 1$ and

$$\phi_n = \phi_n(\alpha_1, \alpha_2, \alpha_3, \dots) = \sum \frac{n!}{k_1!(1!)^{k_1}k_2!(2!)^{k_2}\dots} \alpha_1^{k_1} \alpha_2^{k_2} \dots,$$

where the summation is over all nonnegative integers k_j such that

$$k_1 + 2k_2 + 3k_3 + \dots = n.$$

For references see Bell [1] and Riordan [3, p. 36]. The general coefficient

$$A_n(k_1, k_2, k_3, \dots) = \frac{n!}{k_1!(1!)^{k_1}k_2!(2!)^{k_2}\dots}$$

is integral.

Some arithmetic properties of the polynomial ϕ_n have been given by Bell and some additional properties were obtained by the present writer [2]. In particular, the latter showed that

$$\phi_{pn}(\alpha_1, \alpha_2, \alpha_3, \dots) \equiv \phi_n(\phi_p, \alpha_p, \alpha_{2p}, \dots) \pmod{p},$$

where p is a prime and the first argument on the right is ϕ_p and not α_p .

In the present paper we consider the following problem. Let p be a fixed prime and let $\theta(n)$ denote the number of coefficients $A_n(k_1, k_2, k_3, \dots)$ that are prime to n . Then we can state the following results.

I. Let

¹ Supported in part by NSF grant GP-1593.