BULLETIN OF THE
AMERICAN MATHEMATICAL SOCIETY
Volume 79, Number 4, July 1973

BOUNDED SOLUTIONS OF WHOLE-LINE DIFFERENTIAL EQUATIONS

BY DAVID LOWELL LOVELADY

Communicated by Fred Brauer, December 18, 1972

Let Y be a finite-dimensional linear space with norm $|\ |$, and let $R = (-\infty, \infty)$. Let $\mathscr{A}$ be the algebra of linear functions from Y to Y with induced norm $\|\ \|$, and let A be a locally integrable function from R to $\mathscr{A}$. W. A. Coppel [**2**], [**3**, Theorem 1, p. 131] has determined necessary and sufficient conditions for

$$\text{(NH)} \qquad u'(t) = f(t) + A(t)u(t)$$

to have at least one solution u in $\mathscr{L}^\infty[R^+, Y]$ (where $R^+ = [0, \infty)$) for each f in $\mathscr{L}^\infty[R^+, Y]$. R. Conti [**1**] has solved the same problem for f in $\mathscr{L}^p[R^+, Y]$, $p \geqq 1$, and u in $\mathscr{L}^\infty[R^+, Y]$. In this note we indicate how the results of [**4**] solve both of these problems for equations on R instead of R^+. Let Φ be the fundamental solution for

$$\text{(H)} \qquad v'(t) = A(t)v(t),$$

i.e., Φ is that locally absolutely continuous function from R to $\mathscr{A}$ such that $\Phi(t) = I + \int_0^t A(s)\Phi(s)\, ds$ whenever t is in R.

THEOREM. *Statements* (i) *and* (ii) *are equivalent and statements* (iii) *and* (iv) *are equivalent.*

(i) *If f is in $\mathscr{L}^\infty[R, Y]$ there is a solution u of* (NH) *in $\mathscr{L}^\infty[R, Y]$.*

(ii) *There are three supplementary projections P_{-1}, P_0, and P_1 and a number K such that if t is in R then*

$$\int_{-\infty}^{t} \|\Phi(t)P_{-1}\Phi(s)^{-1}\|\, ds + \left| \int_0^t \|\Phi(t)P_0\Phi(s)^{-1}\|\, ds \right| + \int_t^\infty \|\Phi(t)P_1\Phi(s)^{-1}\|\, ds \leqq K.$$

(iii) *If f is in $\mathscr{L}^p[R, Y]$, $p > 1$, there is a solution u of* (NH) *in $\mathscr{L}^\infty[R, Y]$.*

(iv) *There are P_{-1}, P_0, P_1, and K as in* (ii) *such that if t is in R then*

AMS (MOS) subject classifications (1970). Primary 34A30, 34E99.