

EXISTENCE OF SOLUTIONS OF DIFFERENTIAL EQUATIONS IN BANACH SPACE

BY WILLIAM J. KNIGHT

Communicated by Fred Brauer, July 23, 1973

The results announced here concern the existence of a solution to the general initial value problem

$$(1) \quad x'(t) = f(t, x(t)), \quad x(0) = x_0,$$

in which $x(t)$ lies in a Banach space X for $t \in J = [0, a]$. Recent results for this problem have been announced in this Bulletin by S. N. Chow and J. D. Schuur [1] and by W. E. Fitzgibbon [2]. Related results were obtained earlier by F. Browder [3]. Here however, X is not assumed to be separable or reflexive, although as usual f will be continuous in x with respect to the weak topology on X .

A *pseudo-solution* of (1) is an absolutely continuous function $x: J \rightarrow X$ with pseudo-derivative (see Pettis [4]) satisfying (1). A *strong solution* of (1) is a strongly absolutely continuous function $x: J \rightarrow X$ with strong derivative ($\lim_{h \rightarrow 0} (x(t+h) - x(t))/h$ in norm) satisfying (1) a.e. on J . For notions of absolute continuity, see Hille and Phillips [5, p. 76].

In what follows let B denote an open ball about some point $x_0 \in X$, let $I = [0, b]$ be a compact interval, and let f be a function from $I \times B$ into X .

THEOREM A. *Assume these hypotheses:*

(a) *For a.e. $t \in I$, $f(t, x)$ is continuous in the variable x with respect to the weak topology on B and X .*

(b) *For each strongly absolutely continuous function $y: I \rightarrow B$, $f(t, y(t))$ is Pettis integrable on I .*

(c) *For some null set $N \subset I$, the weak closure of $f((I - N) \times B)$ is weakly compact in X .*

Then (1) has a (possibly nonunique) pseudo-solution on a subinterval $J = [0, a]$ of I .

AMS (MOS) subject classifications (1970). Primary 34G05; Secondary 47H10.

Key words and phrases. Initial value problem, pseudo-solution, pseudo-derivative, strong solution, fixed point, strongly measurable, uniformly convex space.

Copyright © American Mathematical Society 1974