

A GENERALIZED WEIL TYPE REPRESENTATION AND A FUNCTION ANALOGOUS TO e^{-x^2}

BY TOMIO KUBOTA¹

Communicated by D. J. Lewis, April 28, 1975

Let $dV(z)$ be the euclidean measure of \mathbb{C} , and let $n \geq 2$ be a natural number. Put $e(z) = \exp(\pi\sqrt{-1}(z + \bar{z}))$, $\zeta = e(1/n)$, and consider the Hilbert space H_n consisting of all functions Φ on \mathbb{C} such that $\Phi(\zeta z) = \Phi(z)$ and $\|\Phi\| < \infty$, where the norm is coming from the inner product

$$(f_1, f_2) = \int_{\mathbb{C}} f_1(z) \overline{f_2(z)} |z|^{2n-4} dV(z).$$

Denote by $\Phi \rightarrow \Phi^*$ the integral linear transformation given by

$$\Phi^*(t) = \int_{\mathbb{C}} \Phi(z) k(zt) |z|^{2n-4} dV(z)$$

with $k(z) = n^2 \lim_{Y \rightarrow \infty} \int_{|z| < Y} e(z^n/w^n) e(w^n) dV(z)$.

Denote furthermore by $\sigma = (a, b; c, d)$ an element of $G = SL(2, \mathbb{C})$ for which (a, b) is the first row and (c, d) is the second, and define an operator $r_n(\sigma)$ of H_n for three types of elements $\sigma_1 = (a, 0; 0, a^{-1})$, $\sigma_2 = (1, b; 0, 1)$, and $\sigma_3 = (-c^{-1}, 0; c, 0)$ by $(r_n(\sigma_1)\Phi)(t) = |a|^{n(n-1)/2} \Phi(a^2/n t)$, $(r_n(\sigma_2)\Phi)(t) = \Phi(t)e(bt^n)$, and $(r_n(\sigma_3)\Phi)(t) = |c|^{-n(n-1)/2} \Phi^*(c^{-2}/n t)$. Then, it follows from the results, to be announced in [2] in detail, that $r_n(\sigma_i)$ extends multiplicatively to an irreducible unitary representation $\sigma \rightarrow r_n(\sigma)$ of G of class one on H_n belonging to the supplementary series. If $n = 2$, then $k(z)$ reduces to $e(2z) + e(-2z)$, and $\sigma \rightarrow r_n(\sigma)$ reduces essentially to a special case of the representation given in [3].

These results, viewed so to speak from the reverse side, yield as a byproduct a representation theoretic characterization of a special function. Namely, we obtain

THEOREM. *Up to a constant factor, the function $h(t) = tK_{1/n}(2\pi|t|^n)$ is the only function in H_n which is invariant by all $r_n(\sigma)$ with $\sigma \in K = SU(2)$, where $K_{1/n}$ is a modified Bessel function.*

This Theorem follows from the facts, proved in [2], that $h(t)$ is actually invariant by all $r_n(\sigma)$, ($\sigma \in K$), and that the set of all $r_n(\sigma)h(t)$, ($\sigma \in G$), is dense in H_n .

AMS (MOS) subject classifications (1970). Primary 22E45; Secondary 33A40.

Key words and phrases. Unitary representation, Weil representation, special functions, Bessel functions, theta functions, automorphic functions.

¹Research supported in part by NSF Grant-43950(SK-CUCB).