

SINGULAR SOBOLEV CONNECTIONS WITH HOLONOMY

L. M. SIBNER AND R. J. SIBNER

We consider local Sobolev connections on $SU(2)$ bundles over the complement, in R^4 , of a smoothly embedded compact 2-manifold. Finite action implies that a holonomy condition is satisfied and we obtain an a priori estimate for the connection 1-form in terms of curvature and the flat connection carrying the holonomy. The a priori estimate classifies the possible singularities in these connections by the set of flat connections. In a certain case, this leads to smoothness and extendability results.

Let N be a full 4-dimensional neighborhood of the singular set S . The objects of study are connections $D = d + A$ defined on $SU(2)$ bundles over $X = N \setminus S$. We assume that $A \in H^2_{1,\text{loc}}(X)$ and that the action is finite, i.e., the curvature $F = dA + A \wedge A$ is in $L^2(N)$.

The following *holonomy condition* was first stated by Cliff Taubes. Choose coordinates (r, θ, u, v) with (u, v) coordinates on S and (r, θ) coordinates in a plane normal to S . Fixing u and v , and denoting by A_θ the θ component of A , the initial value problem for an $SU(2)$ valued function,

$$\frac{dg_r}{d\theta} + A_\theta g_r = 0, \quad g_r(0) = I,$$

has a unique solution $g_r(\theta)$, with $g_r(2\pi) = J_r \in SU(2)$. The holonomy condition we require is

$$(H) \quad \lim_{r \rightarrow 0} J_r = J^b \text{ exists.}$$

This condition is *gauge invariant* up to conjugacy in $SU(2)$. Our results can be formulated in two theorems.

THEOREM 1. *If A and F are smooth on $N \setminus S$ and $F \in L^2(N)$, then (H) is satisfied for almost all u and v . Up to conjugacy, the limit is independent of u and v .*

Next, assume (H) holds. Locally, the conjugacy class $[J^b] \in SU(2)$ uniquely defines a flat connection $A^b = C d\theta$ with C a constant element of $su(2)$ determined up to a similarity transformation. Our second result uses holonomy to obtain an a priori estimate. We denote by X_0 and N_0 the intersections of X and N with a small open set in R^4 having nonvoid intersection with S .

Received by the editors February 23, 1988.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 35J60, 53C80.

Research of the first author partially supported by NSF grant DMS-8501419.

Research of the second author partially supported by NSF grant INT-8411481.