

$$H^n(x) = \frac{R_n(x)}{(x-1)^n}.$$

The relation (7) gives many generalizations of (9). For example we can take $m = p^e$ in lieu of $m = p$. Further details I hope to give in another paper on Bernoulli numbers and Euler polynomials.

UNIVERSITY OF TEXAS

A THEOREM ON MEAN RULED SURFACES

BY MALCOLM FOSTER

Consider the ruled surface formed by the normals to a surface S along some curve C on S . We ask: What are the curves C for which the line of striction of the ruled surface is the locus of the centers of mean curvature corresponding to C ?

On S we take the lines of curvature parametric. Referred to the moving trihedral of S , the direction-cosines of the normal are $(0, 0, 1)$, and the variations in these are given by*

$$dX = qdu, \quad dY = -p_1dv, \quad dZ = 0.$$

Now the displacement of the central point on each generator of the ruled surface is orthogonal both to the normal and to its neighboring position. Hence we have

$$\delta z = 0, \quad qdu\delta x - p_1dv\delta y + \delta z = 0,$$

which reduce to

$$(1) \quad qdu(\xi du + zqdu) - p_1dv(\eta_1 dv - zp_1 dv) = 0.$$

If in (1) we assign a value to the ratio dv/du , this equation will determine the distance z to the line of striction on the ruled surface defined by this ratio; and if to z we assign a given value, equation (1) will determine the curves, (though not necessarily real), for which this assigned value of z is the distance to the lines of striction.

From (1) we have for the problem at hand,

* Eisenhart, *Differential Geometry of Curves and Surfaces*, pp. 166-174.