

ABSTRACT CAUCHY PROBLEMS OF THE ELLIPTIC TYPE

BY A. V. BALAKRISHNAN

Communicated by Einar Hille, June 13, 1958

Let A be the infinitesimal generator of a strongly continuous one-parameter semigroup $T(\xi)$, $0 < \xi$, of endomorphisms over a B -space X . Suppose it is required to find a function $u(t)$, $0 < t$, with values in X such that:

(i) $u(t)$, $u^1(t)$, \dots , $u^{n-1}(t)$ are absolutely continuous, $u^k(t)$ being the derivative of $u^{k-1}(t)$.

(ii) $u^n(t) = (-1)^{n+1}Au(t)$.

(iii) $\|u^k(t) - u_k\| \rightarrow 0$, as $t \rightarrow 0+$, $k = 0, \dots, n-1$.

We call this an abstract Cauchy problem of the elliptic type ($ACPE_n$). We prove:

THEOREM 1. *The $ACPE_n$ has at most one solution provided*

$$(H_1) \quad \int_1^\infty \|T(\xi)\| \xi^{-\sigma-1} d\xi < \infty \text{ for every } \sigma > 0.$$

THEOREM 2. *Let $n=2$. Let the semi-group $T(\xi)$ satisfy H_1 and let $u(t)$ be any solution of the $ACPE_2$ such that*

$$(H_2) \quad \limsup_{t \rightarrow \infty} t^{-1} \text{Log } \|u(t)\| \leq 0.$$

Then necessarily

$$(1) \quad u(t) = (t/2\pi^{1/2}) \int_0^\infty T(\xi) u_0 \xi^{-3/2} \exp(-t^2/4\xi) d\xi.$$

A slightly different but useful version of Theorem 2 is:

THEOREM 3. *Let $n=2$. Let the semi-group $T(\xi)$ satisfy H_1 . Let $u(t)$, $t > 0$, satisfy (i), (ii), but (iiia) below in place of (iii)*

$$(iiia) \quad \|u(t) - u_0\| \rightarrow 0 \text{ as } t \rightarrow 0+.$$

Then, if $u(t)$ satisfies H_2 in addition, $u(t)$ is again determined by (1). Moreover, if $\|T(\xi)\| \rightarrow 0$ as $\xi \rightarrow \infty$, then any such $u(t)$ has a similar property, viz.:

$$\|u(t)\| \rightarrow 0, \text{ as } t \rightarrow \infty.$$

Results similar in principle have been obtained for other values of n .