

STRUCTURE THEOREM FOR COMMUTATORS OF OPERATORS

BY ARLEN BROWN AND CARL PEARCY

Communicated by P. R. Halmos, July 13, 1964

If \mathcal{H} is a separable (complex) Hilbert space, and A is a (bounded, linear) operator on \mathcal{H} , then A is a *commutator* if there exist operators B and C on \mathcal{H} such that $A = BC - CB$. It was shown by Wintner [8] and also by Wielandt [7] that no nonzero scalar multiple of the identity operator I on \mathcal{H} is a commutator, and this was improved by Halmos [5] who showed that no operator of the form $\lambda I + C$ is a commutator, where $\lambda \neq 0$ and C is a compact operator. The purpose of this note is to announce the following theorem and give some indication of its proof. Details of the results described below will appear elsewhere [2].

THEOREM. *An operator A on a separable Hilbert space \mathcal{H} is a commutator if and only if A is not of the form $\lambda I + C$ where $\lambda \neq 0$ and C is a compact operator.*

This theorem furnishes the solution to several problems concerning commutators posed by Halmos in [4] and [5]. In particular it is interesting to note that the identity operator is the limit in the norm of commutators and that there exists a commutator whose spectrum consists of the number 1 alone.

INDICATION OF THE PROOF. We must show that every operator that is not of the form $\lambda I + C$, with $\lambda \neq 0$ and C compact, is a commutator. These operators fall naturally into two classes; viz., the class of compact operators, which was shown to consist entirely of commutators in [1], and the class consisting of all operators that cannot be written in the form $\lambda I + C$ for any scalar λ (0 or not) and compact C . We denote this latter class by (F) , and the first problem is to obtain a more usable characterization of the operators of this class. To this end we define for an arbitrary operator T on \mathcal{H} the function

$$\eta_T(x) = \|Tx - (Tx, x)x\|, \quad x \in \mathcal{H}, \|x\| = 1,$$

and denote by $\eta_T(\mathfrak{M})$ the supremum over the subspace $\mathfrak{M} \subset \mathcal{H}$ of $\eta_T(x)$.

PROPOSITION 1. *An operator T is of type (F) if and only if $\inf \eta_T(\mathfrak{M}) > 0$ where the infimum is taken over all cofinite-dimensional subspaces \mathfrak{M} of \mathcal{H} .*