

A NEW INVARIANT OF HOMOTOPY TYPE AND SOME DIVERSE APPLICATIONS

BY DANIEL H. GOTTLIEB¹

Communicated by J. Milnor, December 14, 1964

Let X be a connected, locally finite simplicial polyhedron. Let $X^{\mathbf{x}}$ be the space of maps from X to X with the compact-open topology. Let $x_0 \in X$ be taken as a base point in X , then the evaluation map $p: X^{\mathbf{x}} \rightarrow X$ defined by $p(f) = f(x_0)$ for $f \in X^{\mathbf{x}}$ is continuous. Now p induces the homomorphism

$$p_*: \pi_1(X^{\mathbf{x}}, 1_X) \rightarrow \pi_1(X, x_0),$$

where $1_X \in X^{\mathbf{x}}$ is the identity map. Hence $p_*\pi_1(X^{\mathbf{x}}, 1_X)$ is a subgroup of the fundamental group of (X, x_0) .

PROPOSITION 1. *$p_*\pi_1(X^{\mathbf{x}}, 1_X)$ considered as a subgroup of $\pi_1(X, x_0)$ is an invariant of homotopy type.*

In [2], this invariant is studied and theorems are obtained which bear on the study of $X^{\mathbf{x}}$, groups of homeomorphisms, homological group theory and knot theory. Most of these results come from the following theorem.

THEOREM 2. *Let X have the homotopy type of a compact, connected polyhedron with nonzero Euler-Poincaré number. Then $p_*\pi_1(X^{\mathbf{x}}, 1_X) = 0$.*

The proof of this employs Nielsen-Wecken fixed-point class theory ([1] and [5]).

Let $G(X)$ be the group of homeomorphisms of a manifold X , and let $G_0(X)$ be the isotropy group over x_0 . Then there is an exact sequence [3]

$$\cdots \rightarrow \pi_i(G_0(X), 1_X) \xrightarrow{i_*} \pi_i(G(X), 1_X) \xrightarrow{p'_*} \pi_i(X, x_0) \rightarrow \cdots,$$

where $p': G(X) \rightarrow X$ is the evaluation map.

COROLLARY 3. *Let X be as in Theorem 2. Then $p'_*\pi_1(G(X), 1_X) = 0$. In particular, if $\pi_2(X, x_0) = 0$, then $i_*: \pi_1(G_0(X), 1_X) \cong \pi_1(G(X), 1_X)$.*

This follows because $p'_*\pi_1(G(X), 1_X) \subseteq p_*\pi_1(X^{\mathbf{x}}, 1_X)$.

¹ This work was partially supported by NSF Grant 1908.