

## A STRUCTURE THEOREM FOR COMPLETE NONCOMPACT HYPERSURFACES OF NONNEGATIVE CURVATURE

BY H. WU<sup>1</sup>

Communicated by I. Singer, May 20, 1971

The convexity theorem of Sacksteder-van Heijenoort [4] states that if  $M$  is a  $C^\infty$   $n$ -dimensional ( $n > 1$ ) complete orientable Riemannian manifold of nonnegative sectional curvature with positive curvature at one point, then every isometric immersion  $x: M \rightarrow \mathbf{R}^{n+1}$  is an imbedding and  $x(M)$  bounds an open convex subset of  $\mathbf{R}^{n+1}$ ; furthermore  $M$  is diffeomorphic to either  $\mathbf{R}^n$  or  $S^n$  (unit  $n$ -sphere). The purpose of this note is to announce a structure theorem that complements the above result of Sacksteder and van Heijenoort. Full details will appear in a forthcoming monograph on convexity and rigidity of hypersurfaces.

**THEOREM.** *Let  $M$  be a  $C^\infty$  hypersurface in  $\mathbf{R}^{n+1}$  ( $n > 1$ ) which is complete, noncompact, orientable with nonnegative sectional curvature, which is in addition all positive at one point, then:*

(1) *The spherical image of  $M$  in the unit sphere  $S^n$  has a geodesically convex closure, which lies in a closed hemisphere.*

(2) *The total curvature of  $M$  (cf. Chern-Lashof [2]) does not exceed one.*

(3)  *$M$  is a pseudograph (see below for definition) over one of its tangent planes.*

(4)  *$M$  has infinite volume.*

**COROLLARY.** *Suppose the sectional curvature of  $M$  is in fact everywhere positive, then:*

(5) *The spherical map is a diffeomorphism onto a geodesically convex open subset of  $S^n$ . Consequently the spherical image lies in an open hemisphere.*

(6) *Coordinates in  $\mathbf{R}^{n+1}$  can be so chosen that  $M$  is tangent to the hyperplane  $x_{n+1} = 0$  at the origin, and there is a nonnegative strictly convex function (i.e. its Hessian is everywhere positive definite)  $f(x_1, \dots, x_n)$  defined in a convex domain of  $\{x_{n+1} = 0\}$  such that  $M$  is exactly the graph of  $f$ .*

**REMARKS.** (A) A  $C^\infty$  convex hypersurface  $M$  (i.e.  $M$  is the full boundary of an open convex set) in  $\mathbf{R}^{n+1}$  is said to form a *pseudograph*

---

AMS 1969 subject classifications. Primary 5374; Secondary 5375.

<sup>1</sup> Sloan Fellow. Also partially supported by the National Science Foundation.

Copyright © American Mathematical Society 1971