FINITE RICKART C*-ALGEBRAS

BY DAVID HANDELMAN AND JOHN LAWRENCE¹ Communicated by Barbara Osofsky, May 23, 1977

1. A C*-algebra is Rickart [1] if the right annihilator of each element t is generated by a projection 1 - RP(t). If in addition, $xx^* = 1$ implies $x^*x = 1$, the ring is called a finite Rickart C*-algebra. In this note we announce several new results on finite Rickart C*-algebras. Detailed proofs will appear elsewhere.

Two projections e and f in a Rickart C^* -algebra are *-equivalent ($e \stackrel{*}{\sim} f$) if there exists a w such that $ww^* = e$ and $w^*w = f$. Kaplansky asked whether left projections in a Rickart C^* -algebra were *-equivalent to right projections, that is, whether $RP(t) \stackrel{*}{\sim} LP(t)$ [5]. We have the following partial answer.

THEOREM 1 [2]. In a finite Rickart C^* -algebra, left projections are equivalent to right projections. In fact RP(t) and LP(t) are unitarily equivalent for each element t in the algebra.

A consequence of this is the following.

COROLLARY 2 [2]. A simple homomorphic image of a finite Rickart C^* -algebra is a finite AW*-factor.

THEOREM 3 [3]. If T is a finite Rickart C*-algebra that is either abelian or an $n \times n$ matrix ring over some ring for n > 1, then all matrix rings over T are also finite Rickart C*-algebras.

A finite Rickart C^* -algebra is a subdirect product of its simple homomorphic images.

THEOREM 4 [4]. In a finite Rickart C^* -algebra, the intersection of the maximal (two-sided) ideals is zero.

Applying Theorem 4 and Corollary 2, we have

COROLLARY 5 [4]. A finite Rickart C^* -algebra can be embedded in a finite AW^* -algebra.

2. Outline of the method. In [2] we show that a finite Rickart C^* -algebra T has an \aleph_0 -continuous (unit) regular quotient ring R to which the involution on T can be lifted. Certain questions concerning finite Rickart C^* -algebras can then be 'lifted' to this quotient ring. This is implemented by a close study

AMS (MOS) subject classifications (1970). Primary 16A30, 47L05.

¹Research supported by NRC Grant A4540 and University of Waterloo Research Grant 131-7052. Copyright © 1978, American Mathematical Society