

THE MAY-WIGNER STABILITY THEOREM FOR CONNECTED MATRICES

BY HAROLD M. HASTINGS¹

In 1972 R. M. May [4], following empirical results of M. R. Gardner and W. R. Ashby [2], sketched a proof of the following asymptotic stability theorem for large random linear differential systems:

$$(1) \quad dx/dt = Ax.$$

May used deep results of E. P. Wigner [7], the "semicircle law" for eigenvalues of random symmetric matrices; see also Mehta [6].

MAY-WIGNER STABILITY THEOREM. *Let B be an $n \times n$ matrix with $n^2 C$ ($0 < C < 1$) randomly located nonzero entries, each chosen independently from a symmetric distribution with variance α^2 . Let $A = B - I$, and let $P(\alpha, n, C)$ be the probability that the corresponding differential system (1) has a stable equilibrium at 0. Let $\epsilon > 0$. Then $P(\alpha, n, C) \rightarrow 1$ as $n \rightarrow \infty$ provided $\alpha^2 n C < 1 - \epsilon$; conversely, $P(\alpha, n, C) \rightarrow 0$ as $n \rightarrow \infty$ for $\alpha^2 n C > 1 + \epsilon$.*

This result provided a basis for studying the stability of neutral models in both cybernetics [2] and ecology (May [5] and references therein). Unfortunately Wigner's results are extremely complex, and may not apply to all random matrices (G. Sugihara, private communication, see also [6, p. 150]). We therefore sought a conceptually simpler proof.

We announce here a direct proof for matrices with connected underlying graphs. More precisely, the underlying graph of A (with one edge joining i and j if A_{ij} or A_{ji} is nonzero) is asymptotically almost surely connected if

$$C \geq (1 + \epsilon) \log n/n,$$

and asymptotically almost surely not connected if

$$C \leq (1 - \epsilon) \log n/n$$

for any fixed positive ϵ , (Bollobás [1, p. 143]). We assume the former condition holds; in particular the theorem holds for any constant C .

Received by the editors May 26, 1982.

¹Partially supported by NSF grant MCS8102053.

1980 *Mathematics Subject Classification*. Primary 15A52, 34D05, 82A99, 92A15, 92A17.