

- [K] T. Kato, *Non-stationary flows of viscous and ideal fluids in R^3* , J. Funct. Anal. **9** (1972), 296–305.
- [KM] S. Klainerman and A. Majda, *Singular limits of quasilinear hyperbolic systems with large parameters and the incompressible limit of compressible fluids*, Comm. Pure Appl. Math. **34** (1981), 481–524.
- [L] J. Leray, *Sur le mouvement d'un liquide visqueux emplissant l'espace*, Acta Math. **63** (1934), 193–248.
- [L-L] L. Landau and E. Lifschitz, *Fluid mechanics*, Addison-Wesley, New York, 1953.
- [M] A. Majda, *Compressible fluid flow and systems of conservation laws in several space variables*, Appl. Math. Sci., vol. 53, Springer-Verlag, New York, 1984.
- [M-dP] A. Majda and R. DiPerna, *Oscillations and concentrations in weak solutions of the incompressible fluid equations*, Comm. Math. Phys. **108** (1987), 667–689.
- [R.] D. Ruelle, *Large volume limit of distributions of characteristic exponents in turbulence*, Comm. Math. Phys. **87** (1982), 287–302.
- [S] J. Serrin, *The initial value problem for the Navier–Stokes equations*, Non-linear Problems (R. E. Langer, ed.), Univ. of Wisconsin Press, Madison, 1963.
- [T] R. Temam, *Navier–Stokes equations*, North-Holland, Amsterdam, 1977.
- [VF] M. J. Vishik and A. V. Fursikov, *Mathematical problems of statistical hydrodynamics*, Kluwer Acad. Publ., Dordrecht, Holland, 1988.
- [vW] W. von Wahl, *The equations of Navier–Stokes and abstract parabolic equations*, Vieweg and Sohn, Braunschweig and Wiesbaden, 1988.
- [Y] V. I. Yudovitch, *Non-stationary flow of an ideal incompressible liquid*, Zh. Vychisl. Mat. i Mat. Fiz. **3** (1963), 1032–1066.

PETER CONSTANTIN
UNIVERSITY OF CHICAGO

BULLETIN (New Series) OF THE
AMERICAN MATHEMATICAL SOCIETY
Volume 23, Number 2, October 1990
© 1990 American Mathematical Society
0273-0979/90 \$1.00 + \$.25 per page

Mathematical problems from combustion theory, by Jerrold Berbernes and David Eberly. Applied Mathematical Sciences, vol. 83, Springer-Verlag, New York, 1989, 177 pp., \$34.00. ISBN 0-387-97104-1

Combustion involves the liberation of energy by chemical reaction. Typically, the rate of energy release is a strongly sensitive function of temperature. Doubling the temperature, for example, may well increase the rate by a factor of ten thousand. Combustible materials, essentially inert at room temperature, can therefore ignite rapidly and explode when sufficiently heated.