Analytic Torsion and Holomorphic Determinant Bundles

II. Direct Images and Bott-Chern Forms

Jean-Michel Bismut¹, Henri Gillet²*, and Christophe Soulé³

¹ Département de Mathématique, Université Paris-Sud, Bâtiment 425, F-91405 Orsay Cedex, France

² Department of Mathematics, University of Illinois at Chicago, Chicago, Ill 60638 USA ³ C.N.R.S., L.A. 212 and IHES, 35, Route de Chartres, F-91440 Bures/Yvette, France

Abstract. In this paper, we derive the main properties of Kähler fibrations. We introduce the associated Levi-Civita superconnection to construct analytic torsion forms for holomorphic direct images. These forms generalize in any degree the analytic torsion of Ray and Singer. In the case of acyclic complexes of holomorphic Hermitian vector bundles, such forms are calculated by means of Bott-Chern classes.

Contents

Introduction	•	79
1. Kähler Fibrations		81
a) An Euclidean Connection on the Vertical Tangent Space of a Fibration		
b) Invariance Properties of $\nabla^{\mathbb{Z}}$		
c) Kähler Fibrations and the Levi-Civita Connection		
d) A Family of Dirac Operators		
e) A Holomorphic Hermitian Connection on Infinite Dimensional Bundles		
2. Double Transgression for Direct Images and the Heat Equation.		
a) Kähler Fibrations and the Levi-Civita Superconnection		
b) Construction of Chern Character Forms in P		
c) Number Operator and the Levi-Civita Superconnection		
d) Double Transgression of the Chern Character: The Infinitesimal Form		
e) Asymptotic Behavior of the First Transgressed Forms		
f) On Certain Identities Verified by the Levi-Civita Superconnection		
g) The Asymptotics of $\operatorname{Tr}_s[N_u \exp(-A_u^2)]$		108
h) Double Transgression of the Chern Character Forms		
i) The Case where (ξ, v) is Acyclic		116
References		125

Introduction

This is the second of a series of three papers devoted to the study of holomorphic determinant bundles and direct images. Parts I and III of this work will be referred

^{*} Supported by NSF Grant DMS 850248 and by the Sloan Foundation