

## ON A STRONGER VERSION OF WALLIS' FORMULA

V. R. RAO UPPULURI

**In Mathematical Statistics, estimation of parameters which index the probability density functions of random variables is an interesting area. The object of estimation is to look for estimators which have 'desirable' properties. It turns out that the lower bounds on the variances of estimators can be used to derive some inequalities. This is illustrated here in connection with Wallis' formula.**

We shall now cite some references where this idea is used. In the classic book on mathematical probability, Uspensky [10] remarks as follows: "There are many cases in which, by means of considerations belonging to the theory of probability, several identities or inequalities can be established whose direct proof sometimes involves considerable difficulty." He has exemplified this remark very beautifully in several contexts in the book. In 1955, Chassan [1] has given some inequalities involving trigonometric functions, obtained by comparing the variance of a minimum variance estimator with the variance of a less efficient estimator. In 1956, Gurland [4] has given an inequality satisfied by the Gamma function, which was also obtained by comparing the variances of two estimators by using the so-called Cramér-Rao lower bound for the variance of unbiased estimators. In 1959, Olkin [8] has given an extension of Gurland's inequality, by using the multivariate form of the probability density function used by Gurland [4]. In 1962, Gokhale [3] has given a different inequality for the Gamma function than that given by Gurland [4], by using an analogue of the Cramér-Rao lower bound derived by Rao [9].

In 1962, Mann [7] has given a beautiful application of Statistical Inference. By constructing the most powerful regions of a given size, he has illustrated how one can deduce the arithmetic-geometric mean inequality, Hölder inequality, and other well-known inequalities. In a humorous vein he remarks at the end: "Thus we have derived Hölder's inequality from the fact that we cannot increase our knowledge on the milk yield of cows by flipping a coin or by measuring the weight of herrings."

In 1956, Gurland [5] has also given another illustration of the use of Cramér-Rao lower bound for the variance of unbiased estimators, which yielded a closer approximation to  $\pi$ , than the so-called Wallis' result. In this paper we shall pursue this idea and give stronger versions of Wallis' Formula, by using the so-called Bhattacharya bounds for the variance of estimators which is an extension of the Cramér-Rao bound.