

## Research Article

# Compatible and Incompatible Nonuniqueness Conditions for the Classical Cauchy Problem

Josef Diblík<sup>1,2</sup> and Christine Nowak<sup>3</sup>

<sup>1</sup> Department of Mathematics and Descriptive Geometry, Faculty of Civil Engineering, University of Technology, 602 00 Brno, Czech Republic

<sup>2</sup> Department of Mathematics, Faculty of Electrical Engineering and Communication, Brno University of Technology, 616 00 Brno, Czech Republic

<sup>3</sup> Institute for Mathematics, University of Klagenfurt, 9020 Klagenfurt, Austria

Correspondence should be addressed to Christine Nowak, christine.nowak@uni-klu.ac.at

Received 21 September 2010; Revised 23 November 2010; Accepted 10 March 2011

Academic Editor: Yuri V. Rogovchenko

Copyright © 2011 J. Diblík and C. Nowak. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the first part of this paper sufficient conditions for nonuniqueness of the classical Cauchy problem  $\dot{x} = f(t, x)$ ,  $x(t_0) = x_0$  are given. As the essential tool serves a method which estimates the “distance” between two solutions with an appropriate Lyapunov function and permits to show that under certain conditions the “distance” between two different solutions vanishes at the initial point. In the second part attention is paid to conditions that are obtained by a formal inversion of uniqueness theorems of Kamke-type but cannot guarantee nonuniqueness because they are incompatible.

## 1. Introduction

Consider the initial value problem

$$\dot{x} = f(t, x), \quad x(t_0) = x_0, \quad (1.1)$$

where  $t_0 \in \mathbb{R}$ ,  $t \in J := [t_0, t_0 + a]$  with  $a > 0$ ,  $x, x_0 \in \mathbb{R}^n$  and  $f : J \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ .

In the first part (Section 2) we give sufficient conditions for nonuniqueness of the classical  $n$ -dimensional Cauchy problem (1.1). As the essential tool serves a method which estimates the “distance” between two solutions with an appropriate Lyapunov function and permits to show that under certain conditions the “distance” between two different solutions vanishes at the initial point. In the second part (Section 3) we analyze for the one-dimensional case a set of conditions that takes its origin in an inversion of the uniqueness theorem by Kamke (see, e.g., [1, page 56]) but cannot guarantee nonuniqueness since it contains an

inner contradiction. Several attempts were made to get nonuniqueness criteria by using conditions that are (in a certain sense) reverse uniqueness conditions of Kamke type. But this inversion process has to be handled very carefully. It can yield incompatible conditions. This is illustrated by a general set of conditions (in Theorems 3.2, 3.5 and 3.6) that would ensure nonuniqueness, but unfortunately they are inconsistent.

In this paper we study Cauchy problems where  $f$  is continuous at the initial point. Related results can be found in [1–5]. In literature there are several investigations for the discontinuous case [1, 6–13] with different qualitative behaviour.

## 2. Main Result

In the following let  $\mathbb{R}_+ := [0, \infty)$ ,  $b > 0$ ,  $\rho > 0$  and

$$S_\rho^n(x_0) := \{x \in \mathbb{R}^n : \|x - x_0\| < \rho\}, \quad (2.1)$$

where  $\|\cdot\|$  means the Euclidean norm.

*Definition 2.1.* We say that the initial value problem (1.1) has at least two different solutions on the interval  $J$  if there exist solutions  $\varphi(t)$ ,  $\psi(t)$  defined on  $J$  and  $\varphi \neq \psi$ .

The following notions are used in our paper (see, e.g., [14, pages 136 and 137]).

*Definition 2.2.* A function  $\varphi : [0, \rho) \rightarrow \mathbb{R}_+$  is said to belong to the class  $\mathcal{K}_\rho$  if it is continuous, strictly increasing on  $[0, \rho)$  and  $\varphi(0) = 0$ .

*Definition 2.3.* A function  $V : J \times S_\rho^n(0) \rightarrow \mathbb{R}_+$  with  $V(t, 0) \equiv 0$  is said to be positive definite if there exists a function  $\varphi \in \mathcal{K}_\rho$  such that the relation

$$V(t, x) \geq \varphi(\|x\|) \quad (2.2)$$

is satisfied for  $(t, x) \in J \times S_\rho^n(0)$ .

For the convenience of the reader we recall the definition of a uniformly Lipschitzian function with respect to a given variable.

*Definition 2.4.* A function  $V(t, \cdot) : S_\rho^n(0) \rightarrow \mathbb{R}_+$  is said to be Lipschitzian uniformly with respect to  $t \in J$  if for arbitrarily given  $x^* \in S_\rho^n(0)$  there exists a constant  $k = k(x^*)$  such that

$$\|V(t, x_1^*) - V(t, x_2^*)\| \leq k \|x_1^* - x_2^*\| \quad (2.3)$$

holds for every  $t \in J$  and for every  $x_1^*, x_2^*$  within a small neighbourhood of  $x^*$  in  $S_\rho^n(0)$ .

In [1, 15, 16] generalized derivatives of a Lipschitzian function along solutions of an associated differential system are analyzed. A slight modification of Theorem 4.3 [15, Appendix I] is the following lemma.

**Lemma 2.5.** Let  $V : J \times S_\rho^n(0) \rightarrow \mathbb{R}_+$  be continuous and let  $V(t, \cdot) : S_\rho^n(0) \rightarrow \mathbb{R}_+$  be Lipschitzian uniformly with respect to  $t \in J$ . Let  $x_1, x_2 : J \rightarrow S_\rho^n(0)$  be any two solutions of

$$\dot{x} = f(t, x), \quad (2.4)$$

where  $f : J \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is a continuous function. Then for the upper right Dini derivative the equality

$$\begin{aligned} & D^+V(t, x_2(t) - x_1(t)) \\ & := \limsup_{h \rightarrow 0^+} \frac{1}{h} [V(t+h, x_2(t+h) - x_1(t+h)) - V(t, x_2(t) - x_1(t))] \\ & = \limsup_{h \rightarrow 0^+} \frac{1}{h} [V(t+h, x_2(t) - x_1(t) + h(f(t, x_2(t)) - f(t, x_1(t)))) - V(t, x_2(t) - x_1(t))] \end{aligned} \quad (2.5)$$

holds.

In the proof of Theorem 2.8 we require the following lemmas which are slight adaptations of Theorem 1.4.1 [14, page 15] and Theorem 1.3.1 [1, page 10] for the left side of the initial point.

**Lemma 2.6.** Let  $E$  be an open  $(t, u)$ -set in  $\mathbb{R}^2$ , let  $g : E \rightarrow \mathbb{R}$  be a continuous function, and let  $u$  be the unique solution of

$$\dot{u} = g(t, u), \quad u(t_2) = u_2, \quad (2.6)$$

to the left with  $t_2 > t_0$ ,  $(t_2, u_2) \in E$ . Further, we assume that the scalar continuous function  $m : (t_0, t_2] \rightarrow \mathbb{R}$  with  $(t, m(t)) \in E$  satisfies  $m(t_2) \leq u(t_2)$  and

$$D^+m(t) \geq g(t, m(t)), \quad t_0 < t \leq t_2. \quad (2.7)$$

Then

$$m(t) \leq u(t) \quad (2.8)$$

holds as far as the solution  $u$  exists left of  $t_2$  in  $(t_0, t_2]$ .

**Lemma 2.7.** Let  $S := \{(t, x) : t_0 - a \leq t \leq t_0, |x - x_0| \leq b\}$  and  $f : S \rightarrow \mathbb{R}$  be continuous and nondecreasing in  $x$  for each fixed  $t$  in  $[t_0 - a, t_0]$ . Then, the initial value problem (1.1) has at most one solution in  $[t_0 - a, t_0]$ .

**Theorem 2.8** (main result). Suppose that

(i)  $f : J \times S_b^n(x_0) \rightarrow \mathbb{R}^n$  is a continuous function such that

$$M := \sup\{\|f(t, x)\| : t \in J, x \in S_b^n(x_0)\} < \frac{b}{a}. \quad (2.9)$$

Let  $x_1$  be a solution of problem (1.1) on  $J$ . Let, moreover, there exist numbers  $t_1 \in (t_0, t_0+a]$ ,  $r \in (0, 2b)$  and continuous functions  $g : (t_0, t_1] \times \mathbb{R}_+ \rightarrow \mathbb{R}$ ,  $V : [t_0, t_1] \times S_r^n(0) \rightarrow \mathbb{R}_+$  such that

(ii)  $g$  is nondecreasing in the second variable, and the problem

$$\dot{u} = g(t, u), \quad \lim_{t \rightarrow t_0^+} u(t) = 0 \quad (2.10)$$

has a positive solution  $u^*$  on  $(t_0, t_1]$ ;

(iii)  $V$  is positive definite and  $V(t, \cdot) : S_r^n(0) \rightarrow \mathbb{R}_+$  is Lipschitzian uniformly with respect to  $t \in J$ ;

(iv) for  $t_0 < t \leq t_1$ ,  $\|y - x_1(t)\| < r$ , the inequality

$$\dot{V}(t, y - x_1(t)) \geq g(t, V(t, y - x_1(t))) \quad (2.11)$$

holds where

$$\begin{aligned} & \dot{V}(t, y - x_1(t)) \\ & := \limsup_{h \rightarrow 0^+} \frac{1}{h} [V(t+h, y - x_1(t) + h[f(t, y) - f(t, x_1(t))]) - V(t, y - x_1(t))]. \end{aligned} \quad (2.12)$$

Then the set of different solutions of problem (1.1) on interval  $J$  has the cardinality of the continuum.

*Remark 2.9.* If condition (i) is fulfilled then, as it is well known, problem (1.1) is globally solvable and every global solution admits the estimate

$$\|x(t) - x_0\| \leq M(t - t_0), \quad t \in J. \quad (2.13)$$

Moreover, for any local solution  $x_*$  of problem (1.1), defined on some interval  $[t_0, t_1] \subset J$ , there exists a global solution  $x$  of that problem such that  $x(t) = x_*(t)$  for  $t \in [t_0, t_1]$ .

*Remark 2.10.* For the case  $M = 0$  the initial value problem is unique and the assumptions of Theorem 2.8 cannot be satisfied. Therefore, without loss of generality, we assume  $M > 0$  in the proof below.

*Proof.* At first we show that (1.1) has at least two different solutions on  $[t_0, t_1^*]$ , where  $t_1^* \leq t_1$ ,  $t_1^* \leq t_0 + \min\{a, b/(3M)\}$  is sufficiently close to  $t_0$ . We construct a further solution of (1.1) by finding a point  $(t_2, x_2)$  not lying on the solution  $x_1(t)$  and starting from this point backwards to the initial point  $(t_0, x_0)$ .

First we show that there exist values  $t_2$  and  $x_2$ ,  $t_0 < t_2 \leq t_1^*$ ,  $\|x_2 - x_0\| \leq 2b/3$  such that

$$u^*(t_2) = V(t_2, x_2 - x_1(t_2)) \quad (2.14)$$

holds for the nontrivial solution  $u^*(t)$  of  $\dot{u} = g(t, u)$ . From Lemma 2.7 it follows that  $u^*(t)$  is determined uniquely to the left by the initial data  $(t_2, u^*(t_2))$ . We consider the  $\varepsilon$ -tubes

$$S(\varepsilon) := \{(t, x) : t_0 \leq t \leq t_1^*, \|x - x_1(t)\| = \varepsilon\} \quad (2.15)$$

for  $\varepsilon > 0$  around the solution  $x_1(t)$ . There exists  $\varepsilon_1 > 0$  such that  $S(\varepsilon)$  with  $0 < \varepsilon \leq \varepsilon_1 < r$  is contained in the set

$$\left\{ (t, x) : t_0 \leq t \leq t_1^*, \|x - x_0\| \leq \frac{2b}{3} \right\}. \quad (2.16)$$

For  $0 \leq \delta \leq \varepsilon_1$ ,  $t \in [t_0, t_1^*]$  we define

$$\begin{aligned} \Psi(\delta, t) &:= \max_{\|x - x_1(t)\| = \delta} V(t, x - x_1(t)), \\ \Psi(\delta) &:= \max_{t \in [t_0, t_1^*]} \Psi(\delta, t) \equiv \max_{(t, x) \in S(\delta)} V(t, x - x_1(t)). \end{aligned} \quad (2.17)$$

The function  $\Psi(\delta, t)$  is continuous in  $t$  for  $t_0 \leq t \leq t_1^*$ . Since  $\lim_{\delta \rightarrow 0} \Psi(\delta) = 0$ , there exists a  $\delta_2$ ,  $0 < \delta_2 \leq \min\{\varepsilon_1, b/3\}$ , such that  $\Psi(\delta_2) \leq u^*(t_1^*)$ . It is clear that inequalities

$$\Psi(\delta_2, t_1^*) \leq \Psi(\delta_2) \leq u^*(t_1^*) \quad (2.18)$$

and (due to positive definiteness of  $V$ )

$$\Psi(\delta_2, t_0) > 0 = \lim_{t \rightarrow t_0^+} u^*(t) \quad (2.19)$$

hold. We define a function

$$\omega(t) := \Psi(\delta_2, t) - u^*(t), \quad (2.20)$$

continuous on  $[t_0, t_1^*]$ . Taking into account inequalities  $\omega(t_0) > 0$  and  $\omega(t_1^*) \leq 0$  we conclude that there exists  $t_2$ ,  $t_0 < t_2 \leq t_1^*$ , with

$$\Psi(\delta_2, t_2) = u^*(t_2). \quad (2.21)$$

The value  $\Psi(\delta_2, t_2)$  is taken by  $V(t_2, x - x_1(t_2))$  at a point  $x = x_2$  such that  $\|x_2 - x_1(t_2)\| = \delta_2$  and clearly (in view of the construction)  $x_2 \neq x_1(t_2)$ . The above statement is proved and (2.14) is valid for  $(t_2, x_2)$  determined above.

Now consider the initial value problem

$$\dot{x} = f(t, x), \quad x(t_2) = x_2. \quad (2.22)$$

Obviously  $t_2 - t_0 \leq b/(3M)$  since

$$0 < t_2 - t_0 \leq t_1^* - t_0 \leq \min\left\{a, \frac{b}{3M}\right\} \leq \frac{b}{3M} \quad (2.23)$$

and  $\|x_2 - x_0\| \leq 2b/3$  because

$$\begin{aligned} \|x_2 - x_0\| &= \|x_2 - x_1(t_2) + x_1(t_2) - x_0\| \\ &\leq \|x_2 - x_1(t_2)\| + \|x_1(t_2) - x_0\| = \delta_2 + \left\| \int_{t_0}^{t_2} f(s, x_1(s)) ds \right\| \\ &\leq \delta_2 + M(t_2 - t_0) \leq \delta_2 + M \frac{b}{3M} = \delta_2 + \frac{b}{3} \leq \frac{2b}{3}. \end{aligned} \quad (2.24)$$

Peano's theorem implies that there exists a solution  $x_2(t)$  of problem (2.22) on  $t_0 \leq t \leq t_2$ . We will show that  $x_2(t_0) = x_0$ . Set

$$m(t) := V(t, x_2(t) - x_1(t)). \quad (2.25)$$

Note that  $m(t_2) = u^*(t_2)$ . Lemma 2.5 and condition (iv) imply

$$\begin{aligned} D^+ m(t) &:= \limsup_{h \rightarrow 0^+} \frac{m(t+h) - m(t)}{h} \\ &= D^+ V(t, x_2(t) - x_1(t)) \\ &= \dot{V}(t, x_2(t) - x_1(t)) \geq g(t, V(t, x_2(t) - x_1(t))) = g(t, m(t)) \end{aligned} \quad (2.26)$$

for  $t_0 < t \leq t_2$ .

Applying Lemma 2.6 we get  $m(t) \leq u^*(t)$  for  $t_0 < t \leq t_2$ . As  $m(t) \geq 0$  for  $t_0 < t \leq t_2$  and  $m$  is continuous at  $t_0$ , we find  $m(t_0) = 0$ . Therefore we have  $x_2(t_0) = x_1(t_0) = x_0$  and, as noted above,  $x_2(t_2) = x_2 \neq x_1(t_2)$ . Thus problem (1.1) has two different solutions.

According to the well-known Kneser theorem [17, Theorem 4.1, page 15] the set of solutions of problem (1.1) either consists of one element or has the cardinality of the continuum. Consequently, if problem (1.1) has two different solutions on interval  $[t_0, t_1^*]$  and condition (i) is satisfied, then the set of different solutions of problem (1.1) on interval  $J$  has the cardinality of the continuum. The proof is completed.  $\square$

*Remark 2.11.* Note that in the scalar case with  $V(t, x) := |x|$  condition (2.11) has the form

$$(f(t, y) - f(t, x_1(t))) \cdot \text{sign}(y - x_1(t)) \geq g(t, |y - x_1(t)|). \quad (2.27)$$

*Example 2.12.* Consider for  $a = 0.1$ ,  $b = 1$ ,  $t_0 = 0$  and  $x_0 = 0$  the scalar differential equation

$$\dot{x} = f(t, x) := \begin{cases} 2x^{1/3} - \frac{1}{2} \cdot t^{1/2} \cdot \sin \frac{|x|}{t} & \text{if } t \neq 0, \\ 2x^{1/3} & \text{if } t = 0, \end{cases} \quad (2.28)$$

with the initial condition  $x(0) = 0$ . Let us show that the set of different solutions of this problem on interval  $J$  has the cardinality of  $\mathbb{R}$ . Obviously we can set  $x_1(t) \equiv 0$ . Put

$$g(t, u) := 2u^{1/3} - \frac{1}{2} \cdot t^{1/2}, \quad u^*(t) := t^{3/2}, \quad V(t, x) := |x|. \quad (2.29)$$

Conditions (i), (ii), and (iii) are satisfied. Let us verify that the last condition (iv) is valid, too. We get

$$\begin{aligned} \dot{V}(t, y - x_1(t)) &= \dot{V}(t, y) = (\text{sign } y) \cdot \left[ 2y^{1/3} - \frac{1}{2} \cdot t^{1/2} \cdot \sin \frac{|y|}{t} \right] \\ &\geq 2|y|^{1/3} - \frac{1}{2} \cdot t^{1/2} = 2V(t, y)^{1/3} - \frac{1}{2} \cdot t^{1/2} = g(t, V(t, y)) \\ &= g(t, V(t, y - x_1(t))). \end{aligned} \quad (2.30)$$

Thus, all conditions of Theorem 2.8 hold and, consequently, the set of different solutions on  $J$  of given problem has the cardinality of  $\mathbb{R}$ .

### 3. Incompatible Conditions

In this section we show that the formulation of condition (iv) in Theorem 2.8 without knowledge of a solution of the Cauchy problem can lead to an incompatible set of conditions. In the proof of Theorem 3.2 for the one-dimensional case we use the following result given by Nekvinda [18, page 1].

**Lemma 3.1.** *Let  $D \subset \mathbb{R}^2$  and let  $f : D \rightarrow \mathbb{R}$  be a continuous function in  $D$ . Let equation*

$$\dot{x} = f(t, x) \quad (3.1)$$

*has the property of left uniqueness. For any  $t_0 \in \mathbb{R}$  let  $A$  be the set of all  $x_0 \in \mathbb{R}$  such that  $(t_0, x_0) \in D$  and, for some  $\varepsilon > 0$ , the initial-value problem (1.1) has more than one solution in the interval  $[t_0, t_0 + \varepsilon)$ . Then  $A$  is at most countable.*

**Theorem 3.2.** *The set of conditions (i)–(iv):*

- (i)  $f : R_0 \rightarrow \mathbb{R}$  with  $R_0 := \{(t, x) \in J \times \mathbb{R}, |x - x_0| \leq b\}$  is continuous;

- (ii)  $g : (t_0, t_0 + a] \times (0, \infty) \rightarrow \mathbb{R}_+$  is continuous, nondecreasing in the second variable, and has the following property: there exists a continuous function  $u^*(t)$  on  $J$ , which satisfies the differential equation

$$\dot{u}(t) = g(t, u) \quad (3.2)$$

for  $t_0 < t \leq t_0 + a$  with  $u^*(t_0) = 0$  and does not vanish for  $t \neq t_0$ ;

- (iii)  $V : J \times S_{2b}^1(0) \rightarrow \mathbb{R}_+$  is continuous, positive definite, and Lipschitzian uniformly with respect to  $t \in J$ ;

- (iv) for  $t_0 < t \leq t_0 + a$ ,  $|x - x_0| \leq b$ ,  $|y - x_0| \leq b$ ,  $x \neq y$ ,

$$\dot{V}(t, x - y) \geq g(t, V(t, x - y)), \quad (3.3)$$

where we define

$$\dot{V}(t, x - y) := \limsup_{h \rightarrow 0^+} \frac{1}{h} [V(t + h, x - y + h[f(t, x) - f(t, y)]) - V(t, x - y)] \quad (3.4)$$

contains a contradiction.

*Proof.* Any initial value problem

$$\dot{x} = f(t, x), \quad x(t_0) = x^* \quad (3.5)$$

with  $|x^* - x_0| \leq b$  has at least two different solutions due to Theorem 2.8. Thus we have an uncountable set of nonuniqueness points. We show that solutions passing through different initial points are left unique. Suppose that it does not hold. Let  $x_1(t)$  be a solution starting from  $(t_0, x_1)$ , and let  $x_2(t)$  be a solution starting from  $(t_0, x_2)$  with  $x_2 \neq x_1$ . If we assume that these solutions cross at a point  $t_1 > t_0$  and if we set

$$m(t) := V(t, x_1(t) - x_2(t)) \quad (3.6)$$

then  $m(t_0) > 0$ ,  $m(t_1) = 0$ . Therefore there exists a point  $t \in (t_0, t_1)$  such that (we apply Lemma 2.5)

$$D^+ m(t) = D^+ V(t, x_1(t) - x_2(t)) = \dot{V}(t, x_1(t) - x_2(t)) < 0, \quad (3.7)$$

in contradiction to (3.3). Thus we obtain left uniqueness. From Lemma 3.1 we conclude in contrast to the above conclusion that the set of nonuniqueness points  $(t_0, x^*)$  can be at most countable.  $\square$

In [1, Theorem 1.24.1, page 99] the following nonuniqueness result (see [14, Theorem 2.2.7, page 55], too) is given which uses an inverse Kamke's condition (condition (3.9) below).

**Theorem 3.3.** Let  $g(t, u)$  be continuous on  $0 < t \leq a$ ,  $0 \leq u \leq 2b$ ,  $g(t, 0) \equiv 0$ , and  $g(t, u) > 0$  for  $u > 0$ . Suppose that, for each  $t_1$ ,  $0 < t_1 < a$ ,  $u(t) \not\equiv 0$  is a differentiable function on  $0 < t < t_1$ , and continuous on  $0 \leq t < t_1$  for which  $\dot{u}_+(0)$  exists,

$$\begin{aligned} \dot{u} &= g(t, u), & 0 < t < t_1, \\ u(0) &= \dot{u}_+(0) = 0. \end{aligned} \quad (3.8)$$

Let  $f \in C[R_0, \mathbb{R}]$ , where  $R_0 : 0 \leq t \leq a$ ,  $|x| \leq b$ , and, for  $(t, x), (t, y) \in R_0$ ,  $t \neq 0$ ,

$$|f(t, x) - f(t, y)| \geq g(t, |x - y|). \quad (3.9)$$

Then, the scalar problem  $\dot{x} = f(t, x)$ ,  $x(0) = 0$  has at least two solutions on  $0 \leq t \leq a$ .

*Remark 3.4.* In the proof of Theorem 3.3 at first  $f(t, 0) = 0$  is assumed. Putting  $y = 0$  in (3.9) leads to the inequality

$$|f(t, x)| \geq g(t, |x|). \quad (3.10)$$

As  $f(t, x)$  is continuous and  $g(t, u) > 0$  for  $u > 0$  it follows that  $f(t, x)$  must have constant sign for each of the half planes  $x > 0$  and  $x < 0$ . For the upper half plane this implies that

$$\begin{aligned} f(t, x) &\geq g(t, x), \\ f(t, x) &\leq -g(t, x). \end{aligned} \quad (3.11)$$

For the first inequality nonuniqueness is shown in [1]. But a similar argumentation cannot be used for the second inequality as the following example in [5] shows. We consider the initial value problem  $\dot{x} = f(t, x)$ ,  $x(0) = 0$ , with

$$f(t, x) = \begin{cases} -\sqrt{x} & \text{if } x \geq 0 \\ \sqrt{-x} & \text{if } x < 0 \end{cases} \quad (3.12)$$

and  $g(t, u) := \sqrt{u}$ . Thus inequality  $|f(t, x)| = \sqrt{|x|} \geq g(t, |x|)$  holds. In the upper half-plane we have  $f(t, x) \leq -g(t, x)$ . The function  $u(t) = t^2/4$  is a nontrivial solution of the comparison equation. Therefore all assumptions are fulfilled, but the initial value problem has at most one solution because of Theorem 1.3.1 [1, page 10].

The next theorem analyzes in the scalar case (for  $(t_0, x_0) = (0, 0)$ ) that even fulfilling a rather general condition (see condition (3.14) in the following theorem) cannot ensure nonuniqueness since the set of all conditions contains an inner contradiction. The proof was motivated by the paper [5].

**Theorem 3.5.** *There exists no system of three functions  $f$ ,  $g$ , and  $V$  satisfying the following suppositions:*

- (i)  $f : R_0 \rightarrow \mathbb{R}$  with  $R_0 := \{(t, x) \in \mathbb{R} \times \mathbb{R}, 0 \leq t \leq a, 0 \leq x \leq b\}$  is a continuous function;
- (ii) the continuous function  $g : (0, a] \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ ,  $g(t, 0) := 0$  if  $t \in (0, a]$ , has the following property: there exists a continuously differentiable function  $u^*(t)$  on  $0 \leq t \leq a$ , satisfying the differential equation

$$\dot{u} = g(t, u) \quad (3.13)$$

for  $0 < t \leq a$  such that  $u^*(0) = 0$  and  $u^*(t) > 0$  for  $t \neq 0$ ;

- (iii) the continuous function  $V : [0, a] \times S_b^1(0) \rightarrow \mathbb{R}_+$  is positive definite, and for all  $0 < t \leq a$ ,  $0 < x < b$  continuously differentiable;
- (iv) for  $0 < t \leq a$ ,  $0 < y < x \leq b$ ,

$$\dot{V}(t, x - y) \geq g(t, V(t, x - y)) \geq 0, \quad (3.14)$$

where we define

$$\dot{V}(t, x - y) := V'_1(t, x - y) + V'_2(t, x - y) \cdot [f(t, x) - f(t, y)] \quad (3.15)$$

and subscript indices denote the derivative with respect to the first and second argument, respectively;

- (v) there exist a positive constant  $\vartheta$  and a function  $\xi : (0, b] \rightarrow (0, \infty)$  such that for  $0 < t \leq a$  and  $0 < x \leq b$

$$\begin{aligned} 0 \leq V'_1(t, x) \leq \vartheta \cdot \xi(x), \quad 0 < V'_2(t, x) \leq \vartheta \cdot \frac{\xi(x)}{x}, \\ V(t, x) \geq \xi(x); \end{aligned} \quad (3.16)$$

- (vi) for  $t \in [0, a]$  and  $x, y$  with  $0 < y < x \leq b$  the inequality

$$f(t, x) - f(t, y) \geq 0 \quad (3.17)$$

holds.

*Proof.* Let us show that the above properties are not compatible. For fixed numbers  $x, y$  with  $0 < y < x \leq b$  consider the auxiliary function

$$F(t) := \frac{f(t, x) - f(t, y)}{x - y} + 1, \quad t \in [0, a]. \quad (3.18)$$

Clearly,  $F$  is continuous and assumes a (positive) maximum. Set

$$K = \max_{[0,a]} F(t) \geq 1. \quad (3.19)$$

If the function  $g$  fulfills the inequality

$$g(t, u) \leq \Lambda \cdot u \quad (3.20)$$

with a positive constant  $\Lambda$  in a domain  $0 < t \leq A \leq a$ ,  $0 \leq u \leq B$ ,  $B > 0$ , then the initial value problem

$$\dot{u} = g(t, u), \quad u(0) = 0 \quad (3.21)$$

has the unique trivial solution  $u = 0$ . Really, since  $u^*(t) > 0$  for  $t \in (0, a]$ , by integrating inequality

$$\frac{\dot{u}^*(t)}{u^*(t)} \leq \Lambda \quad (3.22)$$

with limits  $t, A^* \in (0, A)$  we get

$$u^*(A^*) \leq u^*(t) \exp[\Lambda(A^* - t)] \quad (3.23)$$

and for  $t \rightarrow 0^+$

$$u^*(A^*) \leq 0 \quad (3.24)$$

which contradicts positivity of  $u^*$ . Therefore problem (3.21) has only the trivial solution. Hence, there exist a sequence  $\{(t_n, u_n)\}$  with  $t_n \in (0, a]$ ,  $u_n > 0$ ,  $\lim_{n \rightarrow \infty} (t_n, u_n) = (0, 0)$  and a sequence  $\{\lambda_n\}$ ,  $\lambda_n > 0$ , with  $\lim_{n \rightarrow \infty} \lambda_n = \infty$  such that the inequality

$$g(t_n, u_n) > \lambda_n u_n \quad (3.25)$$

holds for every  $n$ . Consider now the relation

$$V(t, x) = 0. \quad (3.26)$$

Due to the properties of  $V$  we conclude that for all sufficiently small positive numbers  $t_n, u_n$  (i.e., for all sufficiently large  $n$ ) there exists a (sufficiently small and positive) number  $\tilde{u}_n$  such that the equation

$$V(t_n, x) = u_n \quad (3.27)$$

has the solution  $x = \tilde{u}_n$ . Thus a sequence  $\{\tilde{u}_n\}$  with  $\tilde{u}_n > 0$  and  $\lim_{n \rightarrow \infty} \tilde{u}_n = 0$  corresponds to the sequence  $\{(t_n, u_n)\}$ . For every  $n$  define a number  $j_n$  as

$$j_n = \left\lceil \frac{x - y}{\tilde{u}_n} - 1 \right\rceil, \quad (3.28)$$

where  $\lceil \cdot \rceil$  is the ceiling function. Without loss of generality we can suppose that

$$\frac{x - y}{\tilde{u}_n} > 4. \quad (3.29)$$

Obviously,

$$\frac{x - y}{\tilde{u}_n} - 1 \leq j_n < \frac{x - y}{\tilde{u}_n}. \quad (3.30)$$

Moreover, without loss of generality we can suppose that for every sufficiently large  $n$  the inequality

$$\lambda_n > 2\mathfrak{D}K \quad (3.31)$$

holds. Set

$$\begin{aligned} x_0 &:= y, \\ x_1 &:= y + \tilde{u}_n, \\ x_2 &:= y + 2\tilde{u}_n, \\ &\vdots \\ x_{j_n} &:= y + j_n \cdot \tilde{u}_n, \\ x_{j_n+1} &:= x. \end{aligned} \quad (3.32)$$

Consider for all sufficiently large  $n$  the expression

$$\mathcal{E}_n := j_n V_1'(t_n, \tilde{u}_n) + V_2'(t_n, \tilde{u}_n) \cdot [f(t_n, x) - f(t_n, y)]. \quad (3.33)$$

Then

$$\begin{aligned}
 \mathcal{E}_n &= j_n V_1'(t_n, \tilde{u}_n) + V_2'(t_n, \tilde{u}_n) \cdot \sum_{i=1}^{j_n+1} [f(t_n, x_i) - f(t_n, x_{i-1})] \\
 &= j_n V_1'(t_n, \tilde{u}_n) + V_2'(t_n, \tilde{u}_n) \cdot \sum_{i=1}^{j_n} [f(t_n, x_i) - f(t_n, x_{i-1})] + [f(t_n, x) - f(t_n, x_{j_n})] \\
 &\geq [\text{due to (vi)}] \geq j_n V_1'(t_n, \tilde{u}_n) + V_2'(t_n, \tilde{u}_n) \cdot \sum_{i=1}^{j_n} [f(t_n, x_i) - f(t_n, x_{i-1})] \\
 &= [\text{due to (iv) and (v)}] = j_n V_1'(t_n, \tilde{u}_n) + V_2'(t_n, \tilde{u}_n) \cdot j_n \left[ \frac{-V_1'(t_n, \tilde{u}_n) + \dot{V}(t_n, \tilde{u}_n)}{V_2'(t_n, \tilde{u}_n)} \right] \\
 &\geq [\text{due to (iv)}] \\
 &\geq j_n V_1'(t_n, \tilde{u}_n) + V_2'(t_n, \tilde{u}_n) \cdot j_n \left[ \frac{-V_1'(t_n, \tilde{u}_n) + g(t_n, V(t_n, \tilde{u}_n))}{V_2'(t_n, \tilde{u}_n)} \right] \\
 &= j_n \cdot g(t_n, V(t_n, \tilde{u}_n)) = [\text{due to (3.27)}] = j_n \cdot g(t_n, u_n) \geq [\text{due to (3.25)}] \tag{3.34} \\
 &\geq j_n \lambda_n u_n \geq [\text{due to (3.31)}] \geq j_n u_n \cdot 2\vartheta K \geq [\text{due to (3.30)}] \\
 &\geq \left( \frac{x-y}{\tilde{u}_n} - 1 \right) u_n \cdot 2\vartheta K \\
 &= \left( \frac{x-y}{\tilde{u}_n} - 1 \right) V(t_n, \tilde{u}_n) \cdot 2\vartheta K \\
 &\geq [\text{due to (3.16)}] \geq \left( \frac{x-y}{\tilde{u}_n} - 1 \right) \xi(\tilde{u}_n) \cdot 2\vartheta K \\
 &= (x-y-\tilde{u}_n) \cdot \frac{\xi(\tilde{u}_n)}{\tilde{u}_n} \cdot 2\vartheta K \\
 &\geq [\text{due to (3.29)}] \geq \frac{3}{4} \cdot (x-y) \cdot 2\vartheta K \cdot \frac{\xi(\tilde{u}_n)}{\tilde{u}_n} \\
 &= \frac{3}{2} \cdot (x-y) \cdot \vartheta K \cdot \frac{\xi(\tilde{u}_n)}{\tilde{u}_n} > 0.
 \end{aligned}$$

Estimating the expression  $\mathcal{E}_n$  from above we get (see (3.32))

$$\begin{aligned}
 \mathcal{E}_n &\leq \frac{x-y}{\tilde{u}_n} V_1'(t_n, \tilde{u}_n) + V_2'(t_n, \tilde{u}_n) \cdot [f(t_n, x) - f(t_n, y)] \\
 &\leq [\text{due to (v)}] \tag{3.35} \\
 &\leq \frac{x-y}{\tilde{u}_n} \cdot \vartheta \cdot \xi(\tilde{u}_n) + \vartheta \cdot \frac{\xi(\tilde{u}_n)}{\tilde{u}_n} \cdot (K-1)(x-y) = \vartheta \cdot \frac{\xi(\tilde{u}_n)}{\tilde{u}_n} \cdot K(x-y).
 \end{aligned}$$

These two above estimations yield

$$0 < \frac{3}{2} \cdot (x - y) \cdot \vartheta K \cdot \frac{\xi(\tilde{u}_n)}{\tilde{u}_n} \leq \xi_n \leq (x - y) \cdot \vartheta K \cdot \frac{\xi(\tilde{u}_n)}{\tilde{u}_n}, \quad (3.36)$$

in contrast to  $(3/2) \not\leq 1$ . Since the initially taken points  $x$  and  $y$ ,  $0 < y < x$ , can be chosen arbitrarily close to zero, the theorem is proved.  $\square$

The following result is a consequence of Theorem 3.5 if  $V(t, x) := |x|$ ,  $\xi(x) := x$  and  $\vartheta = 1$ . Condition (3.38) below was discussed previously in [5].

**Theorem 3.6.** *There exists no system of two functions  $f$  and  $g$  satisfying the following suppositions:*

- (i)  $f : R_0 \rightarrow \mathbb{R}$  with  $R_0 := \{(t, x) \in \mathbb{R} \times \mathbb{R}, 0 \leq t \leq a, 0 \leq x \leq b\}$  is a continuous function;
- (ii) the continuous function  $g : (0, a] \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ ,  $g(t, 0) := 0$  if  $t \in (0, a]$ , has the following property: there exists a continuously differentiable function  $u^*(t)$  on  $0 \leq t \leq a$ , satisfying the differential equation

$$\dot{u}(t) = g(t, u) \quad (3.37)$$

for  $0 < t \leq a$  such that  $u^*(0) = 0$  and  $u^*(t) > 0$  for  $t \neq 0$ ;

- (iii) for  $0 < t \leq a$ ,  $0 < y < x \leq b$

$$f(t, x) - f(t, y) \geq g(t, x - y) \geq 0; \quad (3.38)$$

- (iv) for  $0 < y < x \leq b$  the inequality  $f(0, x) - f(0, y) \geq 0$  holds.

*Remark 3.7.* Let us note that in the singular case, that is, when we permit that the function  $f(t, x)$  is not continuous at  $t = 0$ , the given sets of conditions in Theorems 3.5 and 3.6 can be compatible. This can be seen from the proof where the continuity of  $f$  is substantial. Such singular case was considered in [13].

## Acknowledgments

This research was supported by the Grants P201/11/0768 and P201/10/1032 of the Czech Grant Agency (Prague), by the Project FEKT-S-11-2(921) and by the Council of Czech Government MSM 00216 30529.

## References

- [1] R. P. Agarwal and V. Lakshmikantham, *Uniqueness and Nonuniqueness Criteria for Ordinary Differential Equations*, vol. 6 of *Series in Real Analysis*, World Scientific Publishing, River Edge, NJ, USA, 1993.
- [2] J. Kalas, "General nonuniqueness theorem for ordinary differential equations," *Dynamics of Continuous, Discrete and Impulsive Systems*, vol. 3, no. 1, pp. 97–111, 1997.
- [3] J. Kalas, "Nonuniqueness for the solutions of ordinary differential equations," *Czechoslovak Mathematical Journal*, vol. 29, no. 1, pp. 105–112, 1979.

- [4] J. Kalas, "Nonuniqueness results for ordinary differential equations," *Czechoslovak Mathematical Journal*, vol. 48, no. 2, pp. 373–384, 1998.
- [5] C. Nowak and H. Stettner, "Nonuniqueness results for ordinary differential equations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 30, no. 6, pp. 3935–3938, 1997.
- [6] K. Balla, "Solution of singular boundary value problems for non-linear systems of ordinary differential equations," *U.S.S.R. Computational Mathematics and Mathematical Physics*, vol. 20, no. 4, pp. 100–115, 1980.
- [7] V. A. Čečík, "Investigation of systems of ordinary differential equations with a singularity," *Trudy Moskovskogo Matematičeskogo Obščestva*, vol. 8, pp. 155–198, 1959 (Russian).
- [8] J. Diblík and C. Nowak, "A nonuniqueness criterion for a singular system of two ordinary differential equations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 64, no. 4, pp. 637–656, 2006.
- [9] J. Diblík and M. Růžičková, "Inequalities for solutions of singular initial problems for Carathéodory systems via Ważewski's principle," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 69, no. 12, pp. 4482–4495, 2008.
- [10] J. Diblík and M. Růžičková, "Existence of positive solutions of a singular initial problem for a nonlinear system of differential equations," *The Rocky Mountain Journal of Mathematics*, vol. 34, no. 3, pp. 923–944, 2004.
- [11] J. Kalas, "Nonuniqueness theorem for a singular Cauchy problem," *Georgian Mathematical Journal*, vol. 7, no. 2, pp. 317–327, 2000.
- [12] I. T. Kiguradze, *Some Singular Boundary Value Problems for Ordinary Differential Equations*, Tbilisi University Press, Tbilisi, Georgia, 1975.
- [13] C. Nowak, "Some remarks on a paper by M. Samimi on: "Nonuniqueness criteria for ordinary differential equations"," *Applicable Analysis*, vol. 47, no. 1, pp. 39–44, 1992.
- [14] V. Lakshmikantham and S. Leela, *Differential and Integral Inequalities: Theory and Applications. Vol. I: Ordinary Differential Equations*, Academic Press, New York, NY, USA, 1969.
- [15] T. Rouche, P. Habets, and M. Laloy, *Stability Theory by Liapunov's Direct Method*, Springer, New York, NY, USA, 1977.
- [16] T. Yoshizawa, *Stability Theory by Liapunov's Second Method*, The Mathematical Society of Japan, Tokyo, Japan, 1966.
- [17] P. Hartman, *Ordinary Differential Equations*, vol. 38 of *Classics in Applied Mathematics*, SIAM, Philadelphia, Pa, USA, 2002.
- [18] M. Nekvinda, "On uniqueness of solutions of differential equations," *Časopis Pro Pěstování Matematiky*, vol. 108, no. 1, pp. 1–7, 1983.