

On pathwise quadratic variation for càdlàg functions

Henry Chiu* Rama Cont†

Abstract

We revisit Föllmer’s concept of quadratic variation of a càdlàg function along a sequence of time partitions and discuss its relation with the Skorokhod topology. We show that in order to obtain a robust notion of pathwise quadratic variation applicable to sample paths of càdlàg processes, one must reformulate the definition of pathwise quadratic variation as a limit in Skorokhod topology of discrete approximations along the partition. One then obtains a simpler definition which implies the Lebesgue decomposition of the pathwise quadratic variation as a result, rather than requiring it as an extra condition.

Keywords: quadratic variation; semimartingale; pathwise calculus; Ito formula; pathwise integration; cadlag functions; Skorokhod topology.

AMS MSC 2010: 60H05; 26B35.

Submitted to ECP on June 19, 2018, final version accepted on October 30, 2018.

Supersedes arXiv:1806.07290.

Contents

1 Quadratic variation along a sequence of partitions	1
2 Pathwise quadratic variation for cadlag functions	3
3 Quadratic variation for multidimensional functions	7
4 Some applications	9
References	11

1 Quadratic variation along a sequence of partitions

In his seminal paper *Calcul d’Itô sans probabilités* [14], Hans Föllmer introduced a pathwise concept of quadratic variation and used it to provide a pathwise proof of the Itô formula. Föllmer showed that if a function $x \in D([0, T], \mathbb{R})$ has quadratic variation along a sequence $\pi^n = (t_0^n = 0 < \dots < t_j^n < \dots < t_{m(n)}^n = T)$ of time partitions of $[0, T]$ in the sense that for each $t \in [0, T]$ the limit

$$\mu^n := \sum_{[t_j^n, t_{j+1}^n] \in \pi_n} \delta(\cdot - t_j) |x(t_{j+1}^n) - x(t_j^n)|^2 \quad (1.1)$$

*Dept of Mathematics, Imperial College London. E-mail: h.chiu16@imperial.ac.uk

†Laboratoire de Probabilités, Statistiques et Modélisation, CNRS-Sorbonne Université
E-mail: Rama.Cont@math.cnrs.fr

converges weakly to a Radon measure μ_π such that

$$[x]_\pi^c : t \mapsto \mu_\pi([0, t]) - \sum_{s \leq t} |\Delta x(s)|^2 \text{ is a continuous increasing function,} \quad (\text{L})$$

then a pathwise Itô formula may be obtained for functions of x [14]: for any $f \in C^2(\mathbb{R}, \mathbb{R})$,

$$f(x(t)) - f(x(0)) = \int_0^t f'(x) d^\pi x + \frac{1}{2} \int_0^t f''(x(s)) d[x]_\pi^c(s) + \sum_{0 \leq s \leq t} (f(x(s)) - f(x(s-)) - f'(x(s-))\Delta x(s)) \quad (1.2)$$

where $\int_0^t f'(x) d^\pi x$ is a pathwise integral defined as a limit of left Riemann sums computed along the partition:

$$\int_0^t g(x) d^\pi x := \lim_{n \rightarrow \infty} \sum_{\pi_n} g(x(t_j^n)) \cdot (x(t_{j+1}^n) - x(t_j^n)). \quad (1.3)$$

The quantity

$$[x]_\pi(t) = \mu_\pi([0, t]) = [x]_\pi^c(t) + \sum_{s \leq t} |\Delta x(s)|^2$$

is called the quadratic variation of x along π . This result has many interesting applications and has been extended to less regular functions [2, 8, 11, 19] and path-dependent functionals [1, 4, 5, 6, 8]. With the exception of [6, 22, 18], these extensions have focused on continuous paths.

Föllmer's definition [14] contains the condition (L) on the Lebesgue decomposition of the limit μ_π : the atoms of μ_π should correspond exactly to the jumps of x and their mass should be $|\Delta x(t)|^2$ or, equivalently, the discontinuity points of $[x]_\pi$ should coincide with those of x , with $\Delta[x]_\pi(t) = |\Delta x(t)|^2$. This condition can not be removed: as shown by Coquet et al. [9], there are counterexamples of continuous functions x such that (1.1) converges to a limit with atoms. Conversely, one can give examples of discontinuous functions for which (1.1) converges to an atomless measure. If this condition is not satisfied, then the pathwise integral (1.3) fails to satisfy

$$\Delta \left(\int_0^t x(s-) dx(s) \right) (t) = x(t-) \Delta x(t),$$

at each t where condition (L) is not met. On the other hand, this condition (L) is not easy to check and seems to require a link between the path x and the sequence of partitions π , making it difficult to apply to sample paths of stochastic processes.

In this work, we revisit Föllmer's concept of pathwise quadratic variation along a sequence of partitions and show that it has hitherto unsuspected links with the Skorokhod topology. In particular, we show that in order to obtain a robust notion of pathwise quadratic variation applicable to sample paths of càdlàg processes, one must reformulate the definition of the quadratic variation as a limit, in Skorokhod topology, of discrete approximations defined along the partition. This leads to a simpler definition of pathwise quadratic variation which holds in any dimension and, rather than requiring the Lebesgue decomposition of the pathwise quadratic variation as an extra condition, yields it as a consequence.

Outline We begin by recalling Föllmer's definition of pathwise quadratic variation and variations of it which have been used in the literature. We then introduce a new definition of quadratic variation for real-valued càdlàg functions based on the Skorokhod topology and prove equivalence among the various definitions. Section 3 extends the results to vector-valued functions: we show that, unlike Föllmer's original definition in which the one dimensional case plays a special role, our definition applies regardless of dimension, thus simplifying various statements regarding quadratic variation for vector-valued functions. Finally, in Section 4, we show that our approach leads to simple proofs for various properties of pathwise quadratic variation.

2 Pathwise quadratic variation for cadlag functions

Let $\pi := (\pi_n)_{n \geq 1}$ be a sequence of partitions $\pi_n = (t_0^n, \dots, t_{k_n}^n)$ of $[0, \infty)$ into intervals $0 = t_0^n < \dots < t_{k_n}^n < \infty$; $t_{k_n}^n \uparrow \infty$ with vanishing mesh $|\pi_n| \downarrow 0$ on compacts. By convention, $\max(\emptyset \cap \pi_n) := 0$, $\min(\emptyset \cap \pi_n) := t_{k_n}^n$.

Denote $\mathcal{D} := \mathcal{D}([0, \infty), \mathbb{R})$ the space of cadlag functions and $\mathcal{C} := \mathcal{C}([0, \infty), \mathbb{R})$ the subspace of real-valued continuous functions. We equip \mathcal{D} with a metric d which induces the Skorokhod J_1 topology [16, Ch. VI]. Denote $\mathcal{D}_0^+ \subset \mathcal{D}$ to be the subset of non-negative increasing right-continuous functions null at 0.

Definition 2.1 (Föllmer 1981). $x \in \mathcal{D}$ has (finite) quadratic variation $[x]_\pi$ along π if the sequence of measures

$$\mu_n := \sum_{t_i^n \in \pi_n} (x(t_{i+1}^n) - x(t_i^n))^2 \delta(t_i^n) \tag{2.1}$$

converges vaguely on $[0, \infty)$ to a Radon measure μ with $[x]_\pi(t) = \mu([0, t])$, such that $[x]_\pi^c$ defined by $[x]_\pi^c(t) := \mu([0, t]) - \sum_{s \leq t} (\Delta x_s)^2$ is a continuous increasing function. $[x]_\pi(t) = \mu([0, t])$ is then called the quadratic variation of x along π , and admits the following Lebesgue decomposition:

$$[x]_\pi(t) = [x]_\pi^c(t) + \sum_{s \leq t} |\Delta x(s)|^2. \tag{2.2}$$

We denote \mathcal{Q}_0^+ the set of $x \in \mathcal{D}$ satisfying these properties.

At this point let us point out a link between vague and weak convergence of Radon measures on $[0, \infty)$, a link which is well known in the case where the measures are sub-probability measures:

Lemma 2.2. Let v_n and v be non-negative Radon measures on $[0, \infty)$ and $J \subset [0, \infty)$ be the set of atoms of v , the followings are equivalent:

- (i) $v_n \rightarrow v$ vaguely on $[0, \infty)$.
- (ii) $v_n \rightarrow v$ weakly on $[0, T]$ for every $T \notin J$.

Proof. Let $f \in C_K([0, \infty))$ be a compactly supported continuous function. Since J is countable, $\exists T \notin J$; $\text{supp}(f) \subset [0, T]$. Now (ii) $\Rightarrow \int_0^\infty f dv_n = \int_0^T f dv_n \rightarrow \int_0^T f dv = \int_0^\infty f dv \Rightarrow$ (i). Suppose (i) holds, let $T \notin J$ and $f \in \mathcal{C}([0, T], \|\cdot\|_\infty)$. Since $f = (f)^+ - (f)^-$, we may take $f \geq 0$ and define the following extensions:

$$\begin{aligned} \bar{f}^\epsilon(t) &:= f(t) \mathbb{I}_{[0, T]}(t) + f(T) \left(1 + \frac{T-t}{\epsilon}\right) \mathbb{I}_{(T, T+\epsilon]}(t) \\ \underline{f}^\epsilon(t) &:= f(t) \mathbb{I}_{[0, T-\epsilon]}(t) + f(T) \left(\frac{T-t}{\epsilon}\right) \mathbb{I}_{(T-\epsilon, T]}(t), \end{aligned}$$

then $\bar{f}^\epsilon, \underline{f}^\epsilon \in \mathcal{C}_K([0, \infty))$, $0 \leq \underline{f}^\epsilon \leq f \mathbb{I}_{[0, T]} \leq \bar{f}^\epsilon \leq \|f\|_\infty$ and we have

$$\int_0^\infty \underline{f}^\epsilon dv_n \leq \int_0^T f dv_n \leq \int_0^\infty \bar{f}^\epsilon dv_n.$$

Since $v_n \rightarrow v$ vaguely and $T \notin J$, thus, as $n \rightarrow \infty$ we obtain

$$\begin{aligned} 0 &\leq \limsup_n \int_0^T f dv_n - \liminf_n \int_0^T f dv_n \leq \int_0^\infty \bar{f}^\epsilon - \underline{f}^\epsilon dv \\ &\leq \|f\|_\infty v((T-\epsilon, T+\epsilon]) \xrightarrow{\epsilon \rightarrow 0} 0, \end{aligned}$$

hence by monotone convergence

$$\lim_n \int_0^T f dv_n = \lim_\epsilon \int_0^\infty \underline{f}^\epsilon dv = \int_0^T f dv$$

and (ii) follows. □

Proposition 2.3. *If $x \in Q_0^\pi$, then the pointwise limit s of*

$$s_n(t) := \sum_{t_i \in \pi_n} |x(t_{i+1} \wedge t) - x(t_i \wedge t)|^2 \tag{2.3}$$

exists, $s = [x]_\pi$ and s admits the Lebesgue decomposition:

$$s(t) = s^c(t) + \sum_{s \leq t} (\Delta x_s)^2. \tag{2.4}$$

Proof. If $x \in Q_0^\pi$, define

$$q_n(t) := \sum_{\pi_n \ni t_i \leq t} (x(t_{i+1}) - x(t_i))^2,$$

the distribution function of μ_n in (2.1). Since $\mu_n \rightarrow \mu$ vaguely, we have $q_n \rightarrow [x]$ pointwise at all continuity points of $[x]$ (Lemma 2.2 & [12, X.11]). Let I be the set of continuity points of $[x]$. Observe (q_n) is monotonic in $[0, \infty)$ and I is dense in $[0, \infty)$, if $t \notin I$, it follows [12, X.8] that

$$[x](t-) \leq \liminf_n q_n(t) \leq \limsup_n q_n(t) \leq [x](t+) = [x](t).$$

Thus, we may take any subsequence (n_k) such that $\lim_k q_{n_k}(t) =: q(t)$. Since $x \in Q_0^\pi$ and the Lebesgue decomposition (2.2) holds on $[x]$, we have

$$([x](t + \epsilon) - q(t)) + (q(t) - [x](t - \epsilon)) = [x](t + \epsilon) - [x](t - \epsilon) \xrightarrow{\epsilon \rightarrow 0} |\Delta x(t)|^2. \tag{2.5}$$

If $t \pm \epsilon \in I$, $\tilde{\pi}_k := \pi_{n_k}$ and $t_j^{(k)} := \max\{\tilde{\pi}_k \cap [0, t]\}$, the second sum in (2.5) is

$$\lim_k \sum_{\substack{t_i \in \tilde{\pi}_k; \\ t - \epsilon < t_i \leq t}} (x(t_{i+1}) - x(t_i))^2 = \lim_k \sum_{\substack{t_i \in \tilde{\pi}_k; \\ t - \epsilon < t_i < t_j^{(k)}}} (x(t_{i+1}) - x(t_i))^2 + (\Delta x(t))^2 \geq (\Delta x(t))^2$$

by the fact that x is càdlàg and that $t \notin I$.

We see from (2.5) that $q(t) = [x](t)$ as $\epsilon \rightarrow 0$. Since the choice of the convergent subsequence is arbitrary, we conclude that $q_n \rightarrow [x]$ pointwise on $[0, \infty)$. Observe that the pointwise limits of (s_n) and (q_n) coincide i.e.

$$|s_n(t) - q_n(t)| = (x_{t_{i+1}^{(n)}} - x(t))^2 + 2(x_{t_{i+1}^{(n)}} - x(t))(x(t) - x_{t_i^{(n)}}) \tag{2.6}$$

converges to 0 by the right-continuity of x , where $t_i^{(n)} := \max\{\pi_n \cap [0, t]\}$ and that $x \in Q_0^\pi$, Prop. 2.3 follows. □

Denote Q_1^π the set of $x \in \mathcal{D}$ such that (s_n) defined in (2.3) has a pointwise limit s with Lebesgue decomposition given by (2.4). Then $Q_0^\pi \subset Q_1^\pi$ and we have:

Proposition 2.4. *If $x \in \mathcal{Q}_1^\pi$, then the pointwise limit q of*

$$q_n(t) := \sum_{\pi_n \ni t_i \leq t} (x(t_{i+1}) - x(t_i))^2 \tag{2.7}$$

exists. Furthermore $q = s$ and admits the Lebesgue decomposition:

$$q(t) = q^c(t) + \sum_{s \leq t} (\Delta x_s)^2. \tag{2.8}$$

Proof. Since the pointwise limits of (s_n) and (q_n) coincide by (2.6). Prop. 2.4 now follows from $x \in \mathcal{Q}_1^\pi$. \square

Denote \mathcal{Q}_2^π the set of $x \in \mathcal{D}$ such that the quadratic sums (q_n) defined in (2.7) have a pointwise limit q with Lebesgue decomposition (2.8). Then $\mathcal{Q}_0^\pi \subset \mathcal{Q}_1^\pi \subset \mathcal{Q}_2^\pi$ and we have:

Proposition 2.5. *If $x \in \mathcal{Q}_2^\pi$, then $q_n \rightarrow q$ in the Skorokhod topology.*

Proof. Since $x \in \mathcal{Q}_2^\pi$, we have $q_n \rightarrow q$ pointwise on $[0, \infty)$ and that $(q_n), q$ are elements in \mathcal{D}_0^+ . By [16, Thm.VI.2.15], it remains to show that

$$\sum_{s \leq t} (\Delta q_n(s))^2 \xrightarrow{n \rightarrow \infty} \sum_{s \leq t} (\Delta q(s))^2$$

on a dense subset of $[0, \infty)$. Let $t > 0$, define $J^\epsilon := \{s \geq 0 \mid (\Delta X_s)^2 \geq \frac{\epsilon}{2}\}$, $J_n^\epsilon := \{t_i \in \pi_n \mid \exists s \in (t_i, t_{i+1}]; (\Delta X_s)^2 \geq \frac{\epsilon}{2}\} \subset \pi_n$ and observe that

$$\begin{aligned} \sum_{s \leq t} (\Delta q_n(s))^2 &= \sum_{\pi_n \ni t_i \leq t} (x(t_{i+1}) - x(t_i))^4 \\ &= \sum_{J_n^\epsilon \ni t_i \leq t} (x(t_{i+1}) - x(t_i))^4 + \sum_{(J_n^\epsilon)^c \ni t_i \leq t} (x(t_{i+1}) - x(t_i))^4. \end{aligned} \tag{2.9}$$

Since x is càdlàg and that $|\pi_n| \downarrow 0$ on compacts, the first sum in (2.9) converges to $\sum_{J^\epsilon \ni s \leq t} (\Delta x_s)^4$ and the second sum in (2.9)

$$\begin{aligned} \sum_{(J_n^\epsilon)^c \ni t_i \leq t} (x(t_{i+1}) - x(t_i))^4 &\leq \left(\sup_{(J_n^\epsilon)^c \ni t_i \leq t} (x(t_{i+1}) - x(t_i))^2 \right) \sum_{(J_n^\epsilon)^c \ni t_i \leq t} (x(t_{i+1}) - x(t_i))^2 \\ &\leq \epsilon q(t) \end{aligned}$$

for sufficiently large n [6, Appendix A.8] hence

$$\lim_n \sum_{s \leq t} (\Delta q_n(s))^2 = \sum_{J^\epsilon \ni s \leq t} (\Delta x_s)^4 + \overbrace{\limsup_n \sum_{(J_n^\epsilon)^c \ni t_i \leq t} (x(t_{i+1}) - x(t_i))^4}^{\leq \epsilon q(t)}.$$

By the Lebesgue decomposition (2.8), we observe $\sum_{J^\epsilon \ni s \leq t} (\Delta x_s)^4 \leq q(t)^2$ and that

$$\lim_n \sum_{s \leq t} (\Delta q_n(s))^2 = \sum_{s \leq t} (\Delta x_s)^4 = \sum_{s \leq t} (\Delta q(s))^2$$

as $\epsilon \rightarrow 0$. \square

Denote \mathcal{Q}^π the set of càdlàg functions $x \in \mathcal{D}$ such that the limit \tilde{q} of (q_n) exists in (\mathcal{D}, d) . Then $\mathcal{Q}_0^\pi \subset \mathcal{Q}_1^\pi \subset \mathcal{Q}_2^\pi \subset \mathcal{Q}^\pi$ and we have:

Proposition 2.6. *$\mathcal{Q}^\pi \subset \mathcal{Q}_0^\pi$ and $\tilde{q} = [x]_\pi$.*

Proof. Let $x \in \mathcal{Q}^\pi$ and I be the set of continuity points of \tilde{q} . [16, VI.2.1(b.5)] implies that $q_n \rightarrow \tilde{q}$ pointwise on I . Since $q_n \in \mathcal{D}_0^+$ and I is dense on $[0, \infty)$, it follows $\tilde{q} \in \mathcal{D}_0^+$. Denote μ to be the Radon measure of \tilde{q} on $[0, \infty)$, observe the set of atoms of μ is $J := [0, \infty) \setminus I$ and that (q_n) are the distribution functions of the discrete measures (μ_n) in (2.1). Thus, by (Lemma 2.2 & [12, X.11]), we see that $\mu_n \rightarrow \mu$ vaguely on $[0, \infty)$.

If $t > 0$, put $t_i^{(n)} := \max\{\pi_n \cap [0, t)\}$. Since $|\pi_n| \downarrow 0$ on compacts, we have $t_i^{(n)} < t$, $t_i^{(n)} \uparrow t$ and $t_{i+1}^{(n)} \downarrow t$. Observe that

$$\Delta q_n(t) = \begin{cases} (x(t_{i+1}) - x(t_i))^2, & \text{if } t = t_i \in \pi_n. \\ 0, & \text{otherwise.} \end{cases} \tag{2.10}$$

If $\Delta \tilde{q}(t) = 0$, [16, VI.2.1(b.5)] implies that $\Delta q_n(t_i^{(n)}) \rightarrow \Delta \tilde{q}(t)$. Hence, by the fact that x is càdlàg, $(\Delta x(t))^2 = \lim_n \Delta q_n(t_i^{(n)}) = \Delta \tilde{q}(t)$. If $\Delta \tilde{q}(t) > 0$, there exists [16, VI.2.1(a)] a sequence $t'_n \rightarrow t$ such that $\Delta q_n(t'_n) \rightarrow \Delta \tilde{q}(t) > 0$. Using the fact that x is càdlàg, $t'_n \rightarrow t$, (2.10) and [16, VI.2.1(b)], we deduce that (t'_n) must coincide with $(t_i^{(n)})$ for all n sufficiently large, else we will contradict $\Delta \tilde{q}(t) > 0$. Thus, $(\Delta x(t))^2 = \lim_n \Delta q_n(t_i^{(n)}) = \lim_n \Delta q_n(t'_n) = \Delta \tilde{q}(t)$ and the Lebesgue decomposition (2.2) holds on \tilde{q} .

By Def.2.1, we have $\tilde{q} = [x]_\pi$ hence $\mathcal{Q}^\pi \subset \mathcal{Q}_0^\pi$. □

Theorem 2.7. *Let*

- \mathcal{Q}_0^π be the set of $x \in \mathcal{D}$ satisfying Definition 2.1.
- \mathcal{Q}_1^π the set of $x \in \mathcal{D}$ such that (s_n) defined in (2.3) has a pointwise limit s with Lebesgue decomposition given by (2.4).
- \mathcal{Q}_2^π the set of $x \in \mathcal{D}$ such that the quadratic sums (q_n) defined by (2.7) have a pointwise limit q with Lebesgue decomposition (2.8).
- \mathcal{Q}^π the set of càdlàg functions $x \in \mathcal{D}$ such that the limit \tilde{q} of (q_n) exists in (\mathcal{D}, d) .

Then:

- (i) $\mathcal{Q}_0^\pi = \mathcal{Q}_1^\pi = \mathcal{Q}_2^\pi = \mathcal{Q}^\pi$.
- (ii) If $x \in \mathcal{Q}_0^\pi$, then $[x]_\pi = s(x) = q(x) = \tilde{q}(x)$.
- (iii) x has finite quadratic variation along π if and only if the sequence $(q_n)_{n \geq 1}$ defined by

$$q_n(t) := \sum_{\pi_n \ni t_i \leq t} (x(t_{i+1}) - x(t_i))^2$$

converges in (\mathcal{D}, d) .

- (iv) If $(q_n)_{n \geq 1}$ converges in (\mathcal{D}, d) , the limit is equal to $[x]_\pi$.

Proof. These results are a consequence of Prop. 2.3, 2.4, 2.5 and 2.6. □

We see that the two defining properties of $[x]$ in Def. 2.1 are consequences per Thm. 2.7. The following corollary treats the special case of continuous functions.

Corollary 2.8. *Let $x \in \mathcal{Q}^\pi$, s_n defined as in (2.3), (q_n) defined by (2.7).*

- i $q_n \rightarrow [x]$ uniformly on compacts in $[0, \infty)$ if and only if x is continuous.
- ii If $q_n \rightarrow [x]_\pi$ uniformly on compacts in $[0, \infty)$, then $s_n \rightarrow [x]_\pi$ uniformly on compacts.

Proof. (i): The if part follows from Prop. 2.6, (2.2) and [16, VI.1.17(b)]. The only if part: Let $t \geq 0$, it is well known that

$$\Delta q_n(t) \rightarrow \Delta[x]_\pi(t)$$

by uniform convergence. Put $t'_n := \max\{t_i < t | t_i \in \pi_n\}$, since $q_n \rightarrow [x]$ in the Skorokhod topology, we also have

$$\Delta q_n(t'_n) \rightarrow \Delta[x]_\pi(t)$$

by (2.10) and [16, VI.2.1(b)]. If $\Delta[x](t) > 0$, then [16, VI.2.1(b)] implies t'_n must coincide with t for all n large enough, but $t'_n < t$ for all n , hence $\Delta[x](t) = 0$ which implies $\Delta x(t) = 0$ by Prop. 2.6 and (2.2). Since t is arbitrary, we conclude that $x \in \mathcal{C}$.

(ii): Let $T > 0$, $\|\cdot\|(t)$ the supremum norm on $\mathcal{D}([0, T])$ and observe that

$$\|s_n - [x]\|(t) \leq \|q_n - [x]\|(t) + \|s_n - q_n\|(t).$$

Since (i) implies $x \in \mathcal{C}$, (ii) now follows from uniform continuity of x and (2.8). □

Remark 2.9. The converse of (ii) is not true in general.

Remark 2.10. The above conditions cannot be replaced by the pointwise convergence of

$$p_n(t) := \sum_{\pi_n \ni t_{i+1} \leq t} (x(t_{i+1}) - x(t_i))^2$$

together with the Lebesgue decomposition (2.2). To see why, take $t_0 \notin \pi$, put $x(t) := \mathbb{1}_{[t_0, \infty)}(t)$ then obviously $[x](t_0) = \lim s_n(t_0) = \lim q_n(t_0) = 1$ but $\lim p_n(t_0) = 0$.

Remark 2.11. Vovk [22, Sec. 6] proposes a different notion of pathwise quadratic variation along a sequence of partitions, which is shown to coincide with Föllmer's definition under the additional assumption that the sequence of partitions π is refining and exhausts all discontinuity points of the path [22, Prop. 6.3 & 6.4].

This requirement of exhausting all jumps can always be satisfied for a given càdlàg path by adding all discontinuity points to the sequence of partitions. However if one is interested in applying this definition to a process, say a semimartingale, then in general there may exist no sequence of partitions satisfying this condition almost surely across sample paths. And, if this requirement of exhausting all discontinuity points is removed, then Vovk's definition differs from Föllmer's (and therefore, fails to satisfy the Ito formula (1.2) in general).

By contrast, our definition does not require such a condition and easily carries over to stochastic processes without requiring the use of random partitions (see Theorem 4.1).

3 Quadratic variation for multidimensional functions

Denote $\mathcal{D}^m := \mathcal{D}([0, \infty), \mathbb{R}^m)$ and $\mathcal{D}^{m \times m} := \mathcal{D}([0, \infty), \mathbb{R}^{m \times m})$ to be the Skorokhod spaces [3, 21, 17], each of which equipped with a metric d which induces the corresponding Skorokhod J_1 topology [16, Ch. VI]. $\mathcal{C}^m := \mathcal{C}([0, \infty), \mathbb{R}^m)$ the subspace of continuous functions in \mathcal{D}^m . We recall Theorem 2.7 from the one dimensional case $n = 1$ that (Def. 2.1) and (Thm. 2.7.iii) are equivalent.

It is known that $x, y \in \mathcal{Q}^\pi$ does not imply $x + y \in \mathcal{Q}^\pi$ [7, 20] so one cannot for instance define a quadratic covariation $[x, y]_\pi$ of two such functions in the obvious way. This prevents a simple componentwise definition of the finite quadratic variation property for

vector-valued functions. Therefore, the notion of quadratic variation in the multidimensional setting was originally defined in [14] as follows:

Definition 3.1 (Föllmer 1981). *We say that $\mathbf{x} := (x^1, \dots, x^m)^T \in \mathcal{D}^m$ has finite quadratic variation along π if all $x^i, x^i + x^j$ ($1 \leq i, j \leq m$) have finite quadratic variation.*

The quadratic (co)variation $[x^i, x^j]$ is then defined as

$$[x^i, x^j]_\pi(t) := \frac{1}{2} \left([x^i + x^j]_\pi(t) - [x^i]_\pi(t) - [x^j]_\pi(t) \right), \tag{3.1}$$

which admits the following Lebesgue decomposition:

$$[x^i, x^j]_\pi(t) = [x^i, x^j]_\pi^c(t) + \sum_{s \leq t} \Delta x^i(s) \Delta x^j(s). \tag{3.2}$$

The function $[\mathbf{x}]_\pi := ([x^i, x^j]_{1 \leq i, j \leq m})$, which takes values in the cone of symmetric semidefinite positive matrices, is called the quadratic (co)variation of \mathbf{x} .

Note Def. 3.1 requires first introducing the case $m = 1$. The following definition, by contrast, avoids this and directly defines the concept of multidimensional quadratic variation in any dimension:

Definition 3.2. $\mathbf{x} \in \mathcal{D}^m$ has finite quadratic variation $[\mathbf{x}]_\pi$ along π if

$$\mathbf{q}_n(t) := \sum_{\pi_n \ni t_i \leq t} (\mathbf{x}(t_{i+1}) - \mathbf{x}(t_i))(\mathbf{x}(t_{i+1}) - \mathbf{x}(t_i))^T$$

converges to $[\mathbf{x}]_\pi$ in $(\mathcal{D}^{m \times m}, d)$.

We shall now prove the equivalence of these definitions.

Define, for $u, v, w \in \mathcal{D}$

$$q_n^{(u,v)}(t) := \sum_{\pi_n \ni t_i \leq t} (u_{t_{i+1}} - u_{t_i})(v_{t_{i+1}} - v_{t_i})$$

and $q_n^{(w)} := q_n^{(w,w)}$. Note that the Skorokhod topology on (\mathcal{D}^m, d) is strictly finer than the product topology on $(\mathcal{D}, d)^m$ [16, VI.1.21] and that (\mathcal{D}, d) is not a topological vector space [16, VI.1.22]. The following lemma is essential:

Lemma 3.3. *Let $t > 0$. There exists a sequence $t_n \rightarrow t$ such that for all $u, v \in \mathcal{D}$, $(q_n^{(u,v)})$ converges in (\mathcal{D}, d) then*

$$\lim_n \left(\Delta q_n^{(u,v)}(t_n) \right) = \Delta \left(\lim_n q_n^{(u,v)} \right) (t).$$

Note that the sequence t_n is chosen from the partition points of π_n , independently of $u, v \in \mathcal{D}$.

Proof. Define $t_i^{(n)} := \max\{\pi_n \cap [0, t]\}$. Since $|\pi_n| \downarrow 0$ on compacts, we have $t_i^{(n)} < t$, $t_i^{(n)} \uparrow t$ and $t_{i+1}^{(n)} \downarrow t$. Observe that

$$\Delta q_n^{(u,v)}(t) = \begin{cases} (u_{t_{i+1}} - u_{t_i})(v_{t_{i+1}} - v_{t_i}), & \text{if } t = t_i \in \pi_n. \\ 0, & \text{otherwise.} \end{cases} \tag{3.3}$$

Put $\tilde{q} := \lim_n q_n^{(u,v)}$. If $\Delta \tilde{q}(t) = 0$, [16, VI.2.1(b.5)] implies that $\Delta q_n^{(u,v)}(t_i^{(n)}) \rightarrow \Delta \tilde{q}(t)$. If $\Delta \tilde{q}(t) > 0$, there exists [16, VI.2.1(a)] a sequence $t'_n \rightarrow t$ such that $\Delta q_n^{(u,v)}(t'_n) \rightarrow \Delta \tilde{q}(t) > 0$. Using the fact that u, v are càdlàg, $t'_n \rightarrow t$ and (3.3), we deduce that (t'_n) must coincide with $(t_i^{(n)})$ for n sufficiently large, else we will contradict $\Delta \tilde{q}(t) > 0$. Put $t_n := t_i^{(n)}$. \square

Proposition 3.4. *Let $x, y \in \mathcal{Q}^\pi$, then $(q_n^{(x+y)})$ converges in (\mathcal{D}, d) if and only if $(q_n^{(x,y)})$ does. In this case, $x + y \in \mathcal{Q}^\pi$ and $\lim_n q_n^{(x,y)} = \frac{1}{2}([x + y] - [x] - [y])$.*

Proof. Since

$$q_n^{(x+y)} = q_n^{(x)} + q_n^{(y)} + 2q_n^{(x,y)}$$

and that $x, y \in \mathcal{Q}^\pi$, Prop. 3.4 follows from Lemma 3.3 and [16, VI.2.2(a)]. □

Proposition 3.5. *(q_n) converges in $(\mathcal{D}^{m \times m}, d)$ if and only if it converges in $(\mathcal{D}, d)^{m \times m}$.*

Proof. Since the Skorokhod topology on $(\mathcal{D}^{m \times m}, d)$ is strictly finer than the product topology on $(\mathcal{D}, d)^{m \times m}$ [16, VI.1.21], we have $(\mathcal{D}^{m \times m}, d)$ convergence implies $(\mathcal{D}, d)^{m \times m}$ convergence. The other direction follows from the observation that

$$\mathbf{q}_n = \left(q_n^{(x^i, x^j)} \right)_{1 \leq i \leq j \leq n}, \tag{3.4}$$

satisfies Lemma 3.3 and [16, VI.2.2(b)]. □

Theorem 3.6. *Definitions 3.1 and 3.2 are equivalent.*

Proof. This is a consequence of (3.1), (3.4), Prop. 3.4 & 3.5 and Thm. 2.7 □

Corollary 3.7. *If $\mathbf{x} \in \mathcal{D}^m$ has finite quadratic variation, then*

- (i) $\mathbf{q}_n \rightarrow [\mathbf{x}]$ locally uniformly on $[0, \infty)$ if and only if $\mathbf{x} \in \mathcal{C}^n$.
- (ii) $F(\mathbf{q}_n) \rightarrow F([\mathbf{x}])$ for all functionals $F : \mathcal{D}^m \rightarrow \mathbb{R}$ which are J_1 -continuous at $[\mathbf{x}]$.

Proof. This is a consequence of Thm.3.6, (3.2) and [16, VI.1.17.b]. □

Remark 3.8. For \mathbf{x} to have finite quadratic variation, it is sufficient that (\mathbf{q}_n) converges in $(\mathcal{D}, d)^{m \times m}$ due to Prop. 3.5. (i.e. component-wise convergence)

4 Some applications

We now show that our approach yields simple proofs for some properties of pathwise quadratic variation, which turn out to be useful in the study of pathwise approaches to Ito calculus.

Denote $\mathcal{D} := \mathcal{D}([0, \infty), \mathbb{R})$ and $\mathcal{D}^{d \times d} := \mathcal{D}([0, \infty), \mathbb{R}^{d \times d})$ to be the Skorokhod spaces, each of which equipped with a *complete* metric δ which induces the corresponding Skorokhod (a.k.a J_1) topology. Denote \mathcal{F} to be the J_1 Borel sigma algebra of \mathcal{D} (a.k.a the canonical sigma algebra generated by coordinates). Recall that \mathcal{Q}_0^π is the set of paths with finite quadratic variation along π in the sense of Def. 2.1.

Our first application is to give a criterion for a function $x \in \mathcal{D}$ to have finite quadratic variation. In Def. 2.1, one needs to verify the condition (2.2) on the Lebesgue decomposition of the limit μ . The following characterization, by contrast, does not refer to μ :

Property 1. $x \in \mathcal{Q}_0^\pi$ if and only if (q_n) defined by (2.7) is a Cauchy sequence in (\mathcal{D}, δ) .

Proof. This is a consequence of Thm. 2.7 and the completeness of (\mathcal{D}, δ) . □

One of the main advantages of having convergence in the J_1 topology is that it ensures convergence of jumps in a regulated manner. It comes in handy when accessing the limit of $q_n(t_n)$ as $n \rightarrow \infty$.

Property 2. Let $x \in Q_0^\pi$, for each $t \geq 0$, we define $t'_n := \max\{t_i < t | t_i \in \pi_n\}$, then

$$\begin{aligned} t_n \longrightarrow t; t_n \leq t'_n &\implies q_n(t_n-) \longrightarrow [x]_\pi(t-), \\ t_n \longrightarrow t; t_n < t'_n &\implies q_n(t_n) \longrightarrow [x]_\pi(t-), \\ t_n \longrightarrow t; t_n \geq t'_n &\implies q_n(t_n) \longrightarrow [x]_\pi(t), \\ t_n \longrightarrow t; t_n > t'_n &\implies q_n(t_n-) \longrightarrow [x]_\pi(t). \end{aligned}$$

In particular, the sequence (t'_n) is asymptotically unique in the sense that any other sequence (t''_n) meeting the above properties coincides with (t'_n) for n sufficiently large.

Proof. This is a consequence of Thm. 2.7 and [16, VI.2.1]. □

Given a càdlàg process X (i.e. a $(\mathcal{D}, \mathcal{F})$ -measurable random variable), a natural quantity to consider is $\mathbb{P}(X \in Q_0^\pi)$. This only makes sense however if Q_0^π is \mathcal{F} -measurable. This 'natural' property, not easy to show using the original definition (Def. 2.1), becomes simple thanks to Theorem. 2.7:

Property 3 (Measurability of Q_0^π). Q_0^π is \mathcal{F} -measurable.

Proof. By Thm. 2.7, $Q_0^\pi = Q^\pi$ and by definition, Q^π is the J_1 convergence set of

$$x \longmapsto \sum_{\pi_n \ni t_i \leq \cdot} (x(t_{i+1}) - x(t_i))^2,$$

$n \geq 1$ on \mathcal{D} . Since \mathcal{D} is completely metrisable, the claim follows from [12, V.3]. □

Föllmer introduced in [14, 15] the class of processes with finite quadratic variation ('processus à variation quadratique'), defined as càdlàg processes such that the sequence

$$S_n(t) := \sum_{\pi_n \ni t_i \leq t} (X(t_{i+1}) - X(t_i))^2$$

converges in probability for every t to an increasing process $[X]$ with Lebesgue decomposition:

$$[X](t) = [X]^c(t) + \sum_{s \leq t} \Delta X(s)^t \Delta X(s).$$

The pathwise Itô formula (1.2) can be applied to this class of processes, which is strictly larger than the class of semimartingales [10].

Theorem 4.1. Let \mathbf{X} be an \mathbb{R}^d -valued càdlàg process, define a sequence of $(\mathcal{D}^{d \times d}, \delta)$ -valued random variables (\mathbf{q}_n) by

$$\mathbf{q}_n(t) := \sum_{\pi_n \ni t_i \leq t} (\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i))(\mathbf{X}(t_{i+1}) - \mathbf{X}(t_i))^T$$

then the following properties are equivalent:

- (i) \mathbf{X} is a process with finite quadratic variation.
- (ii) (\mathbf{q}_n) converges in probability.
- (iii) (\mathbf{q}_n) is a Cauchy sequence in probability.

In addition,

- iv If (\mathbf{q}_n) converges in probability, the limit is $[\mathbf{X}]$.
- v The convergence of (\mathbf{q}_n) to $[\mathbf{X}]$ is UCP if and only if \mathbf{X} is a continuous process of quadratic variation $[\mathbf{X}]$.
- vi (\mathbf{q}_n) converges (resp. is a Cauchy sequence) in probability if and only if each component sequence of (\mathbf{q}_n) converges (resp. is a Cauchy sequence) in probability.

Proof. We first remark that $(\mathcal{D}^{d \times d}, \delta)$ is a complete separable metric space [16], hence by [13, Lemma 9.2.4], the Cauchy property is equivalent to convergence in probability. By [13, Thm. 9.2.1], we can pass to subsequences and apply Prop. 3.5, Thm. 3.6 & Cor. 3.7 pathwise to \mathbf{X} , the claims follow. \square

Acknowledgments. We thank Rafal Lochowski, Pietro Siorpaes and the referee for useful comments. This project was funded by the EPSRC Doctoral Grant 1824430 ‘Analysis and control of path-dependent random systems’.

References

- [1] Anna Ananova and Rama Cont, *Pathwise integration with respect to paths of finite quadratic variation*, Journal de Mathématiques Pures et Appliquées **107** (2017), 737–757. MR-3650323
- [2] Jean Bertoin, *Temps locaux et intégration stochastique pour les processus de Dirichlet*, Séminaire de Probabilités, XXI, Lecture Notes in Math., vol. 1247, Springer, Berlin, 1987, pp. 191–205. MR-0941983
- [3] Patrick Billingsley, *Convergence of probability measures*, Wiley, 1968. MR-0233396
- [4] Rama Cont, *Functional Ito Calculus and functional Kolmogorov equations*, Stochastic Integration by Parts and Functional Ito Calculus (Lecture Notes of the Barcelona Summer School in Stochastic Analysis, July 2012), Birkhauser Basel, 2016, pp. 115–208. MR-3497715
- [5] Rama Cont and David Fournié, *A functional extension of the Ito formula*, C. R. Math. Acad. Sci. Paris **348** (2009), no. 1-2, 57–61. MR-2586744
- [6] Rama Cont and David-Antoine Fournié, *Change of variable formulas for non-anticipative functionals on path space*, J. Funct. Anal. **259** (2010), no. 4, 1043–1072. MR-2652181
- [7] Rama Cont and Yi Lu, *Weak approximation of martingale representations*, Stochastic Process. Appl. **126** (2016), no. 3, 857–882. MR-3452815
- [8] Rama Cont and Nicholas Perkowski, *Pathwise integration and change of variable formulas for continuous paths with arbitrary regularity*, Transactions of the American Mathematical Society (2018), to appear.
- [9] François Coquet, Adam Jakubowski, Jean Mémin, and Leszek Słominski, *Natural decomposition of processes and weak Dirichlet processes*, pp. 81–116, Springer, Berlin, Heidelberg, 2006. MR-2276891
- [10] François Coquet, Jean Mémin, and Leszek Słominski, *On non-continuous dirichlet processes*, Journal of Theoretical Probability **16** (2003), 197–216. MR-1956828
- [11] Mark Davis, Jan Obloj, and Pietro Siorpaes, *Pathwise stochastic calculus with local times*, Ann. Inst. Henri Poincaré Probab. Stat. **54** (2018), no. 1, 1–21. MR-3765878
- [12] Joseph Doob, *Measure theory*, Springer, 1994. MR-1253752
- [13] Richard M. Dudley, *Real analysis & probability*, Cambridge University Press, 2002. MR-1932358
- [14] H. Föllmer, *Calcul d’Itô sans probabilités*, Seminar on Probability, XV (Univ. Strasbourg, Strasbourg, 1979/1980) (French), Lecture Notes in Math., vol. 850, Springer, Berlin, 1981, pp. 143–150. MR-0622559
- [15] Hans Föllmer, *Dirichlet processes*, Stochastic Integrals: Proceedings of the LMS Durham Symposium, July 7–17, 1980 (David Williams, ed.), Springer, Berlin, 1981, pp. 476–478. MR-0621001

On pathwise quadratic variation for càdlàg functions

- [16] Jean Jacod and Albert Shiryaev, *Limit theorems for stochastic processes*, Springer, 2003. MR-1943877
- [17] Adam Jakubowski, *On the Skorokhod topology*, Annales de l'I.H.P. Probabilités et statistiques **22** (1986), no. 3, 263–285 (eng). MR-0871083
- [18] Rafal M. Lochowski, Nicolas Perkowski, and David J. Prömel, *A superhedging approach to stochastic integration*, Stochastic Processes and their Applications (2018).
- [19] Nicolas Perkowski and David J. Prömel, *Local times for typical price paths and pathwise Tanaka formulas*, Electron. J. Probab. **20** (2015), no. 46, 15. MR-3339866
- [20] Alexander Schied, *On a class of generalized Takagi functions with linear pathwise quadratic variation*, J. Math. Anal. Appl. **433** (2016), no. 2, 974–990. MR-3398747
- [21] Anatoly V. Skorokhod, *Limit theorems for stochastic processes*, Theory of Probability & Its Applications **1** (1956), no. 3, 261–290. MR-0084897
- [22] Vladimir Vovk, *Itô calculus without probability in idealized financial markets.*, Lith. Math. J. **55** (2015), no. 2, 270–290 (English). MR-3347596