

THE LARGEST REAL EIGENVALUE IN THE REAL GINIBRE ENSEMBLE AND ITS RELATION TO THE ZAKHAROV–SHABAT SYSTEM

BY JINHO BAIK¹ AND THOMAS BOTHNER²

¹*Department of Mathematics, University of Michigan, baik@umich.edu*

²*Department of Mathematics, King’s College London, thomas.bothner@kcl.ac.uk*

The real Ginibre ensemble consists of $n \times n$ real matrices \mathbf{X} whose entries are i.i.d. standard normal random variables. In sharp contrast to the complex and quaternion Ginibre ensemble, real eigenvalues in the real Ginibre ensemble attain positive likelihood. In turn, the spectral radius $R_n = \max_{1 \leq j \leq n} |z_j(\mathbf{X})|$ of the eigenvalues $z_j(\mathbf{X}) \in \mathbb{C}$ of a real Ginibre matrix \mathbf{X} follows a different limiting law (as $n \rightarrow \infty$) for $z_j(\mathbf{X}) \in \mathbb{R}$ than for $z_j(\mathbf{X}) \in \mathbb{C} \setminus \mathbb{R}$. Building on previous work by Rider and Sinclair (*Ann. Appl. Probab.* **24** (2014) 1621–1651) and Poplavskiy, Tribe and Zaboronski (*Ann. Appl. Probab.* **27** (2017) 1395–1413), we show that the limiting distribution of $\max_{j: z_j \in \mathbb{R}} z_j(\mathbf{X})$ admits a closed-form expression in terms of a distinguished solution to an inverse scattering problem for the Zakharov–Shabat system. As byproducts of our analysis, we also obtain a new determinantal representation for the limiting distribution of $\max_{j: z_j \in \mathbb{R}} z_j(\mathbf{X})$ and extend recent tail estimates in (*Ann. Appl. Probab.* **27** (2017) 1395–1413) via non-linear steepest descent techniques.

1. Introduction and statement of results. This paper is foremost concerned with the derivation of an integrable system for the limiting distribution function

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(\max_{j: z_j \in \mathbb{R}} z_j(\mathbf{X}) \leq \sqrt{n} + t \right), \quad t \in \mathbb{R},$$

of the largest real eigenvalue of a random matrix $\mathbf{X} \in \mathbb{R}^{n \times n}$ chosen from the real Ginibre ensemble.

DEFINITION 1.1 (Ginibre [23] (1965)). A random matrix $\mathbf{X} \in \mathbb{R}^{n \times n}$ is said to belong to the real Ginibre ensemble (GinOE) if its entries are independently chosen with pdf’s

$$\frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x_{jk}^2}, \quad 1 \leq j, k \leq n.$$

Equivalently, the joint pdf of all the independent entries equals

$$f(\mathbf{X}) = \prod_{1 \leq j, k \leq n} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x_{jk}^2} = (2\pi)^{-\frac{1}{2}n^2} e^{-\frac{1}{2} \sum_{j, k=1}^n x_{jk}^2} = (2\pi)^{-\frac{1}{2}n^2} e^{-\frac{1}{2} \text{tr}(\mathbf{X}\mathbf{X}^T)}.$$

The GinOE displays certain similarities to the classical Gaussian Orthogonal Ensemble (GOE) but the presence of both, real and complex eigenvalues introduces also new phenomena. For instance, on a global scale, Wigner’s semicircle law in the GOE is replaced by the following circular law [15]: let

$$\mu_{\mathbf{X}}(s, t) = \frac{1}{n} \#\{1 \leq j \leq n : \Re z_j(\mathbf{X}) \leq s, \Im z_j(\mathbf{X}) \leq t\}, \quad s, t \in \mathbb{R}$$

Received August 2018; revised June 2019.

MSC2010 subject classifications. Primary 60B20; secondary 45M05, 60G70.

Key words and phrases. Real Ginibre ensemble, extreme value statistics, Riemann–Hilbert problem, Zakharov–Shabat system, inverse scattering theory, Deift–Zhou nonlinear steepest descent method.

denote the empirical spectral distribution of the eigenvalues $\{z_j(\mathbf{X})\}_{j=1}^n$ of a matrix $\mathbf{X} \in \text{GinOE}$, then the random measure $\mu_{\mathbf{X}}/\sqrt{n}$ converges almost surely (as $n \rightarrow \infty$) to the uniform distribution on the unit disk; see Figure 1 below.

REMARK 1.2. The circular law is a universal limiting law: it holds true for any $n \times n$ random matrix \mathbf{X} whose entries are i.i.d. complex random variables with mean zero and variance one; see [31] and references therein to the long and rich history of the circular law.

On a local scale, Figure 1 indicates that fluctuations of the spectral radius $R_n = \max_{1 \leq j \leq n} |z_j(\mathbf{X})|$ around \sqrt{n} behave differently depending on whether $z_j \in \mathbb{R}$ or $z_j \in \mathbb{C} \setminus \mathbb{R}$. And indeed, the above mentioned saturn effect was quantified recently and the following central limit theorem derived.

THEOREM 1.3 (Rider, Sinclair [28] (2014); Poplavskiy, Tribe, Zaboronski [27] (2017)). Let $\{z_j(\mathbf{X})\}_{j=1}^n$ denote the eigenvalues of a $n \times n$ random matrix $\mathbf{X} \in \text{GinOE}$. Then

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(\max_{j: z_j \in \mathbb{C} \setminus \mathbb{R}} |z_j(\mathbf{X})| \leq \sqrt{n} + \sqrt{\frac{\gamma_n}{4}} + \frac{t}{\sqrt{4\gamma_n}} \right) = e^{-\frac{1}{2}e^{-t}}, \quad t \in \mathbb{R}$$

with $\gamma_n = \ln(n/(2\pi(\ln n)^2))$. In addition,

$$(1.1) \quad \lim_{n \rightarrow \infty} \mathbb{P} \left(\max_{j: z_j \in \mathbb{R}} z_j(\mathbf{X}) \leq \sqrt{n} + t \right) = \sqrt{\det(1 - T\chi_t \upharpoonright_{L^2(\mathbb{R})})\Gamma_t}, \quad t \in \mathbb{R},$$

where χ_t is the operator of multiplication by $\chi_{(t, +\infty)}(x)$, the characteristic function of $(t, +\infty) \subset \mathbb{R}$, and $T : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ the trace-class integral operator with kernel

$$(1.2) \quad T(x, y) = \frac{1}{\pi} \int_0^\infty e^{-(x+u)^2} e^{-(y+u)^2} du.$$

Moreover,

$$(1.3) \quad \Gamma_t = 1 - \int_t^\infty G(x)((1 - T\chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1}g)(x) dx$$

with $g(x) = \frac{1}{\sqrt{\pi}}e^{-x^2}$ and $G(x) = \int_{-\infty}^x g(y) dy$.

REMARK 1.4. The limit (1.1) was first established in [28]; however, the result contained an algebraic error in the formula for Γ_t . This mistake was subsequently fixed in [27] which resulted in the compact formula (1.3).




FIG. 1. The circular law for 1000 real (rescaled) Ginibre matrices of varying dimensions $n \times n$ in comparison with the unit circle boundary. We plot $n = 2, 4, 8, 16$ from left to right. A saturn effect is clearly visible on the real line.

When compared to the GOE, (1.1) plays the analogue of the celebrated Tracy–Widom edge law for the largest eigenvalue λ_{\max} , cf. [33]. Indeed we recall that in the GOE, as $n \rightarrow \infty$,

$$\lambda_{\max} \Rightarrow \sqrt{2n} + \frac{1}{\sqrt{2n}^{\frac{1}{6}}} F_1,$$

with the cdf

$$(1.4) \quad F_1(t) = \sqrt{\det(1 - K \chi_t \upharpoonright_{L^2(\mathbb{R})})} \widehat{\Gamma}_t, \quad t \in \mathbb{R},$$

where $K : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ is the integral operator with Airy kernel $K(x, y) = \int_0^\infty \text{Ai}(x + u) \text{Ai}(y + u) du$ in terms of the Airy function $\text{Ai}(z)$, [26]. Moreover (see, e.g., [21], Section 9.7),

$$(1.5) \quad \begin{aligned} \widehat{\Gamma}_t &= 1 - \int_t^\infty A(x) ((1 - K \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1} \text{Ai})(x) dx; \\ A(x) &= \int_{-\infty}^x \text{Ai}(y) dy. \end{aligned}$$

1.1. *An integrable system for (1.1).* The formal similarities between (1.4), (1.5) and (1.1), (1.3) are quite obvious, still while the operator K is of integrable type in the sense of [25], that is, has a kernel of the form

$$(1.6) \quad K(x, y) = \frac{\mathbf{f}^\top(x) \mathbf{g}(y)}{x - y}, \quad \mathbf{f}(z) = \begin{bmatrix} \text{Ai}(z) \\ -\text{Ai}'(z) \end{bmatrix}, \quad \mathbf{g}(z) = \begin{bmatrix} \text{Ai}'(z) \\ \text{Ai}(z) \end{bmatrix},$$

this is not true for T with kernel (1.2); see explicitly [28], Section 4. For this reason, neither the standard Tracy–Widom method [32] used in the derivation of an integrable system (a.k.a. a closed-form expression) for the limiting distribution function (1.1) nor the Riemann–Hilbert problem based techniques of Borodin and Deift [9] are directly applicable. However, as we will show below, the situation with (1.2) is not too bad, since the operator $T \chi_t$ is of integrable type up to Fourier conjugation; see Proposition 3.3 below. This observation combined with certain additional manipulations for the Fredholm determinant and the factor Γ_t in (1.1) (see Sections 2, 3 and 4 below) yields an explicit integrable system for $F(t)$ and a subsequent closed-form, Tracy–Widom-like, formula. In fact, we shall state the sought after closed-form expression for the following generalization of (1.1) that contains a generating function parameter $\gamma \in [0, 1]$,

$$(1.7) \quad \begin{aligned} F(t; \gamma) &:= \sqrt{\det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})} \Gamma_{t\gamma} \\ &\text{with } \Gamma_{t\gamma} := 1 - \gamma \int_t^\infty G(x) ((1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1} g)(x) dx. \end{aligned}$$

The main result of our paper (see Theorem 1.6 below) is a closed-form expression for $F(t; \gamma)$ in terms of a distinguished solution to an inverse scattering problem for the Zakharov–Shabat (ZS) system [1, 34]. As it is standard in scattering theory, we shall formulate this inverse problem as a Riemann–Hilbert problem (RHP):

RIEMANN–HILBERT PROBLEM 1.5. *For any $(x, \gamma) \in \mathbb{R} \times [0, 1]$, determine $\mathbf{X}(z) = \mathbf{X}(z; x, \gamma) \in \mathbb{C}^{2 \times 2}$ such that:*

(1) $\mathbf{X}(z)$ is analytic for $z \in \mathbb{C} \setminus \mathbb{R}$ and has a continuous extension on the closed upper and lower half-planes.

(2) The limiting values $\mathbf{X}_\pm(z) = \lim_{\epsilon \downarrow 0} \mathbf{X}(z \pm i\epsilon)$, $z \in \mathbb{R}$ satisfy the jump condition

$$(1.8) \quad \mathbf{X}_+(z) = \mathbf{X}_-(z) \begin{bmatrix} 1 - |r(z)|^2 & -\bar{r}(z)e^{-2ixz} \\ r(z)e^{2ixz} & 1 \end{bmatrix},$$

$$z \in \mathbb{R} \text{ with } r(z) = r(z; \gamma) = -i\sqrt{\gamma}e^{-\frac{1}{4}z^2}.$$

(3) As $z \rightarrow \infty$, we require the normalization

$$\mathbf{X}(z) = \mathbb{I} + \mathbf{X}_1 z^{-1} + \mathbf{X}_2 z^{-2} + \mathcal{O}(z^{-3}); \quad \mathbf{X}_i = \mathbf{X}_i(x, \gamma) = [X_i^{jk}(x, \gamma)]_{j,k=1}^2.$$

Note that $r(z; \gamma) \in \mathcal{S}(\mathbb{R})$, the Schwartz space on the line, but $\|r\|_\infty = \sup_{z \in \mathbb{R}} |r(z)| < 1$ only for $\gamma \in [0, 1)$. Hence $r(z; \gamma)$, the so-called reflection coefficient, does not belong to the standard Beals–Coifman class of reflection coefficients (cf. [3, 4]) in the case (1.1) most relevant to the GinOE. For this reason, we will prove unique solvability of RHP 1.5 for all $(x, \gamma) \in \mathbb{R} \times [0, 1]$, and thus also existence of the coefficients $\mathbf{X}_i(x, \gamma)$ directly in the sections below. We now present our main result.

THEOREM 1.6. For any $(t, \gamma) \in \mathbb{R} \times [0, 1]$,

$$(1.9) \quad (F(t; \gamma))^2 = \exp \left[-\frac{1}{4} \int_t^\infty (x-t) \left| y \left(\frac{x}{2}; \gamma \right) \right|^2 dx \right] \\ \times \{ \cosh \mu(t; \gamma) - \sqrt{\gamma} \sinh \mu(t; \gamma) \},$$

using the abbreviation

$$\mu(t; \gamma) := -\frac{i}{2} \int_t^\infty y \left(\frac{x}{2}; \gamma \right) dx,$$

and where $y = y(x; \gamma) : \mathbb{R} \times [0, 1] \rightarrow i\mathbb{R}$ equals $y(x; \gamma) := 2iX_1^{12}(x, \gamma)$ in terms of the matrix coefficient $\mathbf{X}_1(x, \gamma)$ in condition (3) of RHP 1.5 above.

Identity (1.9) is the analogue of the Tracy–Widom Painlevé-II formula for $F_1(t)$ in case $\gamma = 1$; see [33]. For $\gamma \in [0, 1)$, our definition (1.7) is motivated by the generating function of the soft-edge scaled $(t, +\infty)$ gap probabilities for the superimposed (cf. [21], Section 6.6) orthogonal ensemble

$$\text{odd}(\text{OE}_n(e^{-x^2}) \cup \text{OE}_n(e^{-x^2})).$$

In this context, (1.9) is the direct analogue of [21], (9.150), modulo the replacement of the Painlevé-II transcendent with the above solution entry $y(x; \gamma)$ of RHP 1.5. Also, somewhat similar formulæ to the type of (1.9) appear in the computation of edge eigenvalue gap probabilities in the GOE and Gaussian Symplectic Ensemble (GSE) in terms of Painlevé transcendents; see [13].

REMARK 1.7. Another possible motivation for the introduction of γ in (1.7) could arise from studying a thinned version of the real eigenvalues in the GinOE, that is, from analyzing the point process

$$\mathfrak{X}^\gamma = \{z_j^\gamma(\mathbf{X}) : z_j^\gamma(\mathbf{X}) \in \mathbb{R}\}_{j=1}^{n(\gamma)}, \quad 1 \leq n(\gamma) \leq n,$$

obtained from $\mathfrak{X} = \{z_j(\mathbf{X}) \in \mathbb{R}\}_{j=1}^n$ with $\mathbf{X} \in \text{GinOE}$ by independently removing each real eigenvalue with likelihood $\gamma \in [0, 1]$. For GOE, the largest eigenvalue distribution function after thinning admits a Painlevé closed-form expression (see [10], (1.6)) in case of GinOE the corresponding result is unknown. It is also not immediately clear whether (1.7) has a probabilistic interpretation in the thinned superimposed orthogonal ensemble. We plan to address this question in a future publication.

REMARK 1.8. We prove existence of $y(x; \gamma)$ for $(x, \gamma) \in \mathbb{R} \times [0, 1]$ in Theorem 3.9 and continuity of $y(x; \gamma), x \in \mathbb{R}$ for any fixed $\gamma \in [0, 1]$ in Lemma 2.1 and Corollary 3.6. Moreover, we show that $y(x; \gamma)$ for $(x, \gamma) \in \mathbb{R} \times [0, 1]$ is purely imaginary and

$$(1.10) \quad y(x; \gamma) = 2i\sqrt{\frac{\gamma}{\pi}}e^{-4x^2}(1 + \mathcal{O}(e^{-4x^2})), \quad x \rightarrow +\infty,$$

that is, the right-hand side in (1.9) is well-defined.

Returning to the aforementioned comparison between GinOE and GOE, we see from (1.9) and [33], (53), that, overall, the main difference in GinOE arises from the presence of the inverse scattering type RHP 1.5 and its solution entry $y(x; \gamma)$ instead of the more common Painlevé transcendents in the Gaussian invariant ensembles. For this reason, we shall briefly review a few selected aspects of the integrability theory of RHP 1.5; see [1, 3, 4, 34] for more details.

1.2. *The Zakharov–Shabat system in a nutshell.* Note that

$$\Psi(z) := \mathbf{X}(z)e^{-ixz\sigma_3}, \quad z \in \mathbb{C} \setminus \mathbb{R}$$

solves a RHP with a x -independent jump on \mathbb{R} , thus $\frac{\partial \Psi}{\partial x} \Psi^{-1}$ is an entire function. In fact, using condition (3) in RHP 1.5 and Liouville’s theorem, we find

$$(1.11) \quad \frac{\partial \Psi}{\partial x} = \left\{ -iz\sigma_3 + 2i \begin{bmatrix} 0 & X_1^{12} \\ -X_1^{21} & 0 \end{bmatrix} \right\} \Psi.$$

But since RHP 1.5 enjoys the symmetry

$$(1.12) \quad \mathbf{X}(z; x, \gamma) = \sigma_1 \overline{\mathbf{X}(\bar{z}; x, \gamma)} \sigma_1, \quad z \in \mathbb{C} \setminus \mathbb{R}; \sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix},$$

we learn that $X_i^{11}(x, \gamma) = \overline{X_i^{22}(x, \gamma)}$ as well as $X_i^{21}(x, \gamma) = \overline{X_i^{12}(x, \gamma)}$, and thus with $y = 2iX_1^{12}$ from (1.11),

$$(1.13) \quad \frac{\partial \Psi}{\partial x} = \left\{ -iz\sigma_3 + \begin{bmatrix} 0 & y \\ \bar{y} & 0 \end{bmatrix} \right\} \Psi \equiv \mathbf{U}(z; x, \gamma) \Psi.$$

This celebrated first-order system, known as the ZS-system, is directly related to several of the most interesting nonlinear evolution equations in $1 + 1$ dimensions which are solvable by the inverse scattering method. For instance, in order to solve the Cauchy problem for the defocusing nonlinear Schrödinger equation,

$$(1.14) \quad \begin{aligned} iy_t + y_{xx} - 2|y|^2y &= 0, & y(x, 0) &= y_0(x) \in \mathcal{S}(\mathbb{R}); \\ y &= y(x, t) : \mathbb{R}^2 \rightarrow \mathbb{C}, \end{aligned}$$

one first computes the reflection coefficient $r(z) \in \mathcal{S}(\mathbb{R})$ associated to the initial data y_0 through the direct scattering transform. A basic fact of the scattering theory for the Zhakarov–Shabat system (1.13) states that this transform, that is, the map $y_0 \rightarrow r$, is a bijection from $\mathcal{S}(\mathbb{R})$ onto $\mathcal{S}(\mathbb{R}) \cap \{r : \|r\|_\infty = \sup_{z \in \mathbb{R}} |r(z)| < 1\}$; cf. [3]. Second, one considers RHP 1.5 above subject to the replacement

$$e^{2ixz} \rightarrow e^{2i(2tz^2+xz)}, \quad t \in \mathbb{R},$$

and provided this problem is solvable, its (unique) solution in turn leads to a solution of (1.14) with $y(x, 0) = y_0(x)$ via the formula $y(x, t) = 2iX_1^{12}(x, t)$. Thus, in order to solve (1.14), one must solve the t -modified RHP 1.5 (a.k.a. the inverse scattering transform) for

the given reflection coefficient $r(z)$, determined under the aforementioned bijection $y_0 \rightarrow r$. Returning now to our context, we see that (1.9) therefore depends on a distinguished solution $y(x; \gamma)$ of the inverse scattering transform for the Zakharov–Shabat system (1.13) subject to the reflection coefficient $r(z; \gamma) = -i\sqrt{\gamma}e^{-\frac{1}{4}z^2}$.

REMARK 1.9. As outlined above, the operator $T\chi_t$ is of integrable type once viewed in Fourier space. This idea was first used in the analysis of single and multitime processes in [5, 6]. Specifically, [5, 6] showed that certain matrix Fredholm determinants are expressible as determinants of integrable matrix kernels, and thus connected to RHPs. As a direct application of this technique, Bertola and Cafasso rederived for instance the Adler–van Moerbeke PDE for the joint distributions of the Airy-2 process by Riemann–Hilbert techniques. Our approach to the GinOE and (1.7) is clearly inspired by these works.

1.3. *Tail asymptotics.* The advantage of the exact formula (1.9) lies in the fact that $y = y(x; \gamma)$ admits a Riemann–Hilbert formulation as outlined in RHP 1.5. Thus its large space/long time behavior can be systematically computed via nonlinear steepest descent techniques [12] and this path leads the way to large tail estimates for (1.7). We summarize our second result.

COROLLARY 1.10. *Let $\gamma \in [0, 1]$ and $F(t; \gamma)$ be defined as in (1.7). Then, as $t \rightarrow +\infty$,*

$$(1.15) \quad F(t; \gamma) = 1 - \frac{\gamma}{4} \operatorname{erfc}(t) + \mathcal{O}(\gamma^{\frac{3}{2}} t^{-1} e^{-2t^2}),$$

in terms of the complementary error function $\operatorname{erfc}(z)$; cf. [26], 7.2.2. On the other hand, as $t \rightarrow -\infty$,

$$(1.16) \quad F(t; \gamma) = e^{\eta_1(\gamma)t} \eta_0(\gamma)(1 + o(1)), \quad \eta_1(\gamma) = \frac{1}{2\sqrt{2\pi}} \operatorname{Li}_{\frac{3}{2}}(\gamma),$$

in terms of the polylogarithm $\operatorname{Li}_s(z)$ (cf. [26], 25.12.10) and with a t -independent positive factor $\eta_0(\gamma)$.

Estimate (1.15) for $\gamma = 1$ is standard and known from [20], say. The leading order in (1.16) for $\gamma = 1$ was obtained by Forrester [22] and also by Poplavskiy, Tribe and Zaboronski [27] using an interesting connection to coalescence processes. Here, we employ nonlinear steepest descent techniques to confirm (1.15) for all $\gamma \in [0, 1]$ and derive (1.16) for $\gamma \in [0, 1)$.

The paper [22] also discusses the thinned version of the real eigenvalues, or equivalently, the generating function for the probability of the number of eigenvalues. Namely, the discussions in the third paragraph of [22], Section 4, claim a leading order term which, after evaluating the integral [22], (4.1), explicitly, is the same as our $\eta_1(\gamma)$ with $\gamma = 2\xi - \xi^2$ where $1 - \xi$ denotes the removal probability of an eigenvalue. As mentioned above in Remark 1.7, it is interesting to consider the relationship between our $F(t; \gamma)$ and the thinned version.

REMARK 1.11. As can be seen from (1.16), the left tails of $F(t; \gamma)$ in the leading order decay to zero exponentially fast for all $\gamma \in (0, 1]$. This is in sharp contrast to the GOE where $\ln F_1(t) \sim \frac{1}{24}t^3$ as $t \rightarrow -\infty$ and $\ln F_1(t; \gamma) \sim -\frac{2v}{3\pi}(-t)^{3/2}$ for $\gamma \in (0, 1)$ fixed with $v = -\ln(1 - \gamma) < +\infty$; cf. [10]. This means that the large negative x behavior of $\mathfrak{S}(y(\frac{x}{2}; \gamma))$ cannot be as sensitive to a small change in $\gamma \in (0, 1]$ near $\gamma = 1$ as the corresponding behavior for the Painlevé-II transcendent $u(x; \gamma)$; see Figure 4 below for a visualization.




FIG. 2. The distribution function $F(t)$ of the largest real GinOE eigenvalue in red versus GOE Tracy–Widom $F_1(t)$ in blue. The plots are generated in MATLAB with $m = 50$ quadrature points using the Nyström method with Gauss–Legendre quadrature. On the left cdfs, on the right pdfs.

1.4. *Numerical comparison.* The closed-form expression (1.9) is not as optimal for numerical purposes as a single Fredholm determinant formula, compare the discussions in [8]. For this reason, we derive a new determinantal formula for the limiting distribution of $\max_{j:z_j \in \mathbb{R}} z_j(\mathbf{X})$ in our third result below. This identity is completely analogous to the Ferrari–Spohn formula [16] in the GOE.

THEOREM 1.12. Let $F(t)$ be defined as in (1.1). Then

$$(1.17) \quad F(t) = \det(1 - S\chi_t \upharpoonright_{L^2(\mathbb{R})}),$$

where $S : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ is the integral operator with kernel

$$(1.18) \quad S(x, y) = \frac{1}{2\sqrt{\pi}} e^{-\frac{1}{4}(x+y)^2}.$$

Identity (1.17) allows us to numerically simulate several statistical quantities of $\max_{j:z_j \in \mathbb{R}} z_j(\mathbf{X})$ by implementing Bornemann’s algorithm [8] in MATLAB. In more detail, we discretize the determinant (1.17) by the Nyström method using a Gauss–Legendre quadrature rule with $m = 50$ quadrature points. Once the values of the cdf are then numerically accessible, computing associated quantities such as moments is straightforward. We summarize a few values in Table 1 below.

In the upcoming figures, we first plot the distribution function $F(t)$ of the largest real eigenvalue in the GinOE in comparison to $F_1(t)$, the cdf of the largest eigenvalue in the GOE; compare Figure 2. After that, we compare the asymptotic expansions (1.15) and (1.16) to our numerical results in Figure 3.

A closed-form computation of $\eta_0(\gamma)$ in (1.16) is beyond the methods developed in this paper. In [22], (2.26), (2.30), Forrester derives a closed-form series representation for $\eta_0(1)$,

TABLE 1
Some moments of the GinOE in comparison to GOE moments

Ensemble	Mean	Variance	Skewness	Kurtosis
GinOE (1.1)	−1.30319	3.97536	−1.76969	5.14560
GOE (1.4)	−1.20653	1.60778	0.29346	0.16524




FIG. 3. We double-check (1.15) on the left and (1.16) on the right (in a semilogarithmic plot) against the numerically computed values of $F(t; 1)$ based on (1.17) with $\eta_0(1) = 0.75277069$. Once more, we have used the Nyström method with a Gauss–Legendre quadrature rule and $m = 50$ quadrature points.

namely

$$\exp\left[\ln 2 - \frac{1}{4} + \frac{1}{4\pi} \sum_{n=2}^{\infty} \frac{1}{n} \left(-\pi + \sum_{m=1}^{n-1} \frac{1}{\sqrt{m(n-m)}}\right)\right] \approx 1.06470738.$$

However, after using a simple approximation for $\eta_0(1)$ obtained by numerically computing the ratio of $F(t; 1)$ from (1.17) and $e^{\eta_1(1)t}$ for large negative t , our result

$$\eta_0(1) = 0.75277069,$$

does not match [22], (2.26), (2.30). This discrepancy needs to be further investigated.¹ Finally, in the remaining Figure 4 we showcase the qualitatively different asymptotic behaviors of $\Im(y(\frac{x}{2}; \gamma))$ on one hand and $u(x; \gamma)$ on the other, compare Remark 1.11.

1.5. *Outline of paper.* Toward the end of our Introduction, we now offer a short outline for the remaining sections of the paper. In Section 2, we first summarize a few basic properties of the operator T on $L^2(\mathbb{R})$ with kernel (1.2) and show that $T\chi_t$ is indeed of integrable type




FIG. 4. We compare $\Im(y(\frac{x}{2}; \gamma))$ on the left to $u(x; \gamma)$ on the right for varying values of γ . While the solution entry $y(x; \gamma)$ to RHP 1.5 decays to zero as $x \rightarrow -\infty$ for all $\gamma \in (0, 1]$, this is not true for the Painlevé-II transcendent.

¹After the submission of this manuscript, there appeared the preprint [17] in which the authors derive an explicit formula for $\eta_0(1)$; see [17], (2), (3). Their result aligns with our above numerical prediction.

[25], up to Fourier conjugation. Still, instead of deriving an integrable system for $F(t; \gamma)$ at this point we employ further simplification steps in Section 3 that allow to match the thereby obtained RHP 3.4 almost immediately with RHP 1.5. This will in turn prove the first part of Theorem 1.6 in (3.26) below once combined with appropriate right tail estimates that we derive by nonlinear steepest descent arguments; compare Section 3.4. These steps are then followed up in Section 4 by an explicit evaluation of $\Gamma_{t\gamma}$ in (1.7) in terms of Riemann–Hilbert data, and the second part in (1.9) is then also proven. While carrying out the aforementioned steps, we derive estimate (1.15) en route and complete the proof of Corollary 1.10 afterwards in Section 5. The nonlinear steepest descent techniques for the left tail are standard except for the appearance of certain collapsing jump contours. For this reason, we provide the necessary small norm estimates of the underlying (unbounded) Cauchy operators in Appendix A. The paper closes with the derivation of (1.17) in Section 6 which heavily relies on the proof technique presented in [16] for the corresponding GOE result.

2. Preliminary steps. We begin with the following result which is standard for, say, the Airy operator (see, for instance, [2], Lemma 6.15) but which does not appear in the literature for T_{χ_t} , to the best of our knowledge.

LEMMA 2.1. *For every $t \in \mathbb{R}$, the self-adjoint operator T_{χ_t} satisfies $0 \leq T_{\chi_t} \leq 1$, and thus $\|T_{\chi_t}\| \leq 1$. Moreover, $1 - \gamma T_{\chi_t}$ is invertible on $L^2(\mathbb{R})$ for all $\gamma \in [0, 1]$.*

PROOF. Recall the Gaussian integral

$$(2.1) \quad e^{-x^2} = \frac{1}{2\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-\frac{1}{4}y^2 + ixy} \, dy, \quad x \in \mathbb{R},$$

and note that for $f \in L^2(\mathbb{R})$,

$$(2.2) \quad \begin{aligned} 0 \leq \langle f, T_{\chi_t} f \rangle_{L^2(\mathbb{R})} &= \frac{1}{\pi} \int_0^{\infty} \left| \int_{-\infty}^{\infty} e^{-(x+u)^2} f_t(x) \, dx \right|^2 \, du \\ &\leq \frac{1}{\pi} \int_{-\infty}^{\infty} \left| \int_{-\infty}^{\infty} e^{-(x+u)^2} f_t(x) \, dx \right|^2 \, du, \end{aligned}$$

where we abbreviate $f_t(x) := f(x)\chi_{(t, +\infty)}(x)$. Hence, with (2.1), we compute

$$\int_{-\infty}^{\infty} e^{-(x+u)^2} f_t(x) \, dx = \frac{1}{\sqrt{2}} \int_{-\infty}^{\infty} e^{-\frac{1}{4}y^2} \widehat{f}_t(-y) e^{iuy} \, dy = \sqrt{\pi} \widehat{g}(-u),$$

where $\widehat{f}_t(y) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f_t(x) e^{-iyx} \, dy$ is the Fourier transform of f_t and $g(y) := e^{-\frac{1}{4}y^2} \times \widehat{f}_t(-y)$. Thus together in (2.2),

$$(2.3) \quad \begin{aligned} 0 \leq \langle f, T_{\chi_t} f \rangle_{L^2(\mathbb{R})} &\leq \int_{-\infty}^{\infty} |\widehat{g}(-u)|^2 \, du = \int_{-\infty}^{\infty} |g(y)|^2 \, dy = \int_{-\infty}^{\infty} e^{-\frac{1}{2}y^2} |\widehat{f}_t(-y)|^2 \, dy \\ &\leq \int_{-\infty}^{\infty} |\widehat{f}_t(-y)|^2 \, dy = \int_{-\infty}^{\infty} |f_t(y)|^2 \, dy \leq \int_{-\infty}^{\infty} |f(y)|^2 \, dy \\ &= \langle f, f \rangle_{L^2(\mathbb{R})}, \end{aligned}$$

using Plancherel’s theorem in the first and third equality. Hence $0 \leq T_{\chi_t} \leq 1$ and by self-adjointness also

$$\|T_{\chi_t}\| = \sup_{\|f\|_{L^2(\mathbb{R})}=1} |\langle f, T_{\chi_t} f \rangle_{L^2(\mathbb{R})}| \leq 1.$$




FIG. 5. An admissible choice for the contour Γ in (2.5).

For invertibility, we assume there is $f \in L^2(\mathbb{R})$, not identically zero, such that $\gamma T \chi_t f = f$ with $\gamma \in (0, 1]$. By (2.3), we must therefore have equality

$$\langle f, T \chi_t f \rangle_{L^2(\mathbb{R})} = \gamma \int_{-\infty}^{\infty} |\widehat{g}(-u)|^2 du,$$

but from (2.2) (without estimating) then also

$$\langle f, T \chi_t f \rangle_{L^2(\mathbb{R})} = \gamma \int_0^{\infty} |\widehat{g}(-u)|^2 du.$$

So $\widehat{g}(u) = 0$ for $u < 0$, that is,

$$(2.4) \quad \int_t^{\infty} e^{-(x+u)^2} f(x) dx = 0 \quad \text{for } u < 0.$$

Using analytic properties of the exponential, we then conclude that (2.4) must also hold for $u > 0$ and, therefore, $f \equiv 0$, a contradiction. \square

Our next steps will make use of a slight generalization of (2.1), namely the following contour integral formula: for any smooth nonself-intersecting contour Γ oriented from $\infty \cdot e^{i\alpha}$ to $\infty \cdot e^{i\beta}$ with $\alpha \in (\frac{3\pi}{4}, \frac{5\pi}{4})$ and $\beta \in (-\frac{\pi}{4}, \frac{\pi}{4})$; see, for example, Figure 5, we have

$$(2.5) \quad e^{-x^2} = \frac{1}{2\sqrt{\pi}} \int_{\Gamma} e^{-\frac{1}{4}\lambda^2 \pm ix\lambda} d\lambda, \quad x \in \mathbb{R}.$$

Now fix $(x, y) \in \mathbb{R}^2$ throughout and substitute (2.5) twice into (1.2), once with the (+) sign in (2.5) and once with the (-) sign:

$$T(x, y) = \frac{1}{(2\pi)^2} \int_{\Gamma_\lambda} \int_{\Gamma_w} e^{-\frac{1}{4}(\lambda^2+w^2)} e^{-i(x\lambda-yw)} \left[\int_0^{\infty} e^{-iu(\lambda-w)} du \right] dw d\lambda.$$

So provided we choose $(\lambda, w) \in \Gamma_\lambda \times \Gamma_w$ such that $\Im w > \Im \lambda$,

$$(2.6) \quad T(x, y) = \frac{1}{(2\pi)^2} \int_{\Gamma_\lambda} \int_{\Gamma_w} \frac{e^{-\frac{1}{4}(\lambda^2+w^2) - i(x\lambda-yw)}}{i(\lambda-w)} dw d\lambda.$$

Next, we use the residue theorem.

LEMMA 2.2. Suppose $w \in \Gamma_w$ satisfies $\Im w > 0$. Then for any $y, t \in \mathbb{R} : y \neq t$,

$$(2.7) \quad \frac{1}{2\pi i} \int_{-\infty}^{\infty} e^{i(\mu-w)(y-t)} \frac{d\mu}{\mu-w} = \chi_{(t, +\infty)}(y).$$




FIG. 6. Red contour $\Sigma_+(R)$ in the upper half-plane consisting of two vertical pieces $\Sigma_{1,3}$ (left, right), top piece Σ_2 as well as $[-R, R] \subset \mathbb{R}$.

PROOF. The integral on the left-hand side in (2.7) is well-defined under the given assumptions and can be evaluated as follows: define the entire function $f(\mu) := e^{i\mu(y-t)}$, $\mu \in \mathbb{C}$ and consider for $y > t$ a contour integral of f along the closed rectangle $\Sigma_+(R)$ shown in Figure 6 where $R > |w| + 1$ and $w \in \mathbb{C} : \Im w > 0$ is fixed. By the residue theorem,

$$(2.8) \quad \frac{1}{2\pi i} \oint_{\Sigma_+(R)} f(\mu) \frac{d\mu}{\mu - w} = e^{iw(y-t)},$$

and since with some $C = C(w) > 0$

$$\left| \int_{\Sigma_{1,3}} f(\mu) \frac{d\mu}{\mu - w} \right| \leq \frac{C}{R(y-t)} \rightarrow 0 \quad \text{as well as}$$

$$\left| \int_{\Sigma_2} f(\mu) \frac{d\mu}{\mu - w} \right| \leq C e^{-R(y-t)} \rightarrow 0, \quad \text{as } R \rightarrow +\infty,$$

identity (2.7) follows for $y > t$ from (2.8) in the limit $R \rightarrow +\infty$.

For $y < t$, we evaluate the contour integral (2.8) along an analogous contour $\Sigma_-(R)$ in the lower half-plane, however this time the residue theorem does not yield a nonvanishing contribution. Hence, (2.7) follows also for $y < t$ in the limit $R \rightarrow +\infty$. \square

Let us now choose $\Gamma_\lambda = \mathbb{R}$ in (2.6) and $\Gamma_w \equiv \Gamma$ as any smooth nonself-intersecting contour in the upper w half-plane. Together with Lemma 2.2, we find

$$(2.9) \quad \begin{aligned} & T(x, y) \chi_{(t, +\infty)}(y) \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{e^{-ix\lambda}}{\sqrt{2\pi}} \underbrace{\left[\frac{1}{(2\pi)^2} \int_{\Gamma} \frac{e^{-\frac{1}{4}(\lambda^2 + w^2) - it(\mu - w)}}{(\lambda - w)(w - \mu)} dw \right]}_{=: E(\lambda, \mu)} \frac{e^{iy\mu}}{\sqrt{2\pi}} d\mu d\lambda. \end{aligned}$$

Thus, provided we let $\mathcal{F} : L^1(\mathbb{R}) \cap L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ denote the standard Fourier transform, that is,

$$\widehat{f}(x) \equiv (\mathcal{F}f)(x) := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-ix\lambda} f(\lambda) d\lambda, \quad x \in \mathbb{R},$$

which extends to a unitary operator on $L^2(\mathbb{R})$ by classical theory, then (2.9) shows that $T\chi_t$ on $L^2(\mathbb{R})$ is simply equal to the operator composition $\mathcal{F}E\mathcal{F}^{-1}$ where $E : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ is the integral operator on $L^2(\mathbb{R})$ with kernel $E(\lambda, \mu)$ given in (2.9). In order to verify certain regularity properties of E (and other operators to follow), it will be convenient to abide to the following convention.

CONVENTION 2.3. From now on, we shall think of our operators $T\chi_t$, E and \mathcal{F} as acting, not on $L^2(\mathbb{R}, dx)$, but on an extended space $L^2(\Omega, |d\lambda|)$ (where $\mathbb{R} \subset \Omega \subset \mathbb{C}$ for some

oriented contour Ω to be specified below and $|\mathrm{d}\lambda|$ is the arc-length measure) and to have kernel

$$T_{\text{ext}}(\lambda, \mu) := T(\lambda, \mu)\chi_{(t, +\infty)}(\mu)\chi_{\mathbb{R}}(\lambda), \quad (\lambda, \mu) \in \Omega \times \Omega,$$

as well as $E_{\text{ext}}(\lambda, \mu) := E(\lambda, \mu)\chi_{\mathbb{R}}(\lambda)\chi_{\mathbb{R}}(\mu)$ and $\mathcal{F}_{\text{ext}}(\lambda, \mu) := \frac{1}{\sqrt{2\pi}}e^{-i\lambda\mu}\chi_{\mathbb{R}}(\lambda)\chi_{\mathbb{R}}(\mu)$.

Provided we modify the distributional kernel of the identity in accordance with Convention 2.3, equation (2.9) thus establishes the operator identity,

$$(2.10) \quad \begin{aligned} 1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})} &= 1 - \gamma T_{\text{ext}} \upharpoonright_{L^2(\Omega)} \\ &= \mathcal{F}_{\text{ext}}(1 - \gamma E_{\text{ext}} \upharpoonright_{L^2(\Omega)})\mathcal{F}_{\text{ext}}^{-1}, \quad \gamma \in [0, 1]. \end{aligned}$$

The main motivation behind Convention 2.3 comes from the following result.

LEMMA 2.4. *The operator E_{ext} on $L^2(\Omega = \mathbb{R} \sqcup \Gamma, |\mathrm{d}\lambda|)$ with kernel given in Convention 2.3 is trace-class.*

PROOF. Observe the factorization $E_{\text{ext}} = E_1 E_2$ where $E_1 : L^2(\Omega, |\mathrm{d}\lambda|) \rightarrow L^2(\Omega, |\mathrm{d}\lambda|)$ has kernel

$$E_1(\lambda, w) = \frac{1}{2\pi} \frac{e^{-\frac{1}{4}\lambda^2 - \frac{1}{8}w^2}}{\lambda - w} \chi_{\mathbb{R}}(\lambda)\chi_{\Gamma}(w),$$

and $E_2 : L^2(\Omega, |\mathrm{d}\lambda|) \rightarrow L^2(\Omega, |\mathrm{d}\lambda|)$ has kernel

$$E_2(w, \mu) = \frac{1}{2\pi} \frac{e^{-\frac{1}{8}w^2 - i\mu(w-w)}}{w - \mu} \chi_{\Gamma}(w)\chi_{\mathbb{R}}(\mu).$$

But both, E_1 and E_2 , are Hilbert–Schmidt integral operators on $L^2(\Omega, |\mathrm{d}\lambda|)$ since

$$\int_{\Omega} \int_{\Omega} |E_j(z_1, z_2)|^2 |\mathrm{d}z_1| |\mathrm{d}z_2| < \infty, \quad j = 1, 2,$$

so $E_{\text{ext}} = E_1 E_2$ is trace-class on $L^2(\Omega, |\mathrm{d}\lambda|)$. \square

Observe that E_{ext} is already of integrable type in the sense of [25] once we use partial fractions. Still, it is preferable to massage E_{ext} a bit further: Introduce $M : L^2(\Omega, |\mathrm{d}\lambda|) \rightarrow L^2(\Omega, |\mathrm{d}\lambda|)$ as multiplication by $e^{-\frac{1}{8}\lambda^2} \chi_{\mathbb{R}}(\lambda)$,

$$(Mf)(\lambda) = e^{-\frac{1}{8}\lambda^2} \chi_{\mathbb{R}}(\lambda) f(\lambda), \quad f \in L^2(\Omega, |\mathrm{d}\lambda|),$$

and note that both operators $E_{\text{ext}} = (E_{\text{ext}}M)M^{-1}$ and $N := M^{-1}E_{\text{ext}}M$ are trace-class on $L^2(\Omega, |\mathrm{d}\lambda|)$.

REMARK 2.5. The operator N has kernel

$$N(\lambda, \mu) = \frac{1}{(2\pi)^2} \int_{\Gamma} \frac{e^{-\frac{1}{8}(\lambda^2 + \mu^2) - \frac{1}{4}w^2 + i\mu(w - \mu)}}{(\lambda - w)(w - \mu)} \mathrm{d}w \chi_{\mathbb{R}}(\lambda)\chi_{\mathbb{R}}(\mu),$$

and is therefore trace-class since it can be factorized into $N = N_1 N_2$ where $N_j : L^2(\Omega, |\mathrm{d}\lambda|) \rightarrow L^2(\Omega, |\mathrm{d}\lambda|)$ have Hilbert–Schmidt kernels

$$(2.11) \quad \begin{aligned} N_1(\lambda, w) &= \frac{1}{2\pi} \frac{e^{-\frac{1}{8}(\lambda^2 + w^2)}}{\lambda - w} \chi_{\mathbb{R}}(\lambda)\chi_{\Gamma}(w), \\ N_2(w, \mu) &= \frac{1}{2\pi} \frac{e^{-\frac{1}{8}(w^2 + \mu^2) + i\mu(w - \mu)}}{w - \mu} \chi_{\Gamma}(w)\chi_{\mathbb{R}}(\mu). \end{aligned}$$




FIG. 7. Red contour $\Sigma(R)$ in the upper half-plane consisting of two vertical pieces $\Sigma_{1,3}$ (left, right), top piece Σ_2 as well as $[-R + i\delta, R + i\delta] \subset \mathbb{R} + i\delta$.

Concluding our preliminary steps, we note that by the conjugation invariance of the Fredholm determinant and Sylvester’s determinant identity [24], Chapter IV, (5.9),

$$\begin{aligned}
 & \det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) \\
 (2.12) \quad & \stackrel{(2.10)}{=} \det(1 - \gamma E_{\text{ext}} \upharpoonright_{L^2(\Omega)}) = \det(1 - \gamma (E_{\text{ext}} M) M^{-1} \upharpoonright_{L^2(\Omega)}) \\
 & = \det(1 - \gamma M^{-1} E_{\text{ext}} M \upharpoonright_{L^2(\Omega)}) = \det(1 - \gamma N \upharpoonright_{L^2(\Omega)}).
 \end{aligned}$$

3. Riemann–Hilbert problem and proof of Theorem 1.6, part 1. The seemingly quickest way to derive the first parts of (1.9) makes use of the factorization $N = N_1 N_2$ in (2.11) and a subsequent operator identity that further simplifies our above $1 - N \upharpoonright_{L^2(\Omega)}$.

3.1. *Fredholm determinant identities.* We begin with the following lemma which improves the statement of Remark 2.5 at the cost of further extending Ω .

LEMMA 3.1. *The integral operators $N_j : L^2(\mathring{\Omega}, |d\lambda|) \rightarrow L^2(\mathring{\Omega}, |d\lambda|)$ with kernels (2.11) defined on the extended space $\mathring{\Omega} = \Omega \sqcup (\mathbb{R} + i\delta)$ for some sufficiently small $\delta > 0$ are trace-class.*

PROOF. We keep $\lambda \in \mathbb{R}$ and $w \in \Gamma$ (as before with Γ any smooth nonself-intersecting contour in the upper half-plane) fixed throughout. Now choose $0 < \delta < \Im w$ and observe that

$$(3.1) \quad \frac{1}{2\pi i} \int_{\mathbb{R} + i\delta} \frac{dv}{(\lambda - v)(v - w)} = \frac{1}{\lambda - w}.$$

Indeed, the integral in the left-hand side of (3.1) is well-defined and with $\Sigma(R)$ as shown in Figure 7, where $R > |w| + 1$, we find from residue theorem

$$(3.2) \quad \frac{1}{2\pi i} \oint_{\Sigma(R)} \frac{dv}{(\lambda - v)(v - w)} = \frac{1}{\lambda - w}.$$

But with some $C = C(w, \lambda) > 0$,

$$\begin{aligned}
 \left| \int_{\Sigma_{1,3}} \frac{dv}{(\lambda - v)(v - w)} \right| & \leq \frac{C}{R} \rightarrow 0 \quad \text{as well as} \\
 \left| \int_{\Sigma_2} \frac{dv}{(\lambda - v)(v - w)} \right| & \leq \frac{C}{R} \rightarrow 0, \quad \text{as } R \rightarrow +\infty,
 \end{aligned}$$

so identity (3.1) follows from (3.2) in the limit $R \rightarrow +\infty$. Now use (3.1) and rewrite (2.11) in the following way:

$$N_1(\lambda, w) = \frac{1}{4i\pi^2} \int_{\mathbb{R} + i\delta} \frac{e^{-\frac{1}{8}(\lambda^2 + w^2)}}{2\pi(\lambda - v)(v - w)} dv \chi_{\mathbb{R}}(\lambda) \chi_{\Gamma}(w),$$

so that $N_1 = N_{11}N_{12}$ where $N_{1j} : L^2(\mathring{\Omega}, |d\lambda|) \rightarrow L^2(\mathring{\Omega}, |d\lambda|)$ have Hilbert–Schmidt kernels

$$N_{11}(\lambda, \nu) = \frac{1}{2\pi i} \frac{e^{-\frac{1}{8}\lambda^2}}{\lambda - \nu} \chi_{\mathbb{R}}(\lambda) \chi_{\mathbb{R}+i\delta}(\nu),$$

$$N_{12}(\nu, w) = \frac{1}{2\pi} \frac{e^{-\frac{1}{8}w^2}}{\nu - w} \chi_{\mathbb{R}+i\delta}(\nu) \chi_{\Gamma}(w).$$

Similarly with (3.2) for $N_2(w, \mu)$ in (2.11),

$$N_2(w, \mu) = \frac{1}{4i\pi^2} \int_{\mathbb{R}+i\delta} \frac{e^{-\frac{1}{8}(w^2+\mu^2)+it(w-\mu)}}{(v-\mu)(v-w)} dv \chi_{\Gamma}(w) \chi_{\mathbb{R}}(\mu),$$

and thus $N_2 = N_{21}N_{22}$ with $N_{2j} : L^2(\mathring{\Omega}, |d\lambda|) \rightarrow L^2(\mathring{\Omega}, |d\lambda|)$ once more Hilbert–Schmidt,

$$N_{21}(w, \nu) = \frac{1}{2\pi i} \frac{e^{-\frac{1}{8}w^2+itw}}{\nu - w} \chi_{\Gamma}(w) \chi_{\mathbb{R}+i\delta}(\nu),$$

$$N_{22}(\nu, \mu) = \frac{1}{2\pi} \frac{e^{-\frac{1}{8}\mu^2-it\mu}}{\nu - \mu} \chi_{\mathbb{R}+i\delta}(\nu) \chi_{\mathbb{R}}(\mu).$$

Hence, N_1 and N_2 are trace-class on $L^2(\mathring{\Omega}, |d\lambda|)$ as claimed. \square

The last lemma allows us to compute for $j = 1, 2$,

$$\text{tr}_{L^2(\mathring{\Omega})} N_j = \int_{\mathring{\Omega}} N_j(\lambda, \lambda) d\lambda = 0,$$

that is, N_j are traceless and even more, they are nilpotent on $L^2(\mathring{\Omega}, |d\lambda|)$ with $N_j^2 = 0$ (simply recall that \mathbb{R} is disjoint from Γ). The last observation lies at the heart of the following useful operator identity.

LEMMA 3.2. *We have on $L^2(\mathring{\Omega}, |d\lambda|)$,*

$$(3.3) \quad (1 - \sqrt{\gamma}N_1)^{-1}(1 - \sqrt{\gamma}(N_1 + N_2))(1 - \sqrt{\gamma}N_2)^{-1} = 1 - \gamma N_1 N_2.$$

PROOF. By nilpotency of N_j , we know that $1 - N_j$ is invertible on $L^2(\mathring{\Omega}, |d\lambda|)$, in fact

$$(1 - \sqrt{\gamma}N_j \upharpoonright_{L^2(\mathring{\Omega})})^{-1} = 1 + \sqrt{\gamma}N_j \upharpoonright_{L^2(\mathring{\Omega})}, \quad j = 1, 2.$$

Now substitute the latter into the left-hand side of (3.3), multiply out and simplify using $N_j^2 = 0$. The desired equality follows at once. \square

Since all factors in (3.3) are of the form identity plus trace-class (by Lemma 3.1 and the triangle inequality for the trace norm) we are allowed to use the multiplicative nature of the Fredholm determinant [24], Chapter II, (5.1), and conclude from (3.3),

$$(3.4) \quad \det(1 + \sqrt{\gamma}N_1 \upharpoonright_{L^2(\mathring{\Omega})}) \det(1 - \sqrt{\gamma}(N_1 + N_2) \upharpoonright_{L^2(\mathring{\Omega})})$$

$$\times \det(1 + \sqrt{\gamma}N_2 \upharpoonright_{L^2(\mathring{\Omega})})$$

$$= \det(1 - \gamma N_1 N_2 \upharpoonright_{L^2(\mathring{\Omega})}).$$

But from the Plemelj–Smithies formula [24], Chapter II, Theorem 3.1, and our previous comments about vanishing traces and nilpotency of N_j ,

$$(3.5) \quad \det(1 - \sqrt{\gamma}N_j \upharpoonright_{L^2(\mathring{\Omega})}) = \exp\left[-\sum_{k=1}^{\infty} \frac{\gamma^{\frac{k}{2}}}{k} \operatorname{tr}_{L^2(\mathring{\Omega})} N_j^k\right] = 1, \quad j = 1, 2,$$

that is, we have just established the following Fredholm determinant equality.

PROPOSITION 3.3. *We have*

$$(3.6) \quad \det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) = \det(1 - G \upharpoonright_{L^2(\mathring{\Omega})}),$$

where $G : L^2(\mathring{\Omega}, |d\lambda|) \rightarrow L^2(\mathring{\Omega}, |d\lambda|)$ is trace class with kernel

$$G(\lambda, \mu; \gamma) = \sqrt{\gamma}(N_1(\lambda, \mu) + N_2(\lambda, \mu)) = \frac{\mathbf{f}^\top(\lambda)\mathbf{g}(\mu)}{\lambda - \mu}, \quad (\lambda, \mu) \in \mathring{\Omega} \times \mathring{\Omega},$$

and

$$(3.7) \quad \mathbf{f}(\lambda) = \sqrt{\frac{\gamma}{2\pi}} e^{-\frac{1}{8}\lambda^2} \begin{bmatrix} \chi_{\mathbb{R}}(\lambda) \\ e^{i\lambda} \chi_{\Gamma}(\lambda) \end{bmatrix}, \quad \mathbf{g}(\mu) = \frac{e^{-\frac{1}{8}\mu^2}}{\sqrt{2\pi}} \begin{bmatrix} \chi_{\Gamma}(\mu) \\ e^{-i\mu} \chi_{\mathbb{R}}(\mu) \end{bmatrix}.$$

PROOF. Use (2.12), (3.4) and (3.5). The formulæ (3.7) for the kernel of $G = \sqrt{\gamma}(N_1 + N_2)$ are read off from (2.11) and this completes our proof. \square

3.2. *Riemann–Hilbert problem.* The following RHP is our starting point for the derivation of the ZS system (1.13) and in turn (1.9). This problem is naturally associated with the integrable operator G in (3.6) by classical theory; cf. [25].

RIEMANN–HILBERT PROBLEM 3.4 (Its, Izergin, Korepin, Slavnov [25] (1990)). *For any $t \in \mathbb{R}, \gamma \in [0, 1]$, determine $\mathbf{Y}(z) = \mathbf{Y}(z; t, \gamma) \in \mathbb{C}^{2 \times 2}$ such that:*

(1) $\mathbf{Y}(z)$ is analytic for $z \in \mathbb{C} \setminus \Omega$ and we orient $\Omega = \mathbb{R} \sqcup \Gamma$ from “left to right” as shown in Figure 8 below.

(2) The boundary values $\mathbf{Y}_{\pm}(z)$ from the left/right side of the oriented contour Ω exist and are related by the jump condition

$$\mathbf{Y}_+(z) = \mathbf{Y}_-(z) \begin{bmatrix} 1 & -i\sqrt{\gamma}e^{-\frac{1}{4}z^2 - iz} \chi_{\mathbb{R}}(z) \\ -i\sqrt{\gamma}e^{-\frac{1}{4}z^2 + iz} \chi_{\Gamma}(z) & 1 \end{bmatrix}, \quad z \in \Omega.$$

(3) As $z \rightarrow \infty$, we have

$$\mathbf{Y}(z) = \mathbb{I} + \mathbf{Y}_1 z^{-1} + \mathcal{O}(z^{-2}), \quad \mathbf{Y}_1 = \mathbf{Y}_1(t, \gamma) = [Y_1^{jk}(t, \gamma)]_{j,k=1}^2.$$




FIG. 8. *The oriented jump contour $\Omega = \mathbb{R} \sqcup \Gamma$ in RHP 3.4 with an admissible choice for Γ in the upper half-plane.*

The general theory of [25] (see also [2], Section 5.6) asserts that for an integrable integral operator (such as our G on $L^2(\mathring{\Omega}, |\mathrm{d}\lambda|)$) its resolvent $R = (1 - G)^{-1} - 1$, if existent, is again of integrable type with kernel

$$(3.8) \quad R(\lambda, \mu; \gamma) = \frac{\mathbf{F}^\top(\lambda)\mathbf{G}(\mu)}{\lambda - \mu}, \quad \mathbf{F}(\lambda) = ((1 - G \upharpoonright_{L^2(\mathring{\Omega})})^{-1}\mathbf{f})(\lambda),$$

$$\mathbf{G}(\mu) = ((1 - G^* \upharpoonright_{L^2(\mathring{\Omega})})\mathbf{g})(\mu).$$

Most importantly, R can be computed in terms of the solution to RHP 3.4,

$$(3.9) \quad \mathbf{F}(z) = \mathbf{Y}_\pm(z)\mathbf{f}(z), \quad \mathbf{G}(z) = (\mathbf{Y}^\top_\pm(z))^{-1}\mathbf{g}(z), \quad z \in \Omega,$$

where the choice of limiting values (\pm) is immaterial. Thus, RHP 3.4 is (uniquely) solvable if and only if $1 - G$ is invertible on $L^2(\mathring{\Omega})$. Furthermore, the solution $\mathbf{Y}(z)$ to RHP 3.4 takes the form

$$(3.10) \quad \mathbf{Y}(z) = \mathbb{I} - \int_\Omega \mathbf{F}(\lambda)\mathbf{g}^\top(\lambda) \frac{\mathrm{d}\lambda}{\lambda - z}, \quad z \notin \Omega.$$

We now prove solvability of RHP 3.4 using two different arguments:

i. *Argument 1:* By Proposition 3.3,

$$(3.11) \quad \det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) = \det(1 - G \upharpoonright_{L^2(\mathring{\Omega})}), \quad t \in \mathbb{R}, \gamma \in [0, 1].$$

The left-hand side in (3.11) is nonvanishing by Lemma 2.1 and since G is trace-class this implies that $1 - G$ is invertible on $L^2(\mathring{\Omega}, |\mathrm{d}\lambda|)$ for all $t \in \mathbb{R}, \gamma \in [0, 1]$; cf. [24], Theorem 6.1. Thus RHP 3.4 is solvable for all $t \in \mathbb{R}, \gamma \in [0, 1]$; cf. [25].

ii. *Argument 2:* We provide a solvability proof for RHP 3.4 based on a vanishing lemma argument (a standard technique in Riemann–Hilbert analysis; cf. [18, 19, 35]).

LEMMA 3.5 (Vanishing lemma). *Let $t \in \mathbb{R}, \gamma \in [0, 1]$ and suppose \mathbf{Y} satisfies conditions (1) and (2) in RHP 3.4 above but instead of condition (3) we enforce*

$$\mathbf{Y}(z) = \mathcal{O}(z^{-1}), \quad z \rightarrow \infty.$$

Then $\mathbf{Y} \equiv 0$.

PROOF. Let Δ denote the open region in between \mathbb{R} and Γ . Introduce the auxiliary function

$$\mathbf{N}(z) := \mathbf{Y}(z) \begin{cases} \begin{bmatrix} 1 & 0 \\ -i\sqrt{\gamma}e^{-\frac{1}{4}z^2+itz} & 1 \end{bmatrix} & z \in \Delta, \\ \mathbb{I} & z \in \mathbb{C} \setminus \overline{\Delta}, \end{cases}$$

and note that $\mathbf{N}(z)$ also satisfies $\mathbf{N}(z) = \mathcal{O}(z^{-1}), z \rightarrow \infty$. However, $\mathbf{N}(z)$ is jump-free on Γ , instead we have collapsed the jumps to the real line,

$$(3.12) \quad \mathbf{N}_+(z) = \mathbf{N}_-(z) \begin{bmatrix} 1 - \gamma e^{-\frac{1}{2}z^2} & -i\sqrt{\gamma}e^{-\frac{1}{4}z^2-itiz} \\ -i\sqrt{\gamma}e^{-\frac{1}{4}z^2+itz} & 1 \end{bmatrix}, \quad z \in \mathbb{R}.$$

Next, define $\mathbf{H}(z) = \mathbf{N}(z)\mathbf{N}^\dagger(\bar{z})$ for $z \in \mathbb{C} \setminus \mathbb{R}$ where \mathbf{N}^\dagger is the Hermitian conjugate of \mathbf{N} . Since \mathbf{H} is analytic in the upper z -plane, continuous down to the real line and decays of

$\mathcal{O}(z^{-2})$ as $z \rightarrow \infty$, we find from Cauchy’s theorem $\int_{\mathbb{R}} \mathbf{H}_+(z) dz = 0$. We add to this equation its Hermitian conjugate,

$$\begin{aligned}
 (3.13) \quad 0 &= \int_{-\infty}^{\infty} \{ \mathbf{N}_+(z) \mathbf{N}_-^\dagger(z) + \mathbf{N}_-(z) \mathbf{N}_+^\dagger(z) \} dz \\
 &\stackrel{(3.12)}{=} 2 \int_{-\infty}^{\infty} \{ \mathbf{N}_-(z) \begin{bmatrix} 1 - \gamma e^{-\frac{1}{2}z^2} & 0 \\ 0 & 1 \end{bmatrix} \mathbf{N}_-^\dagger(z) \} dz.
 \end{aligned}$$

Reading off the diagonal entries in (3.13) we find in turn

$$\begin{aligned}
 &\int_{-\infty}^{\infty} \{ |N_-^{21}(z)|^2 (1 - \gamma e^{-\frac{1}{2}z^2}) + |N_-^{22}(z)|^2 \} dz \\
 &= 0 = \int_{-\infty}^{\infty} \{ |N_-^{11}(z)|^2 (1 - \gamma e^{-\frac{1}{2}z^2}) + |N_-^{12}(z)|^2 \} dz,
 \end{aligned}$$

so that $\mathbf{N}_-(z) \equiv 0$ by continuity of $\mathbf{N}_-(z)$ on \mathbb{R} , and thus with (3.12) also $\mathbf{N}_+(z) \equiv 0$. In summary, $\mathbf{N}(z)$ is analytic for $\Im z > 0$, continuous for $\Im z \geq 0$ and we have

$$|\mathbf{N}(z)| \leq C, \quad \Im z > 0, \quad \sup_{z \in \mathbb{R}} |\mathbf{N}_+(z)| = 0.$$

Hence, by Carlson’s theorem (cf. [29], Theorem 5.1.2 and Corollary 5.1.3), $\mathbf{N}(z) \equiv 0$ for $\Im z \geq 0$. Dealing with $\Im z \leq 0$ in a similar fashion, we establish triviality of $\mathbf{N}(z)$ in the whole z -plane, and thus also $\mathbf{Y}(z) \equiv 0$. \square

COROLLARY 3.6. *The Riemann–Hilbert problem 3.4 for $\mathbf{Y}(z; t, \gamma)$ has a unique solution for every $t \in \mathbb{R}, \gamma \in [0, 1]$. Moreover, the coefficient $\mathbf{Y}_1(t, \gamma)$ is continuous in $t \in \mathbb{R}$ for every $\gamma \in [0, 1]$.*

PROOF. The RHP 3.4 is equivalent to a singular integral equation (cf. [25]), which can be stated using a Fredholm operator of index zero. The above vanishing lemma then states that the kernel of this operator is trivial, that is, the operator itself is onto. Thus the singular integral equation, equivalently the RHP, is solvable. We refer the interested reader to [35] for more on this subject. Once solvability is established, uniqueness follows from a standard Liouville argument: by RHP 3.4, the scalar function $\det \mathbf{Y}(z)$ is entire, approaching unity at infinity, that is, for any solution of the RHP we have $\det \mathbf{Y}(z) \equiv 1$. Thus, given two solutions $\mathbf{Y}_1(z)$ and $\mathbf{Y}_2(z)$ to RHP 3.4, we consider $\mathbf{Q}(z) = \mathbf{Y}_1(z) \mathbf{Y}_2(z)^{-1}$ which is again entire. Since in addition $\mathbf{Q}(z) \rightarrow \mathbb{I}$ as $z \rightarrow \infty$, equality of \mathbf{Y}_1 and \mathbf{Y}_2 follows from another application of Liouville’s theorem. Finally, continuity of $\mathbf{Y}_1(\cdot, \gamma)$ for fixed $\gamma \in [0, 1]$ follows from continuity of the jump matrix, the fact that RHP 3.4 is solvable for all $(t, \gamma) \in \mathbb{R} \times [0, 1]$ and a standard small norm argument. \square

3.3. The ZS-system. We are now prepared to take the first step in the proof of Theorem 1.6, namely the derivation of a closed-form expression for $\det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})$. First, we state a standard connection formula between the same determinant and RHP 3.4.

PROPOSITION 3.7. *For any fixed $t \in \mathbb{R}, \gamma \in [0, 1]$,*

$$(3.14) \quad \frac{\partial}{\partial t} \ln \det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) = -i Y_1^{22}(t, \gamma),$$

in terms of the matrix $\mathbf{Y}_1(t, \gamma) = [Y_1^{jk}(t, \gamma)]_{j,k=1}^2$ defined in condition (3) of RHP 3.4.

PROOF. Through (3.11) and the Jacobi variation formula,

$$\begin{aligned}
 \frac{\partial}{\partial t} \ln \det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) &\stackrel{(3.11)}{=} \frac{\partial}{\partial t} \ln \det(1 - G \upharpoonright_{L^2(\mathring{\Omega})}) \\
 (3.15) \qquad \qquad \qquad &= - \operatorname{tr}_{L^2(\mathring{\Omega})} \left((1 - G)^{-1} \frac{\partial G}{\partial t} \right) \\
 &= - \int_{\mathring{\Omega}} \int_{\mathring{\Omega}} (1 - G)^{-1}(\lambda, \mu) \frac{\partial G}{\partial t}(\mu, \lambda) \, d\mu \, d\lambda.
 \end{aligned}$$

But from (3.7) we can compute explicitly the t -derivative,

$$\frac{\partial G}{\partial t}(\mu, \lambda) = i f_2(\mu) g_2(\lambda), \qquad \mathbf{f}(\lambda) = \begin{bmatrix} f_1(\lambda) \\ f_2(\lambda) \end{bmatrix}, \qquad \mathbf{g}(\mu) = \begin{bmatrix} g_1(\mu) \\ g_2(\mu) \end{bmatrix},$$

which gives back in (3.15),

$$\begin{aligned}
 \frac{\partial}{\partial t} \ln \det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) &= -i \int_{\mathring{\Omega}} ((1 - G \upharpoonright_{L^2(\mathring{\Omega})})^{-1} f_2)(\lambda) g_2(\lambda) \, d\lambda \\
 (3.16) \qquad \qquad \qquad &\stackrel{(3.8)}{=} -i \int_{\mathring{\Omega}} F_2(\lambda) g_2(\lambda) \, d\lambda.
 \end{aligned}$$

Now use (3.10) and take the limit $z \rightarrow \infty, z \notin \Omega$,

$$\mathbf{Y}(z) = \mathbb{I} + \frac{1}{z} \int_{\Omega} \mathbf{F}(\lambda) \mathbf{g}^T(\lambda) \, d\lambda + \mathcal{O}(z^{-2}),$$

so that by comparison with RHP 3.4, condition (3),

$$(3.17) \qquad \mathbf{Y}_1 = \int_{\Omega} \mathbf{F}(\lambda) \mathbf{g}^T(\lambda) \, d\lambda = \int_{\mathring{\Omega}} \mathbf{F}(\lambda) \mathbf{g}^T(\lambda) \, d\lambda.$$

Identity (3.17) used in the right-hand side of (3.16) implies (3.14) and completes our proof. \square

Second, we derive the ZS-system (1.13) from RHP 3.4 as follows: Define (compare the proof of Lemma 3.5)

$$(3.18) \qquad \mathbf{N}(z) := \mathbf{Y}(z) \begin{cases} \begin{bmatrix} 1 & 0 \\ -i\sqrt{\gamma} e^{-\frac{1}{4}z^2 + itz} & 1 \end{bmatrix} & z \in \Delta, \\ \mathbb{I} & z \in \mathbb{C} \setminus \overline{\Delta}, \end{cases}$$

where Δ lies in between \mathbb{R} and Γ . As noted before, $\mathbf{N}(z)$ solves the problem summarized below.

RIEMANN–HILBERT PROBLEM 3.8. *For any $t \in \mathbb{R}, \gamma \in [0, 1]$, the matrix-valued function $\mathbf{N}(z) = \mathbf{N}(z; t, \gamma) \in \mathbb{C}^{2 \times 2}$ has the following properties:*

(1) $\mathbf{N}(z)$ is analytic for $z \in \mathbb{C} \setminus \mathbb{R}$ and has a continuous extension on the closed upper and lower half-planes.

(2) The square integrable boundary values $\mathbf{N}_{\pm}(z) = \lim_{\epsilon \downarrow 0} \mathbf{N}(z \pm i\epsilon), z \in \mathbb{R}$ obey the jump condition

$$(3.19) \qquad \mathbf{N}_+(z) = \mathbf{N}_-(z) \begin{bmatrix} 1 - \gamma e^{-\frac{1}{2}z^2} & -i\sqrt{\gamma} e^{-\frac{1}{4}z^2 - itz} \\ -i\sqrt{\gamma} e^{-\frac{1}{4}z^2 + itz} & 1 \end{bmatrix}, \qquad z \in \mathbb{R}.$$

(3) As $z \rightarrow \infty$, the leading order behavior of $\mathbf{N}(z)$ remains unchanged from condition (3) in RHP 3.4,

$$\mathbf{N}(z) = \mathbb{I} + \mathbf{Y}_1(t, \gamma)z^{-1} + \mathcal{O}(z^{-2}).$$

A simple check between RHP 3.8 and RHP 1.5 for $\mathbf{X}(z; x, \gamma)$ reveals their equality subject to the identifications

$$(3.20) \quad \mathbf{N}(z; t, \gamma) = \mathbf{X}\left(z; \frac{t}{2}, \gamma\right),$$

$$z \in \mathbb{C} \setminus \mathbb{R}, t \in \mathbb{R}, \gamma \in [0, 1], \text{ with } r(z) = r(z; \gamma) = -i\sqrt{\gamma}e^{-\frac{1}{4}z^2}.$$

For this reason, we now establish solvability of RHP 1.5, and thus, in turn, existence of $y(x; \gamma)$:

THEOREM 3.9. *The RHP 1.5 for $\mathbf{X}(z; x, \gamma)$ with $r(z; \gamma) = -i\sqrt{\gamma}e^{-\frac{1}{4}z^2}$ is uniquely solvable for every $(x, \gamma) \in \mathbb{R} \times [0, 1]$.*

PROOF. Observing the similarities between (3.12) and (1.8), one first derives a vanishing lemma in the style of Lemma 3.5, that is, assumes \mathbf{X} satisfies conditions (1) and (2) in RHP 1.5 but instead $\mathbf{X}(z) = \mathcal{O}(z^{-1}), z \rightarrow \infty$. Now define $\mathbf{H}(z) = \mathbf{X}(z)\mathbf{X}^\dagger(\bar{z}), z \in \mathbb{C} \setminus \mathbb{R}$ and conclude $\int_{\mathbb{R}} \mathbf{H}_+(z) dz = 0$ so that, analogous to (3.13),

$$0 = 2 \int_{-\infty}^{\infty} \left\{ \mathbf{X}_-(z) \begin{bmatrix} 1 - |r(z; \gamma)|^2 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{X}_-^\dagger(z) \right\} dz.$$

This equation allows us to deduce $\mathbf{X}_-(z) \equiv 0$, and thus also $\mathbf{X}_+(z) \equiv 0$. By Carlson’s theorem, we then find $\mathbf{X}(z) \equiv 0$ in the whole z -plane and the vanishing lemma is proven. After that, the proof argument of Corollary 3.6 applies verbatim to $\mathbf{X}(z; x, \gamma)$ and Theorem 3.9 follows. \square

Next, a short remark about (obvious) symmetries in RHP 1.5; see also (1.12).

PROPOSITION 3.10. *Besides (3.20) we also have $\mathbf{Y}_1(t, \gamma) = \mathbf{X}_1(\frac{t}{2}, \gamma), (t, \gamma) \in \mathbb{R} \times [0, 1]$ and for any $(x, \gamma) \in \mathbb{R} \times [0, 1]$,*

$$(3.21) \quad \mathbf{X}(z; x, \gamma) = \sigma_1 \overline{\mathbf{X}(\bar{z}; x, \gamma)} \sigma_1, \quad z \in \mathbb{C} \setminus \mathbb{R}; \sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix},$$

from which we learn that

$$(3.22) \quad X_i^{11}(x, \gamma) = \overline{X_i^{22}(x, \gamma)}, \quad X_i^{21}(x, \gamma) = \overline{X_i^{12}(x, \gamma)}, \quad i \in \mathbb{Z}_{\geq 1}.$$

Furthermore, for any $(x, \gamma) \in \mathbb{R} \times [0, 1]$,

$$(3.23) \quad \mathbf{X}(z; x, \gamma) = \sigma_2 \mathbf{X}(-z; x, \gamma) \sigma_2, \quad z \in \mathbb{C} \setminus \mathbb{R}; \sigma_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix},$$

which leads to

$$(3.24) \quad X_i^{11}(x, \gamma) = (-1)^i X_i^{22}(x, \gamma),$$

$$X_i^{21}(x, \gamma) = (-1)^{i+1} X_i^{12}(x, \gamma), \quad i \in \mathbb{Z}_{\geq 1}.$$

PROOF. The connection between \mathbf{Y}_1 and \mathbf{X}_1 follows from (3.20). For (3.21), respectively (3.23), use unique solvability of RHP 1.5 as $\sigma_1 \overline{\mathbf{X}}(\bar{z}; x, \gamma) \sigma_1$, respectively, $\sigma_2 \mathbf{X}(-z; x, \gamma) \sigma_2$, solve the same problem. \square

And finally, the following straightforward and standard steps, compare Section 1.2 above: As we already know,

$$\Psi(z) := \mathbf{X}(z)e^{-ixz\sigma_3}, \quad z \in \mathbb{C} \setminus \mathbb{R},$$

solves (1.11). However, by definition,

$$y = y(x; \gamma) := 2iX_1^{12}(x, \gamma) \Rightarrow \bar{y}(x; \gamma) \stackrel{(3.22)}{=} -2iX_1^{21}(x, \gamma).$$

Moreover, expanding $\frac{\partial \Psi}{\partial x} \Psi^{-1}$ up to $\mathcal{O}(z^{-2})$ as $z \rightarrow \infty$, we obtain from comparison with (1.11),

$$\frac{\partial \mathbf{X}_1}{\partial x} = -2i \begin{bmatrix} -X_1^{12} X_1^{21} & X_2^{12} - X_1^{12} X_1^{22} \\ -X_2^{21} + X_1^{21} X_1^{11} & X_1^{21} X_1^{12} \end{bmatrix},$$

so in the (22)-entry,

$$(3.25) \quad \frac{\partial X_1^{22}}{\partial x} = -\frac{i}{2}|y|^2.$$

We summarize by combining (3.14), Proposition 3.10 and (3.25),

$$\begin{aligned} \frac{\partial^2}{\partial t^2} \ln \det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) &\stackrel{(3.14)}{=} -i \frac{\partial}{\partial t} Y_1^{22}(t, \gamma) = -\frac{i}{2} \frac{\partial}{\partial x'} X_1^{22}(x', \gamma) \Big|_{x'=\frac{t}{2}} \\ &\stackrel{(3.25)}{=} -\frac{1}{4} \left| y\left(\frac{t}{2}; \gamma\right) \right|^2, \end{aligned}$$

so that after integration

$$(3.26) \quad \begin{aligned} &\ln \det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) \\ &= -\frac{1}{4} \int_t^\infty (x - t) \left| y\left(\frac{x}{2}; \gamma\right) \right|^2 dx + c_1(\gamma)t + c_2(\gamma) \end{aligned}$$

for some t -independent c_i . The fastest way to show $c_i = 0$ follows from a comparison of the $t \rightarrow +\infty$ asymptotic expansion in the left- and right-hand side of (3.26).

3.4. *Right tail asymptotics I.* Begin with

$$\det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) = \exp \left[- \sum_{k=1}^\infty \frac{\gamma^k}{k} \operatorname{tr}_{L^2(\mathbb{R})} (T \chi_t)^k \right],$$

and use that, as $t \rightarrow +\infty$,

$$(3.27) \quad \begin{aligned} \operatorname{tr}_{L^2(\mathbb{R})} (T \chi_t) &= \int_t^\infty T(x, x) dx = \frac{1}{2\pi} \sqrt{\frac{\pi}{2}} \int_t^\infty \operatorname{erfc}(\sqrt{2}x) dx \\ &= \frac{1}{\sqrt{2\pi}} \frac{\operatorname{erfc}(\sqrt{2}t)}{8t} (1 + \mathcal{O}(t^{-2})), \end{aligned}$$

together with

$$k \in \mathbb{Z}_{\geq 2}: \quad \left| \operatorname{tr}_{L^2(\mathbb{R})} (T \chi_t)^k \right| \leq C (\operatorname{erfc}(\sqrt{2}t))^k, \quad C > 0, t \rightarrow +\infty.$$

In summary,

LEMMA 3.11. As $t \rightarrow +\infty$,

$$\det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) = 1 - \frac{\gamma}{\sqrt{2\pi}} \frac{\operatorname{erfc}(\sqrt{2}t)}{8t} (1 + \mathcal{O}(t^{-2})).$$

On the other hand, we will now derive the large t -expansion for the right-hand side of (3.26) from a Deift–Zhou nonlinear steepest descent analysis [12] of RHP 3.8. The results will in turn verify (1.10), and thus, after integration in (3.26) and comparison with Lemma 3.11, show that $c_i = 0$. The detailed steps of the nonlinear steepest descent analysis are standard: assume $t > 0$ throughout and first rescale while simultaneously opening lenses,

$$(3.28) \quad \mathbf{T}(z; t, \gamma) := \mathbf{N}(zt; t, \gamma) \begin{cases} \begin{bmatrix} 1 & 0 \\ i\sqrt{\gamma}e^{-t^2\theta_+(z)} & 1 \end{bmatrix} & \Im z \in (0, \delta), \\ \begin{bmatrix} 1 & -i\sqrt{\gamma}e^{-t^2\theta_-(z)} \\ 0 & 1 \end{bmatrix} & \Im z \in (-\delta, 0), \\ \mathbb{I} & \text{else,} \end{cases}$$

with $\delta > 0$ fixed.

We abbreviate $\theta_{\pm}(z) := \frac{1}{4}(z^2 \mp 4iz)$ and keeping the sign charts of $\theta_{\pm}(z)$ in mind (see Figure 9 below) the function $\mathbf{T}(z)$ solves the following RHP.

RIEMANN–HILBERT PROBLEM 3.12. For any $t \in \mathbb{R}_{>0}$, $\gamma \in [0, 1]$ the transformed function $\mathbf{T}(z) = \mathbf{T}(z; t, \gamma) \in \mathbb{C}^{2 \times 2}$ defined in (3.28) has the following properties.

(1) $\mathbf{T}(z)$ is analytic for $z \in \mathbb{C} \setminus \{\Im z = \pm\delta\}$ and both straight lines are oriented from left to right; see Figure 10 below.

(2) Along the two straight lines,

$$\mathbf{T}_+(z) = \mathbf{T}_-(z) \begin{bmatrix} 1 & 0 \\ -i\sqrt{\gamma}e^{-t^2\theta_+(z)} & 1 \end{bmatrix}, \quad \Im z = \delta,$$

and

$$\mathbf{T}_+(z) = \mathbf{T}_-(z) \begin{bmatrix} 1 & -i\sqrt{\gamma}e^{-t^2\theta_-(z)} \\ 0 & 1 \end{bmatrix}, \quad \Im z = -\delta.$$




FIG. 9. The sign charts of $\theta_+(z)$ on the left and $\theta_-(z)$ on the right. We indicate in red values $z \in \mathbb{C}$ where $\Re\theta_{\pm}(z) < 0$ and in blue where $\Re\theta_{\pm}(z) > 0$. Also $\Im\theta_{\pm}(z) = 0$ is highlighted along the dash–dotted black straight lines.




FIG. 10. The oriented jump contour $\Im z = \pm\delta$ in RHP 3.12.

(3) As $z \rightarrow \infty$, the leading order behavior in RHP 3.8 remains formally unchanged,

$$\mathbf{T}(z) = \mathbb{I} + \mathbf{T}_1(t, \gamma)z^{-1} + \mathcal{O}(z^{-2}), \quad \mathbf{T}_1(t, \gamma) = \mathbf{Y}_1(t, \gamma)t^{-1}.$$

The stationary points of the exponents $\theta_{\pm}(z)$ are $z_{\pm} = \pm 2i$ and we denote by γ_{\pm} the steepest descent contours passing through z_{\pm} along which $\Im\theta_{\pm}(z) = 0$. Explicitly,

$$\gamma_+ : \Re z \in \mathbb{R}, \Im z = 2; \quad \gamma_- : \Re z \in \mathbb{R}, \Im z = -2.$$

Note that both contours γ_{\pm} lie in the domain where $\Re\theta_{\pm}(z) > 0$ (see Figure 9) so we are allowed to choose $\delta = 2$ in RHP 3.12 to match those steepest descent contours. After that, applying standard arguments of the classical Laplace method, we derive the following estimates for the jump matrix $\mathbf{G}_{\mathbf{T}}(z; t, \gamma)$ in condition (2) of RHP 3.12.

PROPOSITION 3.13. *There exist positive constants t_0 and c such that*

$$\begin{aligned} \|\mathbf{G}_{\mathbf{T}}(\cdot; t, \gamma) - \mathbb{I}\|_{L^\infty(\Im z = \pm 2, |dz|)} &\leq c\sqrt{\gamma}e^{-t^2}, \\ \|\mathbf{G}_{\mathbf{T}}(\cdot; t, \gamma) - \mathbb{I}\|_{L^2(\Im z = \pm 2, |dz|)} &\leq c\sqrt{\gamma}t^{-\frac{1}{2}}e^{-t^2}, \end{aligned}$$

hold true for all $t \geq t_0$ and $\gamma \in [0, 1]$.

These estimates show by general theory [12] that RHP 3.12 is uniquely solvable in $L^2(\Im z = \pm 2, |dz|)$ for $t \geq t_0, \gamma \in [0, 1]$ and its solution can be computed from the integral equation

$$(3.29) \quad \mathbf{T}(z) = \mathbb{I} + \frac{1}{2\pi i} \int_{\Im \lambda = \pm 2} \mathbf{T}_-(\lambda)(\mathbf{G}_{\mathbf{T}}(\lambda) - \mathbb{I}) \frac{d\lambda}{\lambda - z}, \quad \Im z \neq \pm 2,$$

using that

$$(3.30) \quad \|\mathbf{T}(\cdot; t, \gamma) - \mathbb{I}\|_{L^2(\Im z = \pm 2, |dz|)} \leq c\sqrt{\gamma}t^{-\frac{1}{2}}e^{-t^2} \quad \forall t \geq t_0, \gamma \in [0, 1].$$

In particular, see condition (3) in RHP 3.12 and (3.29),

$$\mathbf{Y}_1(t, \gamma) = t\mathbf{T}_1(t, \gamma) = \frac{it}{2\pi} \int_{\Im \lambda = \pm 2} \mathbf{T}_-(\lambda)(\mathbf{G}_{\mathbf{T}}(\lambda) - \mathbb{I}) d\lambda,$$

so that with (3.30) and Proposition 3.13, as $t \rightarrow +\infty$,

$$(3.31) \quad \begin{aligned} \mathbf{Y}_1(t, \gamma) &= \frac{it}{2\pi} \int_{\Im \lambda = \pm 2} (\mathbf{G}_{\mathbf{T}}(\lambda) - \mathbb{I}) d\lambda + \mathcal{O}(\gamma e^{-2t^2}) \\ &= \frac{\sigma_1 \sqrt{\gamma}}{\sqrt{\pi}} e^{-t^2} + \mathcal{O}(\gamma e^{-2t^2}). \end{aligned}$$

By Proposition 3.10 and the identity $y(x; \gamma) = 2iX_1^{12}(x, \gamma)$ we then find (1.10), namely

$$y(x; \gamma) = 2iX_1^{12}(x, \gamma) = 2iY_1^{12}(2x, \gamma) = 2i\sqrt{\frac{\gamma}{\pi}} e^{-4x^2} + \mathcal{O}(\gamma e^{-8x^2}), \quad x \rightarrow +\infty,$$

and with this back in the right-hand side of (3.26),

$$(3.32) \quad -\frac{1}{4} \int_t^\infty (x-t) \left| y\left(\frac{x}{2}; \gamma\right) \right|^2 dx = \mathcal{O}(\gamma t^{-2} e^{-2t^2}), \quad t \rightarrow +\infty.$$

Hence, comparing Lemma 3.11 to (3.32) in (3.26) we find $c_i = 0$ and have therefore shown

$$(3.33) \quad \det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) = \exp\left[-\frac{1}{4} \int_t^\infty (x-t) \left| y\left(\frac{x}{2}; \gamma\right) \right|^2 dx\right],$$

where $y = y(x; \gamma) : \mathbb{R} \rightarrow i\mathbb{R}$ is related to the inverse scattering problem for (1.13) with the indicated reflection coefficient.² This verifies the first part in (1.9).

4. Proof of Theorem 1.6, part 2. Our goal in this section is to show that the integral

$$(4.1) \quad \Gamma_{t\gamma} = 1 - \gamma \int_t^\infty G(x) ((1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1} g)(x) dx,$$

$$t \in \mathbb{R}; g(x) = \frac{1}{\sqrt{\pi}} e^{-x^2},$$

equals

$$(4.2) \quad \Gamma_{t\gamma} = \cosh\left[-\frac{i}{2} \int_t^\infty y\left(\frac{x}{2}; \gamma\right) dx\right] - \sqrt{\gamma} \sinh\left[-\frac{i}{2} \int_t^\infty y\left(\frac{x}{2}; \gamma\right) dx\right],$$

which, in turn, will complete the proof of Theorem 1.6, identity (1.9), through (1.3). We start by introducing

$$u^\gamma(t) := \gamma \int_t^\infty G(x) ((1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1} g)(x) dx$$

$$= \int_t^\infty Q^\gamma(x; t) \int_{-\infty}^x g^\gamma(v) dv dx, \quad (t, \gamma) \in \mathbb{R} \times [0, 1],$$

where

$$Q^\gamma(x; t) := ((1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1} g^\gamma)(x), \quad g^\gamma(x) := \sqrt{\gamma} g(x), x \in \mathbb{R}.$$

PROPOSITION 4.1. For any $(t, \gamma) \in \mathbb{R} \times [0, 1]$,

$$(4.3) \quad \frac{d}{dt} u^\gamma(t) = -Y_1^{12}(t, \gamma) ((1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1} G^\gamma)(t),$$

$$G^\gamma(t) := \int_{-\infty}^t g^\gamma(y) dy,$$

in terms of Y_1 from RHP 3.4, condition (3).

PROOF. Start by recalling (2.10), the definition of the operator M before Remark 2.5, the definition of the operator N (see Remark 2.5, and Lemma 3.2) identity (3.3),

$$(4.4) \quad 1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}$$

$$= \mathcal{F}_{\text{ext}}(1 - \gamma E_{\text{ext}} \upharpoonright_{L^2(\mathring{\Omega})}) \mathcal{F}_{\text{ext}}^{-1} = \mathcal{F}_{\text{ext}} M (1 - \gamma N \upharpoonright_{L^2(\mathring{\Omega})}) M^{-1} \mathcal{F}_{\text{ext}}^{-1}$$

$$= \mathcal{F}_{\text{ext}} M (1 - \sqrt{\gamma} N_1 \upharpoonright_{L^2(\mathring{\Omega})})^{-1}$$

$$\times (1 - G \upharpoonright_{L^2(\mathring{\Omega})}) (1 - \sqrt{\gamma} N_2 \upharpoonright_{L^2(\mathring{\Omega})})^{-1} M^{-1} \mathcal{F}_{\text{ext}}^{-1}.$$

²Proposition 3.10 together with $y(x; \gamma) = 2iX^{12}(x, \gamma)$ shows that $y(x; \gamma)$ is purely imaginary for $(x, \gamma) \in \mathbb{R} \times [0, 1]$.

With (4.1), we now compute

$$(M^{-1}\mathcal{F}_{\text{ext}}^{-1}g^\gamma)(x) \stackrel{(2.1)}{=} \sqrt{\frac{\gamma}{2\pi}}e^{-\frac{1}{8}x^2}\chi_{\mathbb{R}}(x) \stackrel{(3.7)}{=} f_1(x).$$

Since $f_1 \in L^2(\mathbb{R})$ is supported on the real line only, it thus lies in the kernel of the operator $N_1 : L^2(\mathring{\Omega}, |d\lambda|) \rightarrow L^2(\mathring{\Omega}, |d\lambda|)$, compare (2.11). Hence,

$$\begin{aligned} (4.5) \quad & ((1 - G \upharpoonright_{L^2(\mathring{\Omega})})^{-1}(1 - \sqrt{\gamma}N_1 \upharpoonright_{L^2(\mathring{\Omega})})M^{-1}\mathcal{F}_{\text{ext}}^{-1}g^\gamma)(x) \\ & = ((1 - G \upharpoonright_{L^2(\mathring{\Omega})})^{-1}f_1)(x) \stackrel{(3.8)}{=} F_1(x), \end{aligned}$$

in terms of $F_1(\lambda) = F_1(\lambda; t, \gamma)$ defined in (3.8). But any function in the range of the operator N_2 will be supported on $\Gamma \neq \mathbb{R}$ only, compare again (2.11). Hence we have $\text{Ran}(N_2) \subseteq \text{Ker}(\mathcal{F}_{\text{ext}})$ by Convention 2.3 and, therefore, from (4.4) and (4.5), with $(x, t, \gamma) \in \mathbb{R}^2 \times [0, 1]$,

$$\begin{aligned} (4.6) \quad Q^\gamma(x; t) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-ix\lambda} e^{-\frac{1}{8}\lambda^2} F_1(\lambda) d\lambda \\ &= \frac{1}{\sqrt{2\pi}} \int_{\mathring{\Omega}} e^{-ix\lambda} e^{-\frac{1}{8}\lambda^2} \chi_{\mathbb{R}}(\lambda) F_1(\lambda) d\lambda. \end{aligned}$$

As an important special case, we learn from the last equation that (compare (3.7) and (3.17))

$$(4.7) \quad Q^\gamma(t; t) = Y_1^{12}(t, \gamma), \quad (t, \gamma) \in \mathbb{R} \times [0, 1].$$

At this point, t -differentiate $u^\gamma(t)$, using (4.7),

$$\begin{aligned} (4.8) \quad \frac{d}{dt}u^\gamma(t) &= -Y_{12}(t, \gamma) \int_{-\infty}^t g^\gamma(v) dv \\ &+ \int_t^{\infty} \left[\frac{\partial}{\partial t} Q^\gamma(x; t) \right] \int_{-\infty}^x g^\gamma(v) dv dx \end{aligned}$$

and recall the following basic fact (cf. [32], (2.10), or [21], (9.132))

$$\frac{\partial}{\partial t} Q^\gamma(x; t) = -R^\gamma(x, t)Q^\gamma(t; t) = -R^\gamma(x, t)Y_1^{12}(t, \gamma),$$

where $R^\gamma(x, y)$ is the kernel of the resolvent integral operator $(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1} = 1 + R^\gamma \upharpoonright_{L^2(\mathbb{R})}$. Thus, back in (4.8),

$$\begin{aligned} \frac{d}{dt}u^\gamma(t) &= -Y_{12}(t, \gamma) \int_{-\infty}^t g^\gamma(v) dv - Y_1^{12}(t, \gamma) \int_t^{\infty} R^\gamma(x, t) \int_{-\infty}^x g^\gamma(v) dv dx \\ &= -Y_1^{12}(t, \gamma) \int_t^{\infty} (1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1}(x, t) \int_{-\infty}^x g^\gamma(v) dv dx \\ &= -Y_1^{12}(t, \gamma)((1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1}G^\gamma)(t), \end{aligned}$$

where we used self-adjointness of $T \chi_t$. Identity (4.3) is proven. \square

PROPOSITION 4.2. For any $(t, \gamma) \in \mathbb{R} \times [0, 1]$,

$$(4.9) \quad ((1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1}G^\gamma)(t) = \int_{-\infty}^t ((1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1}g^\gamma)(y) dy.$$

PROOF. Identity (4.9) follows from Proposition B.1 and Corollary B.2 in Appendix B with the operator identifications $K = \gamma T$, that is, $\phi(x) = g^\gamma(x) = \psi(x)$, and the choice of interval $I = (-\infty, 0) \subset \mathbb{R}$. In detail, for $\gamma \in [0, 1)$ we use Corollary B.2 with $x = t$ and the Neumann series representation (recall Lemma 2.1) to deduce

$$(4.10) \quad \begin{aligned} & ((1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1} G^\gamma)(t) \\ &= \int_{-\infty}^0 ((1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1} g^\gamma)(y + t) \, dy, \end{aligned}$$

which is the right-hand side in (4.9) after a shift. For $\gamma = 1$, we take the limit $\gamma \uparrow 1$ in (4.10). □

REMARK 4.3. Identity (4.9) is the γ -generalization of the corresponding equality in [27], (2.6), (2.8), (2.10).

The strategy is now to derive a coupled system of differential equations for the auxiliary function

$$(4.11) \quad \begin{aligned} A^\gamma(t) &:= \int_{-\infty}^t ((1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})})^{-1} g^\gamma)(x) \, dx \\ &\stackrel{(4.6)}{=} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t \left[\int_{\mathring{\Omega}} e^{-ix\lambda} e^{-\frac{1}{8}\lambda^2} \chi_{\mathbb{R}}(\lambda) F_1(\lambda) \, d\lambda \right] dx \end{aligned}$$

and the (closely related) quantity

$$(4.12) \quad \begin{aligned} B^\gamma(t) &:= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t \left[\int_{\mathring{\Omega}} e^{-ix\lambda} e^{-\frac{1}{8}\lambda^2} \chi_{\mathbb{R}}(\lambda) F_2(\lambda) \, d\lambda \right] dx, \\ (t, \gamma) &\in \mathbb{R} \times [0, 1], \end{aligned}$$

where $F_2(\lambda) = F_2(\lambda; t, \gamma)$ is given in (3.8). Imposing boundary conditions, we then compute the unique solutions $(A^\gamma(t), B^\gamma(t))$ and obtain (4.2) through (4.9) and integration in (4.3).

LEMMA 4.4. *The functions $A^\gamma(t)$ and $B^\gamma(t)$ defined in (4.11), (4.12) for $(t, \gamma) \in \mathbb{R} \times [0, 1]$ satisfy the differential equations*

$$(4.13) \quad \frac{dA^\gamma}{dt} = iY_1^{12}(t, \gamma)B^\gamma + Y_1^{12}(t, \gamma), \quad \frac{dB^\gamma}{dt} = -iY_1^{21}(t, \gamma)A^\gamma,$$

with boundary conditions $A^\gamma(t) \rightarrow \sqrt{\gamma}$, $B^\gamma(t) \rightarrow 0$ as $t \rightarrow +\infty$. The quantity $\mathbf{Y}_1(t, \gamma) = [Y_1^{jk}(t, \gamma)]_{j,k=1}^2$ occurred first in RHP 3.4, condition (3).

PROOF. When t -differentiating (4.11) and (4.12), we require the partial derivatives $\frac{\partial F_j}{\partial t}(\lambda; t, \gamma)$ with $\lambda \in \mathbb{R}$. For these use (3.9) with, say, the $(-)$ limiting value in place, and (3.18),

$$(4.14) \quad \mathbf{F}(\lambda) = \mathbf{N}_-(\lambda)\mathbf{f}(\lambda), \quad \lambda \in \mathbb{R}.$$

But $\mathbf{N}(\lambda)e^{-\frac{i}{2}t\lambda\sigma_3}$ obeys the rescaled Zakharov–Shabat system (compare (1.11)),

$$\frac{\partial}{\partial t}(\mathbf{N}(\lambda)e^{-\frac{i}{2}t\lambda\sigma_3}) = \left\{ -\frac{i}{2}\lambda\sigma_3 + i \begin{bmatrix} 0 & Y_1^{12} \\ -Y_1^{21} & 0 \end{bmatrix} \right\} \mathbf{N}(\lambda)e^{-\frac{i}{2}t\lambda\sigma_3},$$

so that in turn from (4.14),

$$\begin{aligned} \frac{\partial \mathbf{F}}{\partial t}(\lambda) &= \left\{ -\frac{i}{2}\lambda\sigma_3 + i \begin{bmatrix} 0 & Y_1^{12} \\ -Y_1^{21} & 0 \end{bmatrix} \right\} \mathbf{F}(\lambda) \\ &\quad + \frac{i}{2}\lambda \mathbf{N}_-(\lambda)\sigma_3 \mathbf{f}(\lambda) + \mathbf{N}_-(\lambda) \frac{\partial \mathbf{f}}{\partial t}(\lambda), \quad \lambda \in \mathbb{R}. \end{aligned}$$

But once we substitute (3.7) and (3.9) into this vector equation we find

$$(4.15) \quad \frac{\partial F_1}{\partial t}(\lambda) = iY_1^{12} F_2(\lambda), \quad \frac{\partial F_2}{\partial t}(\lambda) = i\lambda F_2(\lambda) - iY_1^{21} F_1(\lambda).$$

Now t -differentiate $A^\gamma(t)$ first,

$$\begin{aligned} \frac{dA^\gamma}{dt} &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t \left[\int_{\mathring{\Omega}} e^{-ix\lambda} e^{-\frac{1}{8}\lambda^2} \chi_{\mathbb{R}}(\lambda) \frac{\partial F_1}{\partial t}(\lambda) d\lambda \right] dx \\ &\quad + \frac{1}{\sqrt{2\pi}} \int_{\mathring{\Omega}} e^{-it\lambda} e^{-\frac{1}{8}\lambda^2} \chi_{\mathbb{R}}(\lambda) F_1(\lambda) d\lambda, \end{aligned}$$

use (4.15), (4.12) in the first integral and (3.7), (3.17) in the second, that is,

$$\frac{dA^\gamma}{dt} = iY_1^{12} B^\gamma + Y_1^{12}.$$

Similarly, for $B^\gamma(t)$: differentiate

$$\begin{aligned} \frac{dB^\gamma}{dt} &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t \left[\int_{\mathring{\Omega}} e^{-ix\lambda} e^{-\frac{1}{8}\lambda^2} \chi_{\mathbb{R}}(\lambda) \frac{\partial F_2}{\partial t}(\lambda) d\lambda \right] dx \\ &\quad + \frac{1}{\sqrt{2\pi}} \int_{\mathring{\Omega}} e^{-it\lambda} e^{-\frac{1}{8}\lambda^2} \chi_{\mathbb{R}}(\lambda) F_2(\lambda) d\lambda, \end{aligned}$$

and use (4.15), (4.11) in the first term and (3.7), (3.17) in the second,

$$(4.16) \quad \begin{aligned} \frac{dB^\gamma}{dt} &= -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^t \frac{\partial}{\partial x} \left[\int_{\mathring{\Omega}} e^{-ix\lambda} e^{-\frac{1}{8}\lambda^2} \chi_{\mathbb{R}}(\lambda) F_2(\lambda) d\lambda \right] dx \\ &\quad - iY_1^{21} A^\gamma + Y_1^{22}. \end{aligned}$$

But from the Riemann–Lebesgue lemma (using that $e^{-\frac{1}{8}\lambda^2} \chi_{\mathbb{R}}(\lambda) F_2(\lambda) \in L^1(\mathbb{R})$ by Cauchy–Schwarz)

$$\lim_{|x| \rightarrow \infty} \int_{\mathring{\Omega}} e^{-ix\lambda} e^{-\frac{1}{8}\lambda^2} \chi_{\mathbb{R}}(\lambda) F_2(\lambda) d\lambda = 0,$$

so we can simplify (4.16) further and obtain with (3.7), (3.17) in the end

$$\frac{dB^\gamma}{dt} = -Y_1^{22} - iY_1^{21} A^\gamma + Y_1^{22} = -iY_1^{21} A^\gamma.$$

Since $Y_1^{12} = Y_1^{21}$ (compare Proposition 3.10) is known, the system (4.13) fully determines $(A^\gamma(t), B^\gamma(t))$ once we impose boundary conditions. And for this we can use the asymptotic results derived in Proposition 3.13 and (3.30). In detail, we trace back our steps through (3.9), (3.18) and (3.28),

$$z \in \mathbb{R} : \quad F_1(z) = N_-^{11}(z) f_1(z) = T_-^{11} \left(\frac{z}{t} \right) f_1(z),$$

and use that from (3.29) and (3.30),

$$(4.17) \quad \mathbf{T}_-(z) = \mathbb{I} + \mathcal{O}\left(\frac{e^{-t^2}}{1 + |z|}\right), \quad t \rightarrow +\infty, z \in \mathbb{R}.$$

Thus,

$$A^\gamma(t) = \sqrt{\gamma}G(t)(1 + \mathcal{O}(e^{-t^2})), \quad t \rightarrow +\infty; G(t) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^t e^{-y^2} dy,$$

which shows that $A^\gamma(t) \rightarrow \sqrt{\gamma}$ in the same limit. Quite similarly,

$$z \in \mathbb{R} : \quad F_2(z) = N_-^{21}(z) f_1(z) = T_-^{21}\left(\frac{z}{t}\right) f_1(z),$$

so that with (4.17), $B^\gamma(t) = \mathcal{O}(\sqrt{\gamma}e^{-t^2}) = o(1)$ as $t \rightarrow +\infty$ and this completes our proof. \square

By means of (3.31), we now simply check that the unique solution to the system (4.13) with the imposed boundary conditions is given by

$$(4.18) \quad \begin{aligned} A^\gamma(t) &= \sqrt{\gamma} \cosh\left[\int_t^\infty Y_1^{12}(x, \gamma) dx\right] - \sinh\left[\int_t^\infty Y_1^{12}(x, \gamma) dx\right], \\ (t, \gamma) &\in \mathbb{R} \times [0, 1], \end{aligned}$$

and

$$\begin{aligned} B^\gamma(t) &= i\left(1 - \cosh\left[\int_t^\infty Y_1^{12}(x, \gamma) dx\right] + \sqrt{\gamma} \sinh\left[\int_t^\infty Y_1^{12}(x, \gamma) dx\right]\right), \\ (t, \gamma) &\in \mathbb{R} \times [0, 1]. \end{aligned}$$

Now we combine (4.3), (4.9) and (4.11),

$$\frac{d}{dt} u^\gamma(t) = -Y_1^{12}(t, \gamma) A^\gamma(t),$$

integrate using (4.18) (with the normalization $u^\gamma(t) \rightarrow 0$ as $t \rightarrow \infty$),

$$(4.19) \quad u^\gamma(t) = 1 + \sqrt{\gamma} \sinh\left[\int_t^\infty Y_1^{12}(x, \gamma) dx\right] - \cosh\left[\int_t^\infty Y_1^{12}(x, \gamma) dx\right],$$

and recall that $\Gamma_{t\gamma} = 1 - u^\gamma(t)$. This confirms (4.2), and thus, after combining $\Gamma_{t\gamma}$ with (3.33) also Theorem 1.6.

4.1. *Right tail asymptotics II.* Since we have already established Lemma 3.11, we are now left with (4.2) and its large positive t -expansion. But since with (3.31),

$$Y_1^{12}(t, \gamma) = \sqrt{\frac{\gamma}{\pi}} e^{-t^2} + \mathcal{O}(\gamma e^{-2t^2}), \quad t \rightarrow +\infty, \gamma \in [0, 1],$$

we obtain at once,

$$\int_t^\infty Y_1^{12}(x, \gamma) dx = \frac{\sqrt{\gamma}}{2} \operatorname{erfc}(t) + \mathcal{O}(\gamma t^{-1} e^{-2t^2}),$$

and thus in turn with (4.19) and the relation $\Gamma_{t\gamma} = 1 - u^\gamma(t)$,

LEMMA 4.5. As $t \rightarrow +\infty$,

$$\Gamma_{t\gamma} = 1 - \frac{\gamma}{2} \operatorname{erfc}(t) + \mathcal{O}(\gamma^{\frac{3}{2}} t^{-1} e^{-2t^2}).$$

The combination of Lemma 3.11 and 4.5 with the known asymptotic behavior of $\operatorname{erfc}(z)$ (cf. [26], 7.12.1) proves (1.15) once substituted into (1.7).

5. Left tail asymptotics—proof of Corollary 1.10, expansion (1.16). Our goal in this section is to prove expansion (1.16) for values $\gamma \in (0, 1)^3$ that are either fixed or approach $\gamma = 1$ at a controlled rate. These goals will be achieved by deriving the analogue of (3.31) for $t \rightarrow -\infty$ through a nonlinear steepest descent analysis of RHP 3.8 and subsequent integration in (1.9).

5.1. *Initial steps.* We fix $\gamma \in (0, 1)$ and first rescale similar to (3.28),

$$(5.1) \quad \mathbf{T}(z; t, \gamma) := \mathbf{N}(-zt; t, \gamma), \quad z \in \mathbb{C} \setminus \mathbb{R}, t < 0$$

so that the jump condition for $\mathbf{T}(z; t, \gamma)$ reads as

$$(5.2) \quad \mathbf{T}_+(z) = \mathbf{T}_-(z) \begin{bmatrix} 1 - \gamma e^{-\frac{1}{2}t^2z^2} & -i\sqrt{\gamma}e^{-t^2\theta_+(z)} \\ -i\sqrt{\gamma}e^{-t^2\theta_-(z)} & 1 \end{bmatrix}, \quad z \in \mathbb{R}.$$

Thus, opposed to the $t \rightarrow +\infty$ analysis, the subscripts in $\theta_{\pm}(z)$ have flipped in the exponents, that is, transformation (3.28) is no longer helpful in view of the sign chart Figure 9. Instead we employ a g -function transformation which will swap the diagonal entries in (5.2) and modify the off-diagonal ones accordingly. In detail, the upcoming g -function will be defined in terms of the Cauchy transform of

$$h(x; t, \gamma) := -\ln(1 - \gamma e^{-\frac{1}{2}t^2x^2}), \quad x \in \mathbb{R}, \gamma \in (0, 1), t < 0,$$

and for this reason we shall collect a few of its relevant analytical properties below.

PROPOSITION 5.1. *For any $\gamma \in (0, 1)$ and $t < 0$, the function $h(\cdot; t, \gamma)$ exists in $L^p(\mathbb{R})$ for all $1 \leq p < \infty$ and is real-analytic. Moreover, using the principal branch of the logarithm, that is, $\ln : \mathbb{C} \setminus (-\infty, 0] \rightarrow \mathbb{C}$ with $\ln z = \ln |z| + i \arg z$ and $\arg z \in (-\pi, \pi]$, it extends analytically into the horizontal strip*

$$z \in \Pi_{t\gamma} = \left\{ z \in \mathbb{C} : |\Im z| < \frac{\sqrt{-2 \ln \gamma}}{|t|} \right\},$$

and we have the total integral identity

$$(5.3) \quad \int_{-\infty}^{\infty} h(x; t, \gamma) dx = \frac{\sqrt{2\pi}}{|t|} \text{Li}_{\frac{3}{2}}(\gamma),$$

in terms of the polylogarithm $\text{Li}_s(z)$; cf. [26], 25.12.10.

PROOF. Integrability on \mathbb{R} follows from the inequality $1 - \gamma e^{-\frac{1}{2}t^2x^2} \geq 1 - \gamma > 0$ and the estimate

$$h(x; t, \gamma) = \mathcal{O}(\gamma e^{-\frac{1}{2}t^2x^2}), \quad x \rightarrow \pm\infty.$$

For analyticity, we simply compute the preimage of $(-\infty, 0] \subset \mathbb{R}$ under the map $\mathbb{C} \ni z \mapsto 1 - \gamma e^{-\frac{1}{2}t^2z^2}$,

$$\begin{aligned} \Re(1 - \gamma e^{-\frac{1}{2}t^2z^2}) &= 1 - \gamma e^{-\frac{1}{2}t^2((\Re z)^2 - (\Im z)^2)} \cos(t^2 \Re z \Im z), \\ \Im(1 - \gamma e^{-\frac{1}{2}t^2z^2}) &= \gamma e^{-\frac{1}{2}t^2((\Re z)^2 - (\Im z)^2)} \sin(t^2 \Re z \Im z), \end{aligned}$$

that is, $1 - \gamma e^{-\frac{1}{2}t^2z^2}$ is purely real along the family of curve $\Gamma_{t,n} : t^2 \Re z \Im z = n\pi, n \in \mathbb{Z}$, so in particular real negative on the imaginary axis for $|\Im z| > \sqrt{-2 \ln \gamma}/|t|$ and along the parts of the black colored hyperbolas $\Gamma_{t,n}$ shown in Figure 11 that lie inside the red colored regions.




FIG. 11. The sign charts of $s(z; t, \gamma) := 1 - \gamma e^{-\frac{1}{2}t^2 z^2}$ with $\gamma = 0.9$ for $t = 1$ on the left and $t = 2$ on the right. In red values $z \in \mathbb{C}$ where $\Re s(z; t, \gamma) < 0$ and in blue where $\Re s(z; t, \gamma) > 0$. Also $\Im s(z; t, \gamma) = 0$ along the dash-dotted black curves.

Since $\Pi_{t,\gamma}$ excludes those parts, analyticity of $h(z; t, \gamma)$ follows easily and the remaining integral (5.3) is standard. \square

In order to proceed, we now introduce the g -function,

$$(5.4) \quad g(z) \equiv g(z; t, \gamma) := \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{h(s; t, \gamma)}{s - z} ds, \quad z \in \mathbb{C} \setminus \mathbb{R}$$

and consider the following transformation after (5.1):

$$(5.5) \quad \mathbf{S}(z; t, \gamma) := \mathbf{T}(z; t, \gamma) e^{g(z)\sigma_3}, \quad z \in \mathbb{C} \setminus \mathbb{R}.$$

Since $h(\cdot; t, \gamma)$ is of Hoelder class on \mathbb{R} for all $\gamma \in (0, 1)$ and $t < 0$, the classical Plemelj–Sokhotski formula applies and we arrive at the RHP below.

RIEMANN–HILBERT PROBLEM 5.2. For any $t < 0$ and $\gamma \in (0, 1)$, the function $\mathbf{S}(z) = \mathbf{S}(z; t, \gamma) \in \mathbb{C}^{2 \times 2}$ defined in (5.5) has the following properties:

(1) $\mathbf{S}(z)$ is analytic for $z \in \mathbb{C} \setminus \mathbb{R}$ and admits continuous boundary values on the closed upper and lower half-planes.

(2) Along the real axis, with $\mathbf{S}_{\pm}(z) = \lim_{\epsilon \downarrow 0} \mathbf{S}(z \pm i\epsilon)$, $z \in \mathbb{R}$,

$$(5.6) \quad \mathbf{S}_+(z) = \mathbf{S}_-(z) \begin{bmatrix} 1 & -i\sqrt{\gamma} e^{-t^2 \phi_+(z)} \\ -i\sqrt{\gamma} e^{-t^2 \phi_-(z)} & 1 - \gamma e^{-\frac{1}{2}t^2 z^2} \end{bmatrix}, \quad z \in \mathbb{R},$$

where

$$\phi_{\pm}(z) \equiv \phi_{\pm}(z; t, \gamma) := \theta_{\pm}(z) \pm \frac{1}{i\pi} \text{pv} \int_{-\infty}^{\infty} \frac{h(s; t, \gamma)}{t^2(s - z)} ds, \quad z \in \mathbb{R}.$$

(3) As $z \rightarrow \infty$, $z \notin \mathbb{R}$,

$$\mathbf{S}(z) = \mathbb{I} + \mathbf{S}_1(t, \gamma) z^{-1} + \mathcal{O}(z^{-2}), \quad \mathbf{S}_1(t, \gamma) = \mathbf{T}_1(t, \gamma) - \frac{i\sigma_3}{t\sqrt{2\pi}} \text{Li}_{\frac{3}{2}}(\gamma).$$

³We shall discard the trivial case $\gamma = 0$ for technical purposes.




FIG. 12. The oriented jump contour $\Sigma_{\mathbf{M}}$ in RHP 5.4. We fix $z_1 = -2$, $z_{2,3} = -1 \pm i\delta_{t\gamma}$, $z_{4,5} = 1 \pm i\delta_{t\gamma}$ and $z_6 = 2$ as location of the six vertices.

Our next step relies on the matrix factorization

$$\begin{aligned}
 (5.7) \quad & \begin{bmatrix} 1 & -i\sqrt{\gamma}e^{-t^2\phi_+(z)} \\ -i\sqrt{\gamma}e^{-t^2\phi_-(z)} & 1 - \gamma e^{-\frac{1}{2}t^2z^2} \end{bmatrix} \\
 & = \begin{bmatrix} 1 & 0 \\ -i\sqrt{\gamma}e^{-t^2\phi_-(z)} & 1 \end{bmatrix} \begin{bmatrix} 1 & -i\sqrt{\gamma}e^{-t^2\phi_+(z)} \\ 0 & 1 \end{bmatrix}, \quad z \in \mathbb{R},
 \end{aligned}$$

and the lemma below.

LEMMA 5.3. The function

$$\begin{aligned}
 \phi(z) \equiv \phi(z; t, \gamma) & := \frac{1}{4}(z^2 - 4iz \operatorname{sgn}(\Im z)) + \frac{1}{i\pi} \operatorname{sgn}(\Im z) \int_{-\infty}^{\infty} \frac{h(s; t, \gamma)}{t^2(s - z)} ds \\
 & - \frac{1}{t^2} h(z; t, \gamma), \quad z \in \Pi_{t\gamma} \setminus \mathbb{R}
 \end{aligned}$$

defined with the principal branch of the logarithm (see Proposition 5.1) is analytic in $\Pi_{t\gamma} \setminus \mathbb{R}$ for all $\gamma \in (0, 1)$ and $t < 0$. Moreover,

$$\lim_{\epsilon \downarrow 0} \phi(z \pm i\epsilon) = \phi_{\pm}(z), \quad z \in \mathbb{R}.$$

PROOF. Since $h(\cdot; t, \gamma)$ is of Hoelder class on \mathbb{R} and analytic in $\Pi_{t\gamma}$ by Proposition 5.1, the claims follow easily from the Plemelj–Sokhotski theorem. \square

More precisely, we introduce for any $\gamma \in (0, 1)$ and $t < 0$,

$$(5.8) \quad \mathbf{M}(z; t, \gamma) := \mathbf{S}(z; t, \gamma) \begin{cases} \begin{bmatrix} 1 & i\sqrt{\gamma}e^{-t^2\phi(z)} \\ 0 & 1 \end{bmatrix} & z \in \Omega_1 = \Omega_1(\delta_{t\gamma}), \\ \begin{bmatrix} 1 & 0 \\ -i\sqrt{\gamma}e^{-t^2\phi(z)} & 1 \end{bmatrix} & z \in \Omega_2 = \Omega_2(\delta_{t\gamma}), \\ \mathbb{I} & \text{else,} \end{cases}$$

$$\text{with } 0 < \delta_{t\gamma} := \min \left\{ \frac{\sqrt{-\ln \gamma}}{|t|}, \frac{1}{2} \right\},$$

which leads to the RHP formulated on the red contour $\Sigma_{\mathbf{M}}$ in Figure 12 and with the following characteristics.

RIEMANN–HILBERT PROBLEM 5.4. For any $\gamma \in (0, 1)$ and $t < 0$, the function $\mathbf{M}(z) = \mathbf{M}(z; t, \gamma) \in \mathbb{C}^{2 \times 2}$ has the following properties:

(1) $\mathbf{M}(z)$ is analytic for $z \in \mathbb{C} \setminus \Sigma_{\mathbf{M}}$ and all eight straight lines are oriented from left to right; see Figure 12.

(2) The square integrable limiting values obey

$$\mathbf{M}_+(z) = \mathbf{M}_-(z) \begin{bmatrix} 1 & -i\sqrt{\gamma}e^{-t^2\phi(z)} \\ 0 & 1 \end{bmatrix}, \quad z \in \Sigma_{\mathbf{M}} \cap \{z \in \mathbb{C} : \Im z > 0\};$$

and

$$\mathbf{M}_+(z) = \mathbf{M}_-(z) \begin{bmatrix} 1 & 0 \\ -i\sqrt{\gamma}e^{-t^2\phi(z)} & 1 \end{bmatrix}, \quad z \in \Sigma_{\mathbf{M}} \cap \{z \in \mathbb{C} : \Im z < 0\};$$

as well as

$$\mathbf{M}_+(z) = \mathbf{M}_-(z) \begin{bmatrix} 1 & -i\sqrt{\gamma}e^{-t^2\phi_+(z)} \\ -i\sqrt{\gamma}e^{-t^2\phi_-(z)} & 1 - \gamma e^{-\frac{1}{2}t^2z^2} \end{bmatrix}, \quad z \in \mathbb{R} \setminus [-2, 2].$$

By construction (see Lemma 5.3 and equation (5.7)), there is no jump along $(-2, 2) \subset \mathbb{R}$.

(3) For large $|z|$, the leading order asymptotic behavior of RHP 5.2 is unchanged,

$$\mathbf{M}(z) = \mathbb{I} + \mathbf{S}_1(t, \gamma)z^{-1} + \mathcal{O}(z^{-2}), \quad z \rightarrow \infty.$$

We now proceed with the necessary small norm estimates for the jump matrix $\mathbf{G}_{\mathbf{M}}(z; t, \gamma)$ in condition (2) of RHP 5.4. And since we are about to investigate the limit $\gamma \uparrow 1$ in the upcoming sections we shall already now indicate the γ -dependency in all error estimates.

PROPOSITION 5.5. *There exists positive universal constants t_0, c_j such that for any $\gamma \in (0, 1)$,*

$$\begin{aligned} (5.9) \quad & \|\mathbf{G}_{\mathbf{M}}(\cdot; t, \gamma) - \mathbb{I}\|_{L^\infty(\Sigma_{\mathbf{M}}, |dz|)} \leq c_1 \frac{\gamma^{\frac{1}{4}}e^{-t^2\delta_{t\gamma}}}{1 - \gamma}, \\ & \|\mathbf{G}_{\mathbf{M}}(\cdot; t, \gamma) - \mathbb{I}\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)} \leq c_2 \frac{\gamma^{\frac{1}{4}}e^{-t^2\delta_{t\gamma}}}{1 - \gamma} |t|^{-\frac{1}{2}}, \end{aligned}$$

hold true for all $(-t) \geq t_0$.

PROOF. For $z \in \mathbb{R} \setminus [-2, 2]$, we clearly have $\|\mathbf{G}_{\mathbf{M}}(\cdot; t, \gamma) - \mathbb{I}\| = \mathcal{O}(e^{-t^2})$ in $L^2 \cap L^\infty$, and thus no contribution to the leading order in (5.9). Next, for $z \in \mathbb{C} \setminus \mathbb{R}$,

$$e^{-t^2\phi(z)} = \frac{e^{-t^2\theta_\pm(z)}}{1 - \gamma e^{-\frac{1}{2}t^2z^2}} \exp\left[\pm \frac{i}{\pi} \int_{-\infty}^{\infty} \frac{h(s; t, \gamma)}{s - z} ds\right],$$

so we have to estimate three factors. With the parametrizations $z = z(\lambda) = \lambda \pm i\delta_{t\gamma}, \lambda \in [-1, 1]$, we find

$$|e^{-t^2\theta_\pm(z(\lambda))}| \leq e^{-t^2\delta_{t\gamma} - \frac{1}{4}t^2\lambda^2} \gamma^{-\frac{1}{4}},$$

$$|1 - \gamma e^{-\frac{1}{2}t^2z^2(\lambda)}| = [1 - 2\gamma e^{-\frac{1}{2}t^2(\lambda^2 - \delta_{t\gamma})} \cos(t^2\lambda\delta_{t\gamma}) + \gamma^2 e^{-t^2(\lambda^2 - \delta_{t\gamma}^2)}]^{\frac{1}{2}}$$

and

$$(5.10) \quad \left| \exp\left[\pm \frac{i}{\pi} \int_{-\infty}^{\infty} \frac{h(s; t, \gamma)}{s - z(\lambda)} ds\right] \right| = \exp\left[-\frac{\delta_{t\gamma}}{\pi} \int_{-\infty}^{\infty} \frac{h(s; t, \gamma)}{(s - \lambda)^2 + \delta_{t\gamma}^2} ds\right].$$

Hence,

$$\sup_{\lambda \in [-1, 1]} \left| \frac{e^{-t^2 \theta_{\pm}(z(\lambda))}}{1 - \gamma e^{-\frac{1}{2} t^2 z^2(\lambda)}} \right| \leq \frac{\gamma^{-\frac{1}{4}} e^{-t^2 \delta_{t\gamma}}}{1 - \sqrt{\gamma}},$$

and since the integrand in (5.10) is nonnegative we have therefore established the above L^∞ estimate. For the L^2 estimate, we use Laplace’s method for the factor $\exp[-t^2 \theta_{\pm}(z)]$ and the same reasonings that were used in the second and third while deriving the previous L^∞ estimate. For the four slanted straight lines consider, say, $z = z(\lambda) = \lambda - 2 \pm i\lambda \delta_{t\gamma}$ with $\lambda \in [0, 1]$. Then

$$|e^{-t^2 \theta_{\pm}(z(\lambda))}| = e^{-\frac{1}{4} t^2 (\lambda^2 (1 - \delta_{t\gamma}^2) - 4\lambda(1 - \delta_{t\gamma}) + 4)},$$

as well as

$$\begin{aligned} |1 - \gamma e^{-\frac{1}{2} t^2 z^2(\lambda)}| &= [1 - 2\gamma e^{-\frac{1}{2} t^2 ((\lambda - 2)^2 - \lambda^2 \delta_{t\gamma}^2)} \cos(t^2 \lambda (\lambda - 2) \delta_{t\gamma}) \\ &\quad + \gamma^2 e^{-t^2 ((\lambda - 2)^2 - \lambda^2 \delta_{t\gamma}^2)}]^{1/2}, \end{aligned}$$

and

$$\begin{aligned} (5.11) \quad & \left| \exp \left[\pm \frac{i}{\pi} \int_{-\infty}^{\infty} \frac{h(s; t, \gamma)}{s - z(\lambda)} ds \right] \right| \\ &= \exp \left[-\frac{\lambda \delta_{t\gamma}}{\pi} \int_{-\infty}^{\infty} \frac{h(s; t, \gamma)}{(s - \lambda + 2)^2 + \lambda^2 \delta_{t\gamma}^2} ds \right]. \end{aligned}$$

Thus

$$\sup_{\lambda \in [0, 1]} \left| \frac{e^{-t^2 \theta_{\pm}(z(\lambda))}}{1 - \gamma e^{-\frac{1}{2} t^2 z^2(\lambda)}} \right| \leq \frac{e^{-\frac{1}{4} t^2 (1 - \delta_{t\gamma}^2)}}{1 - \gamma e^{-\frac{1}{2} t^2 (1 - \delta_{t\gamma}^2)}},$$

which is of subleading order for $\gamma \in (0, 1)$ when compared to (5.9); see (5.7). By nonnegativity of the integrand in (5.11), we have therefore completed our proof. \square

Since we are dealing with a (t, γ) -dependent contour in RHP 5.4 (the hexagon in Figure 12 is collapsing to the real axis as $t \rightarrow -\infty$ or $\gamma \uparrow 1$), the general framework of [12] is not directly applicable to Proposition 5.5. Still, using somewhat similar ideas as in [7], the results of Appendix A below guarantee unique solvability of RHP 5.4 in $L^2(\Sigma_{\mathbf{M}}, |dz|)$ for all $(-t) \geq t_0$ and either any fixed $\gamma \in (0, 1)$ or $\gamma \uparrow 1$ at a certain controlled rate.

THEOREM 5.6. *For any fixed $0 < \epsilon < 2$, RHP 5.4 is uniquely solvable in $L^2(\Sigma_{\mathbf{M}}, |dz|)$ for $(-t)$ sufficiently large and all $0 < \gamma \leq 1 - |t|^{-\epsilon}$. Moreover,*

$$\begin{aligned} (5.12) \quad \mathbf{S}_1(t, \gamma) &= \lim_{z \rightarrow \infty} z(\mathbf{M}(z; t, \gamma) - \mathbb{I}) = \mathcal{O}(|t|^{-1+\epsilon} e^{-|t|^{1-\frac{\epsilon}{2}}}) \\ &\forall (-t) \geq t_0, 0 < \gamma \leq 1 - |t|^{-\epsilon}. \end{aligned}$$

5.2. Left tail asymptotics. In order to complete the derivation of (1.16), we first recall (3.14), transformations (3.18), (5.1), (5.5), (5.8),

$$\frac{\partial}{\partial t} \ln \det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) = \frac{1}{\sqrt{2\pi}} \text{Li}_{\frac{3}{2}}(\gamma) + it S_1^{22}(t, \gamma).$$

Thus, with Theorem 5.6 and an indefinite t -integration,

PROPOSITION 5.7. For any fixed $0 < \epsilon < 2$, there exist positive constants $t_0 = t_0(\epsilon)$ and $c = c(\epsilon)$ such that

$$\ln \det(1 - \gamma T \chi_t \upharpoonright_{L^2(\mathbb{R})}) = \frac{t}{\sqrt{2\pi}} \text{Li}_{\frac{3}{2}}(\gamma) + D_1(\gamma) + r_1(t, \gamma),$$

for all $(-t) \geq t_0$ and $0 < \gamma \leq 1 - |t|^{-\epsilon}$. The term $D_1(\gamma)$ is t -independent and we record the error estimate

$$|r_1(t, \gamma)| \leq c(\epsilon) |t|^{\frac{3}{2}\epsilon} e^{-|t|^{1-\frac{\epsilon}{2}}} \quad \forall (-t) \geq t_0 \text{ and } 0 < \gamma \leq 1 - |t|^{-\epsilon}.$$

Next, we use the estimate (based again on the transformations (3.18), (5.1), (5.5), (5.8) and Theorem 5.6)

$$\begin{aligned} \int_t^\infty Y_1^{12}(x, \gamma) dx &= I(\gamma) - \int_{-\infty}^t Y_1^{12}(x, \gamma) dx = I(\gamma) + \int_{-\infty}^t x S_1^{12}(x, \gamma) dx \\ &= I(\gamma) + \mathcal{O}(|t|^{\frac{3}{2}\epsilon} e^{-|t|^{1-\frac{\epsilon}{2}}}), \end{aligned}$$

where $I(\gamma)$ denotes the total integral of $Y_1^{12}(x, \gamma)$ over $x \in \mathbb{R}$. From this, with (4.19) and $\Gamma_{t\gamma} = 1 - u^\gamma(t)$ we obtain in turn

PROPOSITION 5.8. For any fixed $0 < \epsilon < 2$, there exist positive constants $t_0 = t_0(\epsilon)$ and $c = c(\epsilon)$ such that

$$\ln \Gamma_{t\gamma} = D_2(\gamma) + r_2(t, \gamma)$$

for all $(-t) \geq t_0$ and $0 < \gamma \leq 1 - |t|^{-\epsilon}$. The term $D_2(\gamma)$ is t -independent and we record the error estimate

$$|r_2(t, \gamma)| \leq c(\epsilon) |t|^{\frac{3}{2}\epsilon} e^{-|t|^{1-\frac{\epsilon}{2}}} \quad \forall (-t) \geq t_0 \text{ and } 0 < \gamma \leq 1 - |t|^{-\epsilon}.$$

Once we combine Propositions 5.7 and 5.8, Corollary 1.10, expansion (1.16) follows easily.

6. Proof of Theorem 1.12. The following lines are near copies of the argument given in [16] in the derivation of the analogue of (1.17) for the GOE. First, by (1.1) and (1.3),

$$(6.1) \quad (F(t))^2 = \det(1 - T \chi_t \upharpoonright_{L^2(\mathbb{R})}) \Gamma_t = \det(1 - (T + U \otimes V) \upharpoonright_{L^2(t, \infty)}),$$

where $U \otimes V$ denotes the finite rank integral operator on $L^2(t, \infty)$ with kernel

$$(U \otimes V)(x, y) = g(x)G(y), \quad g(x) = \frac{1}{\sqrt{\pi}} e^{-x^2}, \quad G(x) = \int_{-\infty}^x g(y) dy,$$

that is, U is the operator which multiplies by $g(x)$ and V the integral operator with kernel $G(y)$. Indeed, (6.1) follows by noting that for any operator A we have $A(U \otimes V) = (AU \otimes V)$, applying the factorization

$$(6.2) \quad \begin{aligned} &\det(1 - (T + U \otimes V) \upharpoonright_{L^2(t, \infty)}) \\ &= \det(1 - T \upharpoonright_{L^2(t, \infty)}) \det(1 - (1 - T)^{-1}(U \otimes V) \upharpoonright_{L^2(t, \infty)}), \end{aligned}$$

and using the eigenvector/value equation

$$(6.3) \quad \begin{aligned} &((1 - T)^{-1}(U \otimes V) \upharpoonright_{L^2(t, \infty)} F)(x) \\ &= \left[\int_t^\infty G(y) ((1 - T \upharpoonright_{L^2(t, \infty)})^{-1} g)(y) dy \right] F(x), \end{aligned}$$

where

$$F(x) = ((1 - T \upharpoonright_{L^2(t,\infty)})^{-1}g)(x).^4$$

Precisely, (6.3) computes the finite rank operator determinant in (6.2) as

$$\begin{aligned} & \det(1 - (1 - T)^{-1}(U \otimes V) \upharpoonright_{L^2(t,\infty)}) \\ &= 1 - \int_t^\infty G(y)((1 - T \upharpoonright_{L^2(t,\infty)})^{-1}g)(y) \, dy = \Gamma_t, \end{aligned}$$

and (6.1) follows from (6.2) since $\det(1 - T \upharpoonright_{L^2(t,\infty)}) = \det(1 - T \chi_t \upharpoonright_{L^2(\mathbb{R})})$. Second, it will be more convenient to move the t -dependency in the right-hand side of (6.1) into the integral operators,

$$(6.4) \quad (F(t))^2 = \det(1 - (T_t + U_t \otimes V_t) \upharpoonright_{L^2(0,\infty)}), \quad t \in \mathbb{R},$$

where $T_t : L^2(0, \infty) \rightarrow L^2(0, \infty)$ has kernel $T_t(x, y) = T(x + t, y + t)$, compare (1.2), U_t is multiplication by $g(x + t)$ and V_t has kernel $G(y + t)$. Third, we note that $T_t = S_t S_t$ where $S_t : L^2(0, \infty) \rightarrow L^2(0, \infty)$ has kernel

$$S_t(x, y) = \frac{1}{\sqrt{\pi}} e^{-(x+y+t)^2}, \quad x, y > 0.$$

LEMMA 6.1. *For every $t \in \mathbb{R}$, the operator S_t satisfies $\|S_t\| \leq 1$ and $1 \mp S_t$ are invertible on $L^2(0, \infty)$.*

PROOF. Since S_t is self-adjoint, we have for any $f \in L^2(0, \infty)$,

$$\begin{aligned} \|S_t f\|_{L^2(0,\infty)}^2 &= \langle S_t f, S_t f \rangle_{L^2(0,\infty)} = \langle f, S_t^2 f \rangle_{L^2(0,\infty)} = \langle f, T_t f \rangle_{L^2(0,\infty)} \\ &\stackrel{(2.3)}{\leq} \langle f, f \rangle_{L^2(0,\infty)} \end{aligned}$$

and, therefore,

$$\|S_t\| = \sup_{\|f\|_{L^2(0,\infty)}=1} \|S_t f\|_{L^2(0,\infty)} \leq 1.$$

Also, $(1 - S_t)(1 + S_t) = (1 + S_t)(1 - S_t) = 1 - T_t$ on $L^2(0, \infty)$ and since $1 - T_t$ is invertible by Lemma 2.1, so are $1 \mp S_t$. \square

The last lemma allows us to transform the right-hand side in (6.4) through the following factorization:

$$\begin{aligned} & \det(1 - (S_t^2 + U_t \otimes V_t) \upharpoonright_{L^2(0,\infty)}) \\ (6.5) \quad &= \det(1 - S_t \upharpoonright_{L^2(0,\infty)}) \det(1 + S_t - (1 - S_t)^{-1}(U_t \otimes V_t) \upharpoonright_{L^2(0,\infty)}) \\ &= \det(1 - S_t \upharpoonright_{L^2(0,\infty)}) \det(1 + S_t - (U_t \otimes W) \upharpoonright_{L^2(0,\infty)}), \end{aligned}$$

⁴Identity (6.3) follows directly from the definitions of the underlying operators U, V and $(1 - T)^{-1}$. A similar computation in the GOE can be found in [21], Chapter 9.7, (9.128).

where W multiplies by the characteristic function $\chi_0(x)$.⁵ To get to (6.5), we have used that for any operator A we have $A(U \otimes V) = (U \otimes A^*V)$ in terms of the real adjoint A^* and that

$$G(x+t) = 1 - \int_0^\infty S_t(x, y) dy = ((1 - S_t \upharpoonright_{L^2(0, \infty)})\chi_0)(x),$$

$$\chi_0(x) = \begin{cases} 1 & x \geq 0, \\ 0 & x < 0. \end{cases}$$

Continuing with (6.5), another factorization yields

$$\begin{aligned} & \det(1 + S_t - (U_t \otimes W) \upharpoonright_{L^2(0, \infty)}) \\ &= \det(1 + S_t \upharpoonright_{L^2(0, \infty)}) \det(1 - (1 + S_t)^{-1}(U_t \otimes W) \upharpoonright_{L^2(0, \infty)}) \\ (6.6) \quad &= \det(1 + S_t \upharpoonright_{L^2(0, \infty)}) \\ &\quad \times \left\{ 1 - \int_0^\infty \chi_0(y) ((1 + S_t \upharpoonright_{L^2(0, \infty)})^{-1}g)(y+t) dy \right\}, \end{aligned}$$

where we have used a variation of the eigenvector/value trick (6.3) in the last equality. Since

$$g(x+t) = \int_0^\infty S_t(x, y)\delta_0(y) dy = (S_t\delta_0)(x); \quad \int_0^\infty f(x)\delta_0(x) dx := f(0),$$

for any test function f we can simplify the second factor in (6.6) further,

$$\begin{aligned} & 1 - \int_0^\infty \chi_0(y) ((1 + S_t \upharpoonright_{L^2(0, \infty)})^{-1}g)(y+t) dy \\ (6.7) \quad &= \int_0^\infty \chi_0(y) ((1 - (1 + S_t)^{-1}S_t \upharpoonright_{L^2(0, \infty)})\delta_0)(y) dy \\ &= \int_0^\infty \chi_0(y) ((1 + S_t \upharpoonright_{L^2(0, \infty)})^{-1}\delta_0)(y) dy \\ &= \langle \chi_0, (1 + S_t \upharpoonright_{L^2(0, \infty)})^{-1}\delta_0 \rangle_{L^2(0, \infty)}. \end{aligned}$$

The proof of Theorem 1.12 would thus be completed through (6.5), (6.6), (6.7) and the identity $\det(1 - S_t \upharpoonright_{L^2(0, \infty)}) = \det(1 - S\chi_t \upharpoonright_{L^2(\mathbb{R})})$ if we manage to prove the following:

$$\begin{aligned} (6.8) \quad & \det(1 - S_t \upharpoonright_{L^2(0, \infty)}) \\ &= \det(1 + S_t \upharpoonright_{L^2(0, \infty)}) \langle \chi_0, (1 + S_t \upharpoonright_{L^2(0, \infty)})^{-1}\delta_0 \rangle_{L^2(0, \infty)}, \quad t \in \mathbb{R}; \end{aligned}$$

or equivalently (taking logarithmic derivatives, then observing the unity normalization of all three factors in (6.8) as $t \rightarrow +\infty$, using Lemma 6.1 and self-adjointness),

$$(6.9) \quad \operatorname{tr}_{L^2(0, \infty)} \left((1 - S_t^2)^{-1} \frac{dS_t}{dt} \right) = -\frac{1}{2} \frac{d}{dt} \ln \langle \delta_0, (1 + S_t \upharpoonright_{L^2(0, \infty)})^{-1}\chi_0 \rangle_{L^2(0, \infty)}.$$

But integrating by parts in the left-hand side of (6.9) with $\frac{d}{dt} S_t(x, y) = \frac{d}{dx} S_t(x, y) = \frac{d}{dy} S_t(x, y)$ shows that (cf. [16], Lemma 2),

$$\operatorname{tr}_{L^2(0, \infty)} \left((1 - S_t^2)^{-1} \frac{dS_t}{dt} \right) = -\frac{1}{2} \langle \delta_0, ((1 - S_t^2)^{-1}S_t \upharpoonright_{L^2(0, \infty)})\delta_0 \rangle_{L^2(0, \infty)},$$

⁵Evidently, χ_0 and the later on used δ_0 are not in $L^2(0, \infty)$. Still, using regularity and decay properties of the involved integral kernels all subsequent Fredholm determinant and inner product manipulations are justifiable, we refer the interested reader to [33], Section VIII.

so we need to establish the equality

$$\begin{aligned}
 (6.10) \quad & \langle \delta_0, ((1 - S_t^2)^{-1} S_t \upharpoonright_{L^2(0,\infty)}) \delta_0 \rangle_{L^2(0,\infty)} \\
 &= \frac{d}{dt} \ln \langle \delta_0, (1 + S_t \upharpoonright_{L^2(0,\infty)})^{-1} \chi_0 \rangle_{L^2(0,\infty)}.
 \end{aligned}$$

LEMMA 6.2 ([16], Lemma 3). *Let Δ_0 denote multiplication by $\delta_0(x)$, and $D (= \frac{d}{dx})$ differentiation. Then*

$$(6.11) \quad \frac{d}{dt} (1 + S_t)^{-1} = (1 - S_t^2)^{-1} S_t D + (1 - S_t^2)^{-1} S_t \Delta_0 (1 + S_t)^{-1}.$$

PROOF. Integrating by parts, we easily find that $DS_t = -S_t \Delta_0 - S_t D$, or equivalently

$$-(1 - S_t) DS_t = S_t D (1 + S_t) + S_t \Delta_0.$$

Multiplying this last identity by $(1 - S_t^2)^{-1}$ from the left and $(1 + S_t)^{-1}$ from the right (recall Lemma 6.1),

$$-(1 + S_t)^{-1} DS_t (1 + S_t)^{-1} = (1 - S_t^2)^{-1} S_t D + (1 - S_t^2)^{-1} S_t \Delta_0 (1 + S_t)^{-1},$$

and the last left-hand side equals precisely $\frac{d}{dt} (1 + S_t)^{-1}$ because of the simple fact that

$$\frac{d}{dt} (1 + S_t)^{-1} = -(1 + S_t)^{-1} \frac{dS_t}{dt} (1 + S_t)^{-1},$$

and $\frac{d}{dt} S_t(x, y) = \frac{d}{dx} S_t(x, y)$. Identity (6.11) is thus proven. \square

We now use (6.11) in the right-hand side of (6.10) (suppressing at times the Hilbert space references for compact notation),

$$\begin{aligned}
 & \frac{d}{dt} \ln \langle \delta_0, (1 + S_t \upharpoonright_{L^2(0,\infty)})^{-1} \chi_0 \rangle_{L^2(0,\infty)} \\
 &= \frac{\langle \delta_0, \frac{d}{dt} (1 + S_t \upharpoonright_{L^2(0,\infty)})^{-1} \chi_0 \rangle_{L^2(0,\infty)}}{\langle \delta_0, (1 + S_t \upharpoonright_{L^2(0,\infty)})^{-1} \chi_0 \rangle_{L^2(0,\infty)}} \\
 &\stackrel{D\chi_0=0}{=} \frac{\langle \delta_0, (1 - S_t^2)^{-1} S_t \Delta_0 (1 + S_t)^{-1} \chi_0 \rangle}{\langle \delta_0, (1 + S_t)^{-1} \chi_0 \rangle} \\
 &= \frac{\langle \delta_0, (1 - S_t^2)^{-1} S_t \delta_0 \rangle \langle \delta_0, (1 + S_t)^{-1} \chi_0 \rangle}{\langle \delta_0, (1 + S_t)^{-1} \chi_0 \rangle} \\
 &= \langle \delta_0, ((1 - S_t^2)^{-1} S_t \upharpoonright_{L^2(0,\infty)}) \delta_0 \rangle_{L^2(0,\infty)},
 \end{aligned}$$

which is (6.10). In turn, the proof of Theorem 1.12 is thus completed.

APPENDIX A: SMALL NORM ESTIMATES FOR COLLAPSING CONTOURS

The jump contour Σ_M of RHP 5.4 collapses for large $(-t)$ to the real axis and we shall invoke ideas from [7] in the solution of the underlying singular integral equation ([7] does not apply verbatim to RHP 5.4 as we are not dealing with contracting disk contours, for those the scaling invariance of $z^{-1} dz$ is central).

Write $\Sigma_M = \bigcup_{j=1}^8 \Sigma_j$ as union of the eight straight lines shown in Figure 13 below and suppose $\rho \in L^2(\Sigma_M, |dz|)$ is a 2×2 matrix-valued function which is Lipschitz on Σ_j and that satisfies

$$(A.1) \quad \rho(z) = \mathbb{I} + \frac{1}{2\pi i} \int_{\Sigma_M} \rho(\lambda) (\mathbf{G}_M(\lambda) - \mathbb{I}) \frac{d\lambda}{\lambda - z_-}, \quad z \in \Sigma_j,$$




FIG. 13. The oriented jump contour $\Sigma_{\mathbf{M}} = \bigcup_{j=1}^8 \Sigma_j$ in RHP 5.4.

where z_- denotes the limiting value of the integral from the right-hand side. The explicit form of the jump matrix $\mathbf{G}_{\mathbf{M}}(z) = \mathbf{G}_{\mathbf{M}}(z; t, \gamma)$ is stated in RHP 5.4, condition (2).

LEMMA A.1. On each Σ_j , the function

$$\mathbf{M}(z) := \mathbb{I} + \frac{1}{2\pi i} \int_{\Sigma_{\mathbf{M}}} \rho(\lambda)(\mathbf{G}_{\mathbf{M}}(\lambda) - \mathbb{I}) \frac{d\lambda}{\lambda - z}, \quad z \in \mathbb{C} \setminus \Sigma_{\mathbf{M}},$$

satisfies

$$\mathbf{M}_+(z) = \mathbf{M}_-(z)\mathbf{G}_{\mathbf{M}}(z), \quad z \in \Sigma_j.$$

PROOF. Since $\mathbf{M}_-(z) = \rho(z)$ for $z \in \Sigma_j$ the jump behavior of the Cauchy transform implies

$$\begin{aligned} \mathbf{M}_+(z) - \mathbf{M}_-(z) &= \rho(z)(\mathbf{G}_{\mathbf{M}}(z) - \mathbb{I}) \\ &= \mathbf{M}_-(z)(\mathbf{G}_{\mathbf{M}}(z) - \mathbb{I}), \quad z \in \Sigma_j. \end{aligned} \quad \square$$

We shall now solve (A.1) in $L^2(\Sigma_{\mathbf{M}}, |dz|)$ by the Neumann series

$$\begin{aligned} \rho(z) &= \mathbb{I} + \sum_{k=1}^{\infty} \rho_k(z); \\ \rho_k(z) &= \frac{1}{2\pi i} \int_{\Sigma_{\mathbf{M}}} \rho_{k-1}(\lambda)(\mathbf{G}_{\mathbf{M}}(\lambda) - \mathbb{I}) \frac{d\lambda}{\lambda - z_-}, \quad z \in \Sigma_{\mathbf{M}}, k \in \mathbb{Z}_{\geq 1} \\ \rho_0(z) &= \mathbb{I}, \end{aligned}$$

and thus need to estimate $\rho_k(z)$. Recall that $L^2(\Sigma_{\mathbf{M}}, |dz|)$ is the space of (matrix-valued) measurable functions such that

$$\|f\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)} = \left\{ \int_{\Sigma_{\mathbf{M}}} \|f(z)\|^2 |dz| \right\}^{\frac{1}{2}} < \infty.$$

Let $C_{\Sigma_j}^{\pm}$ denote the Cauchy operators on $L^2(\Sigma_{\mathbf{M}}, |dz|)$,

$$(C_{\Sigma_j}^{\pm} \rho)(z) = \lim_{\epsilon \downarrow 0} \int_{\Sigma_j} \frac{\rho(s)}{s - (z \pm i\epsilon)} \frac{ds}{2\pi i}, \quad z \in \Sigma_j,$$

which obey (cf. [30], Chapter II, or [2], Section 5.5)

$$\begin{aligned} (A.2) \quad C_{\Sigma_j}^+ - C_{\Sigma_j}^- &= \text{id}, \quad C_{\Sigma_j}^+ + C_{\Sigma_j}^- = iH_{\Sigma_j}, \quad \text{a.e. on } \Sigma_j; \\ (H_{\Sigma_j} \rho)(z) &= \frac{1}{\pi} \text{pv} \int_{\Sigma_j} \rho(\lambda) \frac{ds}{z - s}. \end{aligned}$$

PROPOSITION A.2 ([11], Theoreme I). *If an oriented contour $\Gamma \subset \mathbb{C}$, given by the parametric equations*

$$\Gamma = \{z \in \mathbb{C} : \Re z = t \in \mathbb{R}, \Im z = \phi(t)\}$$

satisfies a uniform Lipschitz condition, that is, there exists $M \geq 0$ such that

$$|\phi(x) - \phi(y)| \leq M|x - y|,$$

then there exists a universal $C_0 > 0$ such that

$$(A.3) \quad \|H_\Gamma f\|_{L^2(\Gamma, |dz|)} \leq C_0(1 + M)^{10} \|f\|_{L^2(\Gamma, |dz|)}.$$

Observe that our eight pieces Σ_j fit into the context of Proposition A.2 with a (t, γ) -independent constant M , thus we are now prepared to derive our central estimate.

THEOREM A.3. *For any $\gamma \in (0, 1)$ and $t < 0$, let $\Sigma_{\mathbf{M}} = \bigcup_{j=1}^8 \Sigma_j$ denote the (t, γ) -dependent contour of RHP 5.4. Then there exists a universal constant $K > 0$ such that*

$$(A.4) \quad \|H_{\Sigma_{\mathbf{M}}} f\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)} \leq K \ln(\delta_{t\gamma}^{-1}) \|f\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)}.$$

PROOF. We show that for some $\widehat{K} > 0$,

$$\begin{aligned} & \left| \langle H_{\Sigma_{\mathbf{M}}}(\chi_{\Sigma_j} f), \chi_{\Sigma_k} g \rangle_{L^2(\Sigma_{\mathbf{M}}, |dz|)} \right| \\ & \leq \widehat{K} \ln(\delta_{t\gamma}^{-1}) \|f\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)} \|g\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)}, \quad 1 \leq j, k \leq 8. \end{aligned}$$

Indeed, for $j = k$ this follows at once from (A.3) and for $j \neq k$ we use the following estimates (derived from the Cauchy–Schwarz inequality while using polar coordinates and standard manipulations)

$$\begin{aligned} & \int_{-1}^1 \int_{-1}^1 \frac{|f(x + i\delta_{t\gamma})g(y - i\delta_{t\gamma})|}{\sqrt{(x - y)^2 + 4\delta_{t\gamma}^2}} dx dy \\ & \leq C_1 \ln(\delta_{t\gamma}^{-1}) \|f\|_{L^2(\Sigma_4, |dz|)} \|g\|_{L^2(\Sigma_5, |dz|)}; \\ & \int_0^\infty \int_0^\infty \frac{|f(x)g(y)|}{\sqrt{(x - y)^2 + \delta_{t\gamma}^2 x^2}} dx dy \\ & \leq C_2 \ln(\delta_{t\gamma}^{-1}) \|f\|_{L^2(0, \infty)} \|g\|_{L^2(0, \infty)}, \quad C_j > 0; \\ & \int_0^\infty \int_0^\infty \frac{|f(x)g(y)|}{x + y} dx dy \leq C_3 \|f\|_{L^2(0, \infty)} \|g\|_{L^2(0, \infty)}. \end{aligned} \quad \square$$

From estimate (A.4), we derive the operator norm estimate

$$\|H_{\Sigma_{\mathbf{M}}}\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)} \leq K \ln(\delta_{t\gamma}^{-1}), \quad K > 0 \text{ universal}$$

and then in turn, with (A.2),

$$(A.5) \quad \|C_{\Sigma_{\mathbf{M}}}^\pm\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)} \leq \frac{1}{2}(K + 1) \ln(\delta_{t\gamma}^{-1}) \quad \forall (-t) \geq t_0, \gamma \in (0, 1).$$

Returning now to the iterates $\{\rho_k(z)\}_{k=0}^\infty$ introduced above, we have

$$\rho_k(z) = (C_{\Sigma_{\mathbf{M}}}^- [\rho_{k-1}(\mathbf{G}_{\mathbf{M}} - \mathbb{I})])(z), \quad z \in \Sigma_{\mathbf{M}}.$$

But

$$\|\rho_{k-1}(\mathbf{G}_{\mathbf{M}} - \mathbb{I})\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)} \leq \|\mathbf{G}_{\mathbf{M}}(\cdot; t, \gamma) - \mathbb{I}\|_{L^\infty(\Sigma_{\mathbf{M}})} \cdot \|\rho_{k-1}\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)},$$

and for all $(-t) \geq t_0, \gamma \in (0, 1)$,

$$\begin{aligned} \|\rho_1\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)} &= \|C_{\Sigma_{\mathbf{M}}}^-(\mathbf{G}_{\mathbf{M}} - \mathbb{I})\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)} \\ &\leq \frac{1}{2}(K + 1) \ln(\delta_{t\gamma}^{-1}) \cdot \|\mathbf{G}_{\mathbf{M}}(\cdot; t, \gamma) - \mathbb{I}\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)}, \end{aligned}$$

so that with Proposition 5.5,

PROPOSITION A.4. *There exist positive universal constants t_0, c such that for any $\gamma \in (0, 1)$,*

$$\|\rho_k\|_{L^2(\Sigma_{\mathbf{M}_\gamma}, |dz|)} \leq \left(c \ln(\delta_{t\gamma}^{-1}) \frac{e^{-t^2 \delta_{t\gamma}}}{1 - \gamma} \right)^k |t|^{-\frac{1}{2}} \quad \forall (-t) \geq t_0, k \in \mathbb{Z}_{\geq 1}.$$

Thus, given any $0 < \epsilon < 2$, Proposition A.4 implies convergence of the Neumann series $\mathbb{I} + \sum_{k=1}^{\infty} \rho_k(z)$ in $L^2(\Sigma_{\mathbf{M}}, |dz|)$ for sufficiently large $(-t) \geq t_0$ and any $\gamma \in (0, 1)$ such that $0 < \gamma \leq 1 - |t|^{-\epsilon}$. At this point, using $\rho(z)$, we define

$$(A.6) \quad \mathbf{M}(z) := \mathbb{I} + \frac{1}{2\pi i} \int_{\Sigma_{\mathbf{M}}} \rho(\lambda)(\mathbf{G}_{\mathbf{M}}(\lambda) - \mathbb{I}) \frac{d\lambda}{\lambda - z}, \quad z \in \mathbb{C} \setminus \Sigma_{\mathbf{M}},$$

which coincides with the function $\mathbf{M}(z)$ defined by (5.7) (compare Lemma A.1 and the argument in [7], (A.37)–(A.39), near a triple point, a point on $\Sigma_{\mathbf{M}}$ where three arcs meet). Thus, compare RHP 5.4,

$$\mathbf{S}_1(t, \gamma) = \lim_{z \rightarrow \infty} z(\mathbf{M}(z) - \mathbb{I}) = \frac{i}{2\pi} \int_{\Sigma_{\mathbf{M}}} \rho(\lambda)(\mathbf{G}_{\mathbf{M}}(\lambda) - \mathbb{I}) d\lambda = \sum_{k=0}^{\infty} \mathbf{S}_{1k}(t, \gamma),$$

where

$$\mathbf{S}_{1k}(t, \gamma) = \frac{i}{2\pi} \int_{\Sigma_{\mathbf{M}}} \rho_k(\lambda)(\mathbf{G}_{\mathbf{M}}(\lambda) - \mathbb{I}) d\lambda.$$

Since for $(-t) \geq t_0$ and $\gamma \in (0, 1)$, with $C_j > 0$,

$$\begin{aligned} k \geq 1: \quad \|\mathbf{S}_{1k}(t, \gamma)\| &\leq C_1 \|\rho_k\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)} \|\mathbf{G}_{\mathbf{M}}(\cdot; t, \gamma) - \mathbb{I}\|_{L^2(\Sigma_{\mathbf{M}}, |dz|)} \\ &\leq C_1 \left(c \ln(\delta_{t\gamma}^{-1}) \frac{e^{-t^2 \delta_{t\gamma}}}{1 - \gamma} \right)^k \frac{e^{-t^2 \delta_{t\gamma}}}{1 - \gamma} |t|^{-1}, \end{aligned}$$

$$\begin{aligned} k = 0: \quad \|\mathbf{S}_{1k}(t, \gamma)\| &\leq C_2 \|\mathbf{G}_{\mathbf{M}}(\cdot; t, \gamma) - \mathbb{I}\|_{L^1(\Sigma_{\mathbf{M}}, |dz|)} \\ &\leq C_2 \frac{e^{-t^2 \delta_{t\gamma}}}{1 - \gamma} |t|^{-1}, \end{aligned}$$

we can now sum all inequalities from $k = 0$ to $k = +\infty$: for any fixed $\epsilon \in (0, 2)$,

$$\mathbf{S}_1(t, \gamma) = \mathcal{O}(|t|^{-1+\epsilon} e^{-|t|^{1-\frac{\epsilon}{2}}}) \quad \forall (-t) \geq t_0, 0 < \gamma \leq 1 - |t|^{-\epsilon}.$$

This completes the proof of Theorem 5.6.

APPENDIX B: PERMUTING RESOLVENT AND INTEGRATION

Let ϕ and ψ be two functions on \mathbb{R} that decay exponentially fast at $+\infty$. Introduce

$$K(x, y) := \int_0^{\infty} \phi(x + s)\psi(y + s) ds, \quad x, y \in \mathbb{R},$$

and the corresponding integral operator K on $L^2(\mathbb{R})$ with kernel $K(x, y)$. For any function f , we shall denote by $f_y(x) := f(x + y)$ the horizontal translation of f by $-y$. Then we have the following.

PROPOSITION B.1. For any $x, y, t \in \mathbb{R}$ and $k \in \mathbb{Z}_{\geq 1}$, we have

$$(B.1) \quad ((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \phi_y)(x) = ((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \phi_{x-t})_y(t).$$

PROOF. We prove (B.1) inductively using

$$K(x, y) = \int_t^\infty \phi(x + u - t) \psi(y + u - t) \, du.$$

Indeed, for $k = 1$, we have

$$\begin{aligned} (K \chi_t \upharpoonright_{L^2(\mathbb{R})} \phi_y)(x) &= \int_t^\infty K(x, s) \phi(s + y) \, ds \\ &= \int_t^\infty \int_t^\infty \phi(x + u - t) \psi(s + u - t) \phi(s + y) \, du \, ds, \end{aligned}$$

so that by Fubini's theorem,

$$(K \chi_t \upharpoonright_{L^2(\mathbb{R})} \phi_y)(x) = \int_t^\infty K(y + t, u) \phi(u + x - t) \, du = (K \chi_t \upharpoonright_{L^2(\mathbb{R})} \phi_{x-t})_y(t).$$

For general k , assuming that (B.1) is true for $k - 1$,

$$\begin{aligned} &((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \phi_y)(x) \\ &= \int_t^\infty K(x, s) ((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^{k-1} \phi_y)(s) \, ds \\ &= \int_t^\infty K(x, s) ((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^{k-1} \phi_{s-t})_y(t) \, ds \\ &= \int_t^\infty \int_t^\infty K(x, s) (K \chi_t \upharpoonright_{L^2(\mathbb{R})})^{k-1}(t + y, u) \phi(u + s - t) \, du \, ds, \end{aligned}$$

and thus again by Fubini's theorem

$$((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \phi_y)(x) = \int_t^\infty (K \chi_t \upharpoonright_{L^2(\mathbb{R})})^{k-1}(t + y, u) (K \chi_t \upharpoonright_{L^2(\mathbb{R})} \phi_{u-t})(x) \, du.$$

Now apply the base case $k = 1$ result and derive

$$\begin{aligned} ((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \phi_y)(x) &= \int_t^\infty (K \chi_t \upharpoonright_{L^2(\mathbb{R})})^{k-1}(t + y, u) (K \chi_t \upharpoonright_{L^2(\mathbb{R})} \phi_{x-t})(u) \, du \\ &= ((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \phi_{x-t})(t + y) = ((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \phi_{x-t})_y(t), \end{aligned}$$

which completes the proof. \square

The above Proposition B.1 leads to the following useful corollary.

COROLLARY B.2. Let

$$\Phi(x) := \int_I \phi_y(x) \, dy,$$

where I is a subset of \mathbb{R} . Then, for any $x, t \in \mathbb{R}$ and $k \in \mathbb{Z}_{\geq 1}$,

$$(B.2) \quad ((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \Phi)(x) = \int_I ((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \phi_{x-t})_y(t) \, dy.$$

PROOF. We have by linearity and (B.1),

$$\begin{aligned} ((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \Phi)(x) &= (K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \left[\int_I \phi_y(\cdot) \, dy \right](x) \\ &= \int_I ((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \phi_y)(x) \, dy \\ &\stackrel{(B.1)}{=} \int_I ((K \chi_t \upharpoonright_{L^2(\mathbb{R})})^k \phi_{x-t})_y(t) \, dy. \end{aligned} \quad \square$$

Acknowledgments. T. Bothner acknowledges support of the AMS and the Simons Foundation through a travel grant and is grateful to B. Rider, P. Deift, and P. Miller for stimulating discussions about this project. Both authors initiated this work during the 2017 PCMI summer session on random matrices, funded in part by the NSF Grant DMS-1441467. The authors would also like to thank P. Forrester for bringing [22] to their attention.

The first author was supported by the NSF Grant DMS-1664692.

REFERENCES

- [1] ABLOWITZ, M. J. and CLARKSON, P. A. (1991). *Solitons, Nonlinear Evolution Equations and Inverse Scattering*. London Mathematical Society Lecture Note Series **149**. Cambridge Univ. Press, Cambridge. MR1149378 <https://doi.org/10.1017/CBO9780511623998>
- [2] BAIK, J., DEIFT, P. and SUIDAN, T. (2016). *Combinatorics and Random Matrix Theory*. Graduate Studies in Mathematics **172**. Amer. Math. Soc., Providence, RI. MR3468920
- [3] BEALS, R. and COIFMAN, R. R. (1984). Scattering and inverse scattering for first order systems. *Comm. Pure Appl. Math.* **37** 39–90. MR0728266 <https://doi.org/10.1002/cpa.3160370105>
- [4] BEALS, R., DEIFT, P. and TOMEI, C. (1988). *Direct and Inverse Scattering on the Line*. Mathematical Surveys and Monographs **28**. Amer. Math. Soc., Providence, RI. MR0954382 <https://doi.org/10.1090/surv/028>
- [5] BERTOLA, M. and CAFASSO, M. (2012). The transition between the gap probabilities from the Pearcey to the airy process—A Riemann–Hilbert approach. *Int. Math. Res. Not. IMRN* **7** 1519–1568. MR2913183 <https://doi.org/10.1093/imrn/rnr066>
- [6] BERTOLA, M. and CAFASSO, M. (2012). Riemann–Hilbert approach to multi-time processes: The Airy and the Pearcey cases. *Phys. D* **241** 2237–2245. MR2998125 <https://doi.org/10.1016/j.physd.2012.01.003>
- [7] BLEHER, P. M. and KUIJLAARS, A. B. J. (2007). Large n limit of Gaussian random matrices with external source. III. Double scaling limit. *Comm. Math. Phys.* **270** 481–517. MR2276453 <https://doi.org/10.1007/s00220-006-0159-1>
- [8] BORNEMANN, F. (2010). On the numerical evaluation of Fredholm determinants. *Math. Comp.* **79** 871–915. MR2600548 <https://doi.org/10.1090/S0025-5718-09-02280-7>
- [9] BORODIN, A. and DEIFT, P. (2002). Fredholm determinants, Jimbo–Miwa–Ueno τ -functions, and representation theory. *Comm. Pure Appl. Math.* **55** 1160–1230. MR1908746 <https://doi.org/10.1002/cpa.10042>
- [10] BOTHNER, T. and BUCKINGHAM, R. (2018). Large deformations of the Tracy–Widom distribution I: Non-oscillatory asymptotics. *Comm. Math. Phys.* **359** 223–263. MR3781450 <https://doi.org/10.1007/s00220-017-3006-7>
- [11] COIFMAN, R. R., MCINTOSH, A. and MEYER, Y. (1982). L’intégrale de Cauchy définit un opérateur borné sur L^2 pour les courbes lipschitziennes. *Ann. of Math. (2)* **116** 361–387. MR0672839 <https://doi.org/10.2307/2007065>
- [12] DEIFT, P. and ZHOU, X. (1993). A steepest descent method for oscillatory Riemann–Hilbert problems. Asymptotics for the MKdV equation. *Ann. of Math. (2)* **137** 295–368. MR1207209 <https://doi.org/10.2307/2946540>
- [13] DEIFT, P. A. (1999). *Orthogonal Polynomials and Random Matrices: A Riemann–Hilbert Approach*. Courant Lecture Notes in Mathematics **3**. New York Univ., New York. MR1677884
- [14] DIENG, M. (2005). Distribution functions for edge eigenvalues in orthogonal and symplectic ensembles: Painlevé representations. *Int. Math. Res. Not.* **37** 2263–2287. MR2181265 <https://doi.org/10.1155/IMRN.2005.2263>
- [15] EDELMAN, A. (1988). Eigenvalues and condition numbers of random matrices. *SIAM J. Matrix Anal. Appl.* **9** 543–560. MR0964668 <https://doi.org/10.1137/0609045>

- [16] FERRARI, P. L. and SPOHN, H. (2005). A determinantal formula for the GOE Tracy–Widom distribution. *J. Phys. A* **38** L557–L561. MR2165698 <https://doi.org/10.1088/0305-4470/38/33/L02>
- [17] FITZGERALD, W., TRIBE, R. and ZABORONSKI, O. Sharp asymptotics for Fredholm Paffians related to interacting particle systems and random matrices. Preprint. Available at [arXiv:1905.03754](https://arxiv.org/abs/1905.03754).
- [18] FOKAS, A. S., MUĞAN, U. and ZHOU, X. (1992). On the solvability of Painlevé I, III and V. *Inverse Probl.* **8** 757–785. MR1185598
- [19] FOKAS, A. S. and ZHOU, X. (1992). On the solvability of Painlevé II and IV. *Comm. Math. Phys.* **144** 601–622. MR1158763
- [20] FORRESTER, P. and NAGAO, T. (2007). Eigenvalue statistics of the real Ginibre ensemble. *Phys. Rev. Lett.* **99**.
- [21] FORRESTER, P. J. (2010). *Log-Gases and Random Matrices*. London Mathematical Society Monographs Series **34**. Princeton Univ. Press, Princeton, NJ. MR2641363 <https://doi.org/10.1515/9781400835416>
- [22] FORRESTER, P. J. (2015). Diffusion processes and the asymptotic bulk gap probability for the real Ginibre ensemble. *J. Phys. A* **48** 324001, 14. MR3376016 <https://doi.org/10.1088/1751-8113/48/32/324001>
- [23] GINIBRE, J. (1965). Statistical ensembles of complex, quaternion, and real matrices. *J. Math. Phys.* **6** 440–449. MR0173726 <https://doi.org/10.1063/1.1704292>
- [24] GOHBERG, I., GOLDBERG, S. and KRUPNIK, N. (2000). *Traces and Determinants of Linear Operators. Operator Theory: Advances and Applications* **116**. Birkhäuser, Basel. MR1744872 <https://doi.org/10.1007/978-3-0348-8401-3>
- [25] ITS, A. R., IZERGIN, A. G., KOREPIN, V. E. and SLAVNOV, N. A. (1990). Differential equations for quantum correlation functions. *Internat. J. Modern Phys. B* **4** 1003–1037. MR1064758 <https://doi.org/10.1142/S0217979290000504>
- [26] NIST. Digital Library of Mathematical Functions. Available at <http://dlmf.nist.gov>.
- [27] POPLAVSKYI, M., TRIBE, R. and ZABORONSKI, O. (2017). On the distribution of the largest real eigenvalue for the real Ginibre ensemble. *Ann. Appl. Probab.* **27** 1395–1413. MR3678474 <https://doi.org/10.1214/16-AAP1233>
- [28] RIDER, B. and SINCLAIR, C. D. (2014). Extremal laws for the real Ginibre ensemble. *Ann. Appl. Probab.* **24** 1621–1651. MR3211006 <https://doi.org/10.1214/13-AAP958>
- [29] SIMON, B. (2015). *Basic Complex Analysis. A Comprehensive Course in Analysis, Part 2A*. Amer. Math. Soc., Providence, RI. MR3443339 <https://doi.org/10.1090/simon/002.1>
- [30] STEIN, E. M. (1970). *Singular Integrals and Differentiability Properties of Functions*. Princeton Mathematical Series, No. 30. Princeton Univ. Press, Princeton, NJ. MR0290095
- [31] TAO, T. and VU, V. (2010). Random matrices: Universality of ESDs and the circular law. *Ann. Probab.* **38** 2023–2065. MR2722794 <https://doi.org/10.1214/10-AOP534>
- [32] TRACY, C. A. and WIDOM, H. (1994). Fredholm determinants, differential equations and matrix models. *Comm. Math. Phys.* **163** 33–72. MR1277933
- [33] TRACY, C. A. and WIDOM, H. (1996). On orthogonal and symplectic matrix ensembles. *Comm. Math. Phys.* **177** 727–754. MR1385083
- [34] ZAKHAROV, V. E. and SHABAT, A. B. (1971). Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media. *Ž. Èksper. Teoret. Fiz.* **61** 118–134. MR0406174
- [35] ZHOU, X. (1989). The Riemann–Hilbert problem and inverse scattering. *SIAM J. Math. Anal.* **20** 966–986. MR1000732 <https://doi.org/10.1137/0520065>