

LIMIT THEOREMS FOR SOME CRITICAL SUPERPROCESSES

YAN-XIA REN, RENMING SONG AND RUI ZHANG

ABSTRACT. Let $X = \{X_t, t \geq 0; \mathbb{P}_\mu\}$ be a critical superprocess starting from a finite measure μ . Under some conditions, we first prove that $\lim_{t \rightarrow \infty} t\mathbb{P}_\mu(\|X_t\| \neq 0) = \nu^{-1}\langle \phi_0, \mu \rangle$, where ϕ_0 is the eigenfunction corresponding to the first eigenvalue of the infinitesimal generator L of the mean semigroup of X , and ν is a positive constant. Then we show that, for a large class of functions f , conditioning on $\|X_t\| \neq 0$, $t^{-1}\langle f, X_t \rangle$ converges in distribution to $\langle f, \psi_0 \rangle_m W$, where W is an exponential random variable, and ψ_0 is the eigenfunction corresponding to the first eigenvalue of the dual of L . Finally, if $\langle f, \psi_0 \rangle_m = 0$, we prove that, conditioning on $\|X_t\| \neq 0$, $(t^{-1}\langle \phi_0, X_t \rangle, t^{-1/2}\langle f, X_t \rangle)$ converges in distribution to $(W, G(f)\sqrt{W})$, where $G(f) \sim \mathcal{N}(0, \sigma_f^2)$ is a normal random variable, and W and $G(f)$ are independent.

1. Introduction

1.1. Motivation. It is well known that if $\{Z_n, n \geq 0\}$ is a critical (single type) branching process with finite second moment, then

$$(1.1) \quad \lim_{n \rightarrow \infty} nP(Z_n > 0) = \frac{2}{\sigma^2}$$

Received August 17, 2015; received in final form November 16, 2015.

The research of the first author is supported by NSFC (Grant No. 11271030 and 11128101) and Specialized Research Fund for the Doctoral Program of Higher Education.

The research of the second author is supported in part by a grant from the Simons Foundation (208236).

The third author is supported by the China Scholarship Council. Corresponding author.

2010 *Mathematics Subject Classification.* Primary 60F05, 60J80. Secondary 60J25, 60J35.

and

$$(1.2) \quad \lim_{n \rightarrow \infty} P\left(\frac{1}{n}Z(n) > \frac{\sigma^2}{2}x \mid Z(n) > 0\right) = e^{-x}, \quad x \geq 0,$$

where σ^2 is the variance of the offspring distribution. The first result, due to Kolmogorov [34], says that the non-extinction rate is of order $1/n$ as $n \rightarrow \infty$. The second result, due to Yaglom [54], says that conditioned on non-extinction at time n , the total population size in generation n grows like n . For references to these results in English, one can see, for example, [21] and [25]. For probabilistic proofs of these results, see Lyons, Pemantle and Peres [40]. For continuous time critical branching processes $\{Z_t, t \geq 0\}$, Athreya and Ney [4, Theorem 3 and Lemma 2 on p. 113] proved the following limit theorem:

$$(1.3) \quad \lim_{t \rightarrow \infty} P\left(\frac{1}{t}Z(t) > \frac{\sigma^2}{2}x \mid Z(t) > 0\right) = e^{-x}, \quad x \geq 0,$$

where σ^2 is a positive constant determined by the branching rate and the variance of the offspring distribution.

For discrete time multitype critical branching processes $\{\mathbf{Z}(n), n \geq 0\}$, Athreya and Ney [4] gave two limit theorems under the finite second moment condition, see [4, Section V.5]. Let \mathbf{v} be a positive left eigenvector of the mean matrix associated with the eigenvalue 1. The first order limit theorem says that if $\mathbf{w} \cdot \mathbf{v} > 0$, then

$$(1.4) \quad \lim_{n \rightarrow \infty} P\left(\frac{\mathbf{Z}(n) \cdot \mathbf{w}}{n} > x \mid \mathbf{Z}(n) > 0\right) = e^{-x/\gamma_1}, \quad x \geq 0,$$

where $\gamma_1 := \gamma_1(\mathbf{w})$ is a positive constant. The second order limit theorem says that if $\mathbf{w} \cdot \mathbf{v} = 0$, then

$$(1.5) \quad \lim_{n \rightarrow \infty} P\left(\frac{\mathbf{Z}(n) \cdot \mathbf{w}}{\sqrt{n}} > x \mid \mathbf{Z}(n) > 0\right) = \int_x^\infty f(y) dy, \quad x \in \mathbb{R},$$

where

$$f(y) = \frac{1}{2\gamma_2} e^{-|y|/\gamma_2}, \quad y \in \mathbb{R},$$

and $\gamma_2 := \gamma_2(\mathbf{w})$ is a positive constant. The limit result (1.4) is a generalization of (1.2) from the single type case to the multitype case, and was first proved by Joffe and Spitzer [22]. The limit result (1.5) was first proved in Ney [42].

For continuous time multitype critical branching processes, Athreya and Ney [5] proved two limit theorems, similar to (1.4) and (1.5) respectively, under the finite second moment condition, see [5, Theorems 1 and 2].

For limit theorems of critical branching processes (single type or multitype) without the finite second moment condition, one can see, for instance, [20], [43], [49], [51], [52], [53] and the references therein.

Asmussen and Hering [3] discussed similar questions for critical branching Markov processes $\{Y_t, t \geq 0\}$ in a general space E under the so-called condition (M) (see [3, p. 156]) on the first moment semigroup of $\{Y_t, t \geq 0\}$. For each fixed $t \geq 0$, Y_t is a random measure on E . For any finite measure μ on E and any measurable function f on E , we use $\|\mu\|$ to denote the total mass of μ and $\langle f, \mu \rangle$ to denote the integral of f with respect to μ . In [3, Proposition 3.3 on p. 201], Asmussen and Hering discussed the finite time extinction property of branching Markov processes. [3, Theorem 3.4 on p. 202] provided the rate of non-extinction, more precisely, it was shown that

$$\lim_{t \rightarrow \infty} tP_\mu(\|Y_t\| \neq 0) = \nu^{-1} \int_E \phi_0(x)\mu(dx)$$

uniformly in μ with $\|\mu\| = n$ for any integer positive n , where ν is a positive constant and ϕ_0 is the first eigenfunction of the first moment semigroup of $\{Y_t, t \geq 0\}$. [3, Theorem 3.8 on p. 204] gave a result similar to (1.4), while [3, Theorem 3.3 on p. 297] gave a result similar to (1.5). [3, Section 4, Chapter VI] discussed the limit theorems for the critical branching Markov processes with infinite second moments.

As far as we know, not much has been done regarding limiting theorems of $\langle f, Y_t \rangle$ for critical branching Markov processes conditioned on $\|Y_t\| \neq 0$, since the book [3]. For critical superprocesses conditioned on non-extinction at time t , Evans and Perkins [19] obtained results similar to (1.1) and (1.4) when $\varphi(x, z) = z^2$, $\beta(x) \equiv 1$ and the spatial process satisfies some ergodicity conditions. [19] did not consider central limit theorem type results. We note in passing that [19] also obtained results similar to (1.4) conditioned on remote survival. See [7] for similar results for multitype Dawson–Watanabe processes conditioned on remote survival.

The main purpose of this paper is to establish limit theorems similar to (1.1), (1.4) and a central limit type theorem for critical superprocesses, under the finite second moment condition and other very general, easy to check conditions. Here is a summary of our main results. Let $X = \{X_t, t \geq 0; \mathbb{P}_\mu\}$ be a critical superprocess starting from a finite measure μ . Under some conditions to be specified later, we first prove that $\lim_{t \rightarrow \infty} t\mathbb{P}_\mu(\|X_t\| \neq 0) = \nu^{-1}\langle \phi_0, \mu \rangle$, where ϕ_0 is the eigenfunction corresponding to the first eigenvalue of the infinitesimal generator L of the mean semigroup of X , and ν is a positive constant. Then we show that, for a large class of functions f , conditioning on $\|X_t\| \neq 0$, $t^{-1}\langle f, X_t \rangle$ converges in distribution to $\langle f, \psi_0 \rangle_m W$, where W is an exponential random variable, and ψ_0 is the eigenfunction corresponding to the first eigenvalue of the dual of L . Finally, if $\langle f, \psi_0 \rangle_m = 0$, we prove that, conditioning on $\|X_t\| \neq 0$, $(t^{-1}\langle \phi_0, X_t \rangle, t^{-1/2}\langle f, X_t \rangle)$ converges in distribution to $(W, G(f)\sqrt{W})$, where $G(f) \sim \mathcal{N}(0, \sigma_f^2)$ is a normal random variable, and W and $G(f)$ are independent.

In our recent papers [44], [46], we established some spatial central limit theorems for supercritical superprocesses. See also [1], [41], [45], [47] for related results for supercritical branching Markov processes and supercritical superprocesses. Our original motivation for the present paper is to establish spatial central limit theorems for critical superprocesses. One of the main tools of the papers above is the analytical and spectral properties of the Feynman–Kac semigroup of the spatial process, which also play an important role in this paper. We will assume that the dual, with respect to a certain measure, of the semigroup of the spatial process is a Markov semigroup. See the next subsection for details.

For branching Markov processes, there is a clear particle picture. This particle structure was used essentially in proving the central limit theorems for supercritical branching Markov processes in [1], [45], [47]. For superprocesses, the particle picture is less clear. In this case, the backbone decomposition or the excursion measures are frequently used to describe the ‘infinitesimal particles’. [41], [44] used the backbone decomposition to establish central limit theorems for supercritical super-OU processes, while [46] used the excursion measures of superprocesses to prove central limit theorems for general supercritical superprocesses. In this paper, we will also use the excursion measure to prove our central limit theorem. Up to now, there is no known backbone decomposition for critical superprocesses conditioned on survival up to t yet.

1.2. Superprocesses and assumptions. In this subsection, we describe the superprocesses we are going to work with and formulate our assumptions. Since one of the main tools of this paper is the analytic properties of the semigroup of the spatial process, we will need to assume that the semigroup of the spatial process has a dual with respect to a certain measure m and the dual semigroup is Markovian.

Suppose that E is a locally compact separable metric space and that m is a σ -finite Borel measure on E with full support. Suppose that ∂ is a separate point not contained in E . ∂ will be interpreted as the cemetery point. We will use E_∂ to denote $E \cup \{\partial\}$. Every function f on E is automatically extended to E_∂ by setting $f(\partial) = 0$. We will assume that $\xi = \{\xi_t, \Pi_x\}$ is a Hunt process on E and $\zeta := \inf\{t > 0 : \xi_t = \partial\}$ is the lifetime of ξ . We will use $\{P_t : t \geq 0\}$ to denote the semigroup of ξ . We will use $\mathcal{B}(E)$ ($\mathcal{B}^+(E)$) to denote the set of (non-negative) Borel measurable functions on E , and use $\mathcal{B}_b(E)$ ($\mathcal{B}_b^+(E)$) to denote the set of (non-negative) bounded Borel measurable functions on E .

The superprocess $X = \{X_t : t \geq 0\}$ we are going to work with is determined by three parameters: a spatial motion $\xi = \{\xi_t, \Pi_x\}$ on E which is a Hunt process, a branching rate function $\beta(x)$ on E which is a non-negative bounded

measurable function and a branching mechanism φ of the form

$$(1.6) \quad \begin{aligned} \varphi(x, z) &= -a(x)z + b(x)z^2 \\ &+ \int_{(0, +\infty)} (e^{-zy} - 1 + zy)n(x, dy), \quad x \in E, z \geq 0, \end{aligned}$$

where $a \in \mathcal{B}_b(E)$, $b \in \mathcal{B}_b^+(E)$ and n is a kernel from E to $(0, \infty)$ satisfying

$$(1.7) \quad \sup_{x \in E} \int_{(0, +\infty)} y^2 n(x, dy) < \infty.$$

The assumption (1.7) is the counterpart of the second moment condition in [25]. In the multitype continuous time branching process case, one does not need to take the supremum and explicitly assume that β is bounded, since the state space of the spatial process, that is, the type space, is finite. Under this assumption, the superprocess X has finite second moments (see (2.13) below). In our paper, we will not consider the special case that $\beta(\cdot)(b(\cdot) + n(\cdot, (0, \infty))) = 0$, a.e.-m.

Let $\mathcal{M}_F(E)$ be the space of finite measures on E , equipped with topology of weak convergence. The superprocess X is a Markov process taking values in $\mathcal{M}_F(E)$. The existence of such superprocesses is well-known, see, for instance, [13], [15] or [39]. For any $\mu \in \mathcal{M}_F(E)$, we denote the law of X with initial configuration μ by \mathbb{P}_μ . As usual, $\langle f, \mu \rangle := \int_E f(x)\mu(dx)$ and $\|\mu\| := \langle 1, \mu \rangle$. Throughout this paper, a real-valued function $u(t, x)$ on $[0, \infty) \times E_\partial$ is said to be locally bounded if for any $t > 0$, $\sup_{s \in [0, t], x \in E_\partial} |u(s, x)| < \infty$. According to [39, Theorem 5.12], there is a Hunt process $X = \{\Omega, \mathcal{G}, \mathcal{G}_t, X_t, \mathbb{P}_\mu\}$ taking values in $\mathcal{M}_F(E)$, such that, for every $f \in \mathcal{B}_b^+(E)$ and $\mu \in \mathcal{M}_F(E)$,

$$(1.8) \quad -\log \mathbb{P}_\mu(e^{-\langle f, X_t \rangle}) = \langle u_f(t, \cdot), \mu \rangle,$$

where $u_f(t, x)$ is the unique locally bounded non-negative solution to the equation

$$(1.9) \quad u_f(t, x) + \Pi_x \int_0^t \Psi(\xi_s, u_f(t - s, \xi_s)) ds = \Pi_x f(\xi_t), \quad x \in E_\partial,$$

where $\Psi(x, z) = \beta(x)\varphi(x, z)$, $x \in E$ and $z \geq 0$, while $\Psi(\partial, z) = 0$, $z \geq 0$. Since $f(\partial) = 0$, we have $u_f(t, \partial) = 0$ for any $t \geq 0$. In this paper, the superprocess we deal with is always this Hunt realization. The function β is usually called the branching rate, and φ is called the branching mechanism. For more general superprocesses, β can be a measure on E . For the process X in this paper, β can be absorbed to φ . That is to say, we could have, without loss of generality, supposed that $\beta \equiv 1$. To be consistent with the formulations of our previous papers [44], [45], [46], [47], we keep β as a function.

Define

$$(1.10) \quad \alpha(x) := \beta(x)a(x) \quad \text{and} \quad A(x) := \beta(x) \left(2b(x) + \int_0^\infty y^2 n(x, dy) \right).$$

Then, by our assumptions, $\alpha(x) \in \mathcal{B}_b(E)$ and $A(x) \in \mathcal{B}_b^+(E)$. Thus there exists $K > 0$ such that

$$(1.11) \quad \sup_{x \in E} (|\alpha(x)| + A(x)) \leq K.$$

For any $f \in \mathcal{B}_b(E)$ and $(t, x) \in (0, \infty) \times E$, define

$$(1.12) \quad T_t f(x) := \Pi_x [e^{\int_0^t \alpha(\xi_s) ds} f(\xi_t)].$$

It is well known that $T_t f(x) = \mathbb{P}_{\delta_x} \langle f, X_t \rangle$ for every $x \in E$.

Our standing assumption on ξ is that there exists a family of continuous strictly positive functions $\{p(t, x, y) : t > 0\}$ on $E \times E$ such that, for any $t > 0$ and nonnegative function f on E ,

$$P_t f(x) = \int_E p(t, x, y) f(y) m(dy).$$

Define

$$a_t(x) := \int_E p(t, x, y)^2 m(dy), \quad \widehat{a}_t(x) := \int_E p(t, y, x)^2 m(dy).$$

In this paper, we assume the following assumption.

- ASSUMPTION 1.1. (i) For any $t > 0$, $\int_E p(t, x, y) m(dx) \leq 1$.
 (ii) For any $t > 0$, we have

$$(1.13) \quad \begin{aligned} e_t &:= \int_E a_t(x) m(dx) = \int_E \widehat{a}_t(x) m(dx) \\ &= \int_E \int_E p(t, x, y)^2 m(dy) m(dx) < \infty. \end{aligned}$$

Moreover, the functions $x \rightarrow a_t(x)$ and $x \rightarrow \widehat{a}_t(x)$ are continuous on E .

Note that, in Assumption 1.1(i), the integration is with respect to the first space variable. It implies that the dual semigroup $\{\widehat{P}_t : t \geq 0\}$ of $\{P_t : t \geq 0\}$ with respect to m defined by

$$\widehat{P}_t f(x) = \int_E p(t, y, x) f(y) m(dy)$$

is Markovian. Assumption 1.1(ii) is a pretty weak L^2 condition and it allows us to apply results on operator semigroups in Hilbert spaces.

By Hölder’s inequality, we have

$$(1.14) \quad p(t + s, x, y) = \int_E p(t, x, z) p(s, z, y) m(dz) \leq (a_t(x))^{1/2} (\widehat{a}_s(y))^{1/2}.$$

It is well known and easy to check that, $\{P_t : t \geq 0\}$ and $\{\widehat{P}_t : t \geq 0\}$ are strongly continuous contraction semigroups on $L^2(E, m)$, see [47] for a proof. Recall that $\{P_t : t \geq 0\}$ is a strongly continuous contraction semigroup on $L^2(E, m)$ means that, for any $f \in L^2(E, m)$, $\lim_{t \rightarrow 0} \|P_t f - f\|_2 = 0$ and $\|P_t f\|_2 \leq \|f\|_2$ for all $t \geq 0$. We will use $\langle \cdot, \cdot \rangle_m$ to denote inner product

in $L^2(E, m)$. Since $p(t, x, y)$ is continuous in (x, y) , by (1.14) and Assumption 1.1(ii), using the dominated convergence theorem, we get that, for any $f \in L^2(E, m)$, $P_t f$ and $\widehat{P}_t f$ are continuous.

It follows from Assumption 1.1(ii) that, for each $t > 0$, P_t and $\{\widehat{P}_t\}$ are compact operators on $L^2(E, m)$. Let \widetilde{L} and $\widehat{\widetilde{L}}$ be the infinitesimal generators of the semigroups $\{P_t\}$ and $\{\widehat{P}_t\}$ in $L^2(E, m)$ respectively. Define $\widetilde{\lambda}_0 := \sup \Re(\sigma(\widetilde{L})) = \sup \Re(\sigma(\widehat{\widetilde{L}}))$. By Jentzsch's theorem (Theorem V.6.6 on p. 337 of [48]), $\widetilde{\lambda}_0$ is an eigenvalue of multiplicity 1 for both \widetilde{L} and $\widehat{\widetilde{L}}$, and that an eigenfunction $\widetilde{\phi}_0$ of \widetilde{L} corresponding to $\widetilde{\lambda}_0$ can be chosen to be strictly positive m -almost everywhere with $\|\widetilde{\phi}_0\|_2 = 1$ and an eigenfunction $\widetilde{\psi}_0$ of $\widehat{\widetilde{L}}$ corresponding to $\widetilde{\lambda}_0$ can be chosen to be strictly positive m -almost everywhere with $\langle \widetilde{\phi}_0, \widetilde{\psi}_0 \rangle_m = 1$. Thus for m -almost every $x \in E$,

$$e^{\widetilde{\lambda}_0 t} \widetilde{\phi}_0(x) = P_1 \widetilde{\phi}_0(x), \quad e^{\widetilde{\lambda}_0 t} \widetilde{\psi}_0(x) = \widehat{P}_1 \widetilde{\psi}_0(x).$$

Hence, $\widetilde{\phi}_0$ and $\widetilde{\psi}_0$ can be chosen to be continuous and strictly positive everywhere on E .

Our second assumption is the following.

ASSUMPTION 1.2. (i) $\widetilde{\phi}_0$ is bounded.

(ii) The semigroup $\{P_t : t \geq 0\}$ is intrinsically ultracontractive, that is, there exists $c_t > 0$ such that

$$(1.15) \quad p(t, x, y) \leq c_t \widetilde{\phi}_0(x) \widetilde{\psi}_0(y).$$

Assumption 1.2 is a pretty strong assumption on the semigroup $\{P_t : t \geq 0\}$. However, this assumption is satisfied in a lot of cases. In Section 1.4, we will give many examples where Assumptions 1.1 and 1.2 are satisfied. Here we only give one very special example. If E consists of finitely many points and $\xi = \{\xi_t : t \geq 0\}$ is a conservative irreducible Markov process on E , then ξ satisfies Assumptions 1.1 and 1.2 for some finite measure m on E with full support. So, as special cases, our results give the analogs of the results of Athreya and Ney [5] for critical super-Markov chains.

We will prove in Lemma 2.1 that there exists a function $q(t, x, y)$ on $(0, \infty) \times E \times E$ which is continuous in (x, y) for each $t > 0$ such that

$$(1.16) \quad e^{-Kt} p(t, x, y) \leq q(t, x, y) \leq e^{Kt} p(t, x, y), \quad (t, x, y) \in (0, \infty) \times E \times E$$

and that for any bounded Borel function f and any $(t, x) \in (0, \infty) \times E$,

$$T_t f(x) = \int_E q(t, x, y) f(y) m(dy).$$

It follows immediately that

$$(1.17) \quad \|T_t f\|_2 \leq e^{Kt} \|P_t f\|_2 \leq e^{Kt} \|f\|_2.$$

In [47], we have proved that $\{T_t : t \geq 0\}$ is a strongly continuous semigroup on $L^2(E, m)$. Let $\{\widehat{T}_t : t > 0\}$ be the adjoint operators on $L^2(E, m)$ of $\{T_t : t > 0\}$, that is, for $f, g \in L^2(E, m)$,

$$\int_E f(x)T_t g(x)m(dx) = \int_E g(x)\widehat{T}_t f(x)m(dx)$$

and

$$\widehat{T}_t f(x) = \int_E q(t, y, x)f(y)m(dy).$$

We have proved in [47] that $\{\widehat{T}_t : t \geq 0\}$ is also a strongly continuous semigroup on $L^2(E, m)$. We claim that, for all $t > 0$ and $f \in L^2(E, m)$, $T_t f$ and $\widehat{T}_t f$ are continuous. In fact, since $q(t, x, y)$ is continuous in (x, y) , by (1.14), (1.16) and Assumption 1.1(ii), using the dominated convergence theorem, we get that, for any $f \in L^2(E, m)$, $T_t f$ and $\widehat{T}_t f$ are continuous.

By Assumption 1.1(ii) and (1.16), we get that

$$\int_E \int_E q^2(t, x, y)m(x)m(dy) \leq e^{2Kt} \int_E \int_E p^2(t, x, y)m(x)m(dy) < \infty.$$

Thus, for each $t > 0$, T_t and $\{\widehat{T}_t\}$ are compact operators on $L^2(E, m)$. Let L and \widehat{L} be the infinitesimal generators of the semigroups $\{T_t\}$ and $\{\widehat{T}_t\}$ in $L^2(E, m)$ respectively. Define $\lambda_0 := \sup \Re(\sigma(L)) = \sup \Re(\sigma(\widehat{L}))$. By Jentzsch's theorem, λ_0 is an eigenvalue of multiplicity 1 for both L and \widehat{L} , and that an eigenfunction ϕ_0 of L corresponding to λ_0 can be chosen to be strictly positive m -almost everywhere with $\|\phi_0\|_2 = 1$ and an eigenfunction ψ_0 of \widehat{L} corresponding to λ_0 can be chosen to be strictly positive m -almost everywhere with $\langle \phi_0, \psi_0 \rangle_m = 1$. Thus for m -almost every $x \in E$,

$$e^{\lambda_0 t} \phi_0(x) = T_t \phi_0(x), \quad e^{\lambda_0 t} \psi_0(x) = \widehat{T}_t \psi_0(x).$$

Hence, ψ_0 and ϕ_0 can be chosen to be continuous and strictly positive everywhere on E .

Using Assumption 1.2, the boundedness of α and an argument similar to that used in the proof of [12, Theorem 3.4], one can show the following:

- (i) ϕ_0 is bounded.
- (ii) The semigroup $\{T_t : t \geq 0\}$ is intrinsically ultracontractive, that is, there exists $c_t > 0$ such that

$$(1.18) \quad q(t, x, y) \leq c_t \phi_0(x) \psi_0(y).$$

The condition (M) on [3, p. 156] is a condition similar in spirit to the intrinsic ultracontractivity of $\{T_t : t \geq 0\}$. This condition is not very easy to check. Essentially the only examples given in [3] satisfying this condition are branching diffusion processes in bounded smooth domains. Our Assumption 1.2 is in terms of the intrinsic ultracontractivity of $\{P_t : t \geq 0\}$. Intrinsic ultracontractivity has been studied intensively in the last 30 years and there are many

results on the intrinsic ultracontractivity of semigroups. Using these results, we will give in Section 1.4 many examples satisfying Assumption 1.2.

Let λ_∞ be the L^∞ -growth bound of the semigroup T_t , that is,

$$\lambda_\infty := \lim_{t \rightarrow \infty} \frac{1}{t} \log \|T_t\|_{\infty, \infty}.$$

It is easy to see that $\lambda_0 \leq \lambda_\infty$. Note that λ_0 gives the rate of local growth when it is positive, and implies local extinction otherwise. While if $\lambda_\infty \neq 0$, then in some sense, the exponential growth/decay rate of $\langle 1, X_t \rangle$, the total mass of X_t , is λ_∞ , see [18]. According to [28, Theorem 2.7], under Assumptions 1.1 and 1.2, there exist constants $\gamma > 0$ and $c > 0$ such that, for any $(t, x, y) \in (1, \infty) \times E \times E$, we have

$$(1.19) \quad |e^{-\lambda_0 t} q(t, x, y) - \phi_0(x)\psi_0(y)| \leq ce^{-\gamma t} \phi_0(x)\psi_0(y).$$

Hence for $(t, x, y) \in (1, \infty) \times E \times E$, we have

$$(1.20) \quad e^{-\lambda_0 t} q(t, x, y) \geq (1 - ce^{-\gamma t}) \phi_0(x)\psi_0(y).$$

Since $q(t, x, \cdot) \in L^1(E, m)$, we have $\psi_0 \in L^1(E, m)$. Therefore, by (1.19), for $t > 1$, $\|T_t\|_{\infty, \infty} \leq (1 + c)\|\phi_0\|_\infty \langle 1, \psi_0 \rangle_m e^{\lambda_0 t}$, which implies $\lambda_0 = \lambda_\infty$.

The main interest of this paper is on critical superprocesses, so we assume the following.

ASSUMPTION 1.3. $\lambda_0 = 0$.

Define $q_t(x) := \mathbb{P}_{\delta_x}(\|X_t\| = 0)$. Note that, since $\mathbb{P}_{\delta_x}\|X_t\| = T_t 1(x) > 0$, we have $\mathbb{P}_{\delta_x}(\|X_t\| = 0) < 1$. In this paper, we also assume the following.

ASSUMPTION 1.4. For any $t > 0$ and $x \in E$, $q_t(x) \in (0, 1)$. And, there exists $t_0 > 0$ such that,

$$(1.21) \quad \inf_{x \in E} q_{t_0}(x) > 0.$$

In Section 2.2, we will give a sufficient condition (in term of the function Ψ) for Assumption 1.4. In Lemma 3.3, we will show that, under our assumptions, $\lim_{t \rightarrow \infty} q_t(x) = 1$, uniformly in $x \in E$.

1.3. Main results. In this subsection, we will state our main results. In the following, we use the notation

$$\mathbb{P}_{t, \mu}(\cdot) := \mathbb{P}_\mu(\cdot | \|X_t\| \neq 0).$$

Recall that the process X is defined on (Ω, \mathcal{G}) . Suppose that, for each $t > 0$, Y_t is a measurable map from (Ω, \mathcal{G}) to a Polish space S and that Z is an S -valued random variable on a probability space $(\tilde{\Omega}, \tilde{\mathcal{G}}, P)$, we write

$$Y_t |_{\mathbb{P}_{t, \mu}} \xrightarrow{d} Z,$$

if $\lim_{t \rightarrow \infty} \mathbb{P}_{t, \mu}[f(Y_t)] = P[f(Z)]$ for all bounded continuous real-valued functions f on S .

Define

$$(1.22) \quad \nu := \frac{1}{2} \langle A(\phi_0)^2, \psi_0 \rangle_m.$$

It is easy to see that $0 < \nu < \infty$. Define

$$\mathcal{C}_p := \{f \in \mathcal{B}(E) : \langle |f|^p, \psi_0 \rangle_m < \infty\}$$

and $\mathcal{C}_p^+ := \mathcal{C}_p \cap \mathcal{B}^+(E)$. By (1.18) and the fact that $q(t, x, y)$ is continuous, using the dominated convergence theorem, we get that, for $f \in \mathcal{C}_1$, $T_t f(x)$ is continuous. Since $\psi_0 \in L^1(E, m)$, $\mathcal{B}_b(E) \subset \mathcal{C}_p$. Moreover, by Hölder’s inequality, we get $\mathcal{C}_2 \subset \mathcal{C}_1$.

THEOREM 1.5. For any non-zero $\mu \in \mathcal{M}_F(E)$,

$$(1.23) \quad \lim_{t \rightarrow \infty} t \mathbb{P}_\mu (\|X_t\| \neq 0) = \nu^{-1} \langle \phi_0, \mu \rangle.$$

Furthermore, the convergence above is uniform in μ with $\mu(E) \leq M$, where $M > 0$ is any constant.

THEOREM 1.6. If $f \in \mathcal{C}_2$ then, for any non-zero $\mu \in \mathcal{M}_F(E)$, we have

$$(1.24) \quad t^{-1} \langle f, X_t \rangle |_{\mathbb{P}_{t,\mu}} \xrightarrow{d} \langle f, \psi_0 \rangle_m W,$$

where W is an exponential random variable with parameter $1/\nu$. In particular, we have

$$(1.25) \quad t^{-1} \langle \phi_0, X_t \rangle |_{\mathbb{P}_{t,\mu}} \xrightarrow{d} W.$$

REMARK 1.7. (1) The distributional limit $\langle f, \psi_0 \rangle_m W$ in Theorem 1.6 does not depend on the starting measure μ .

(2) Since $1 \in \mathcal{B}_b(E) \subset \mathcal{C}_2$, thus the limit result above implies that

$$t^{-1} \langle 1, X_t \rangle |_{\mathbb{P}_{t,\mu}} \xrightarrow{d} \langle 1, \psi_0 \rangle_m W,$$

which says that, conditioned on no-extinction at time t , the growth rate of the total mass $\langle 1, X_t \rangle$ is t as $t \rightarrow \infty$.

It is well known, see, for instance, [24, Theorem A2.3] that the collection of Radon measures on E equipped with the vague topology forms a Polish space. Let $\rho(\cdot, \cdot)$ be a metric on the space of Radon measures on E compatible with the vague topology. Let l be the finite (deterministic) measure on E defined by $l(dx) = \psi_0(x)m(dx)$.

COROLLARY 1.8. For any $f \in \mathcal{C}_2$ and non-zero $\mu \in \mathcal{M}_F(E)$, it holds that, as $t \rightarrow \infty$,

$$(1.26) \quad \frac{\langle f, X_t \rangle}{\langle \phi_0, X_t \rangle} \Big|_{\mathbb{P}_{t,\mu}} \xrightarrow{d} \langle f, \psi_0 \rangle_m.$$

Moreover, for any non-zero $\mu \in \mathcal{M}_F(E)$ and $\varepsilon > 0$,

$$\lim_{t \rightarrow \infty} \mathbb{P}_{t,\mu} \left(\rho \left(\frac{X_t}{\langle \phi_0, X_t \rangle}, l \right) \geq \varepsilon \right) = 0.$$

The above corollary can be thought of as a “weak” law of large numbers. Thus it is natural to consider a corresponding central limit type theorem. For this, we need to find constants a_t such that

$$a_t \left(\frac{X_t}{\langle \phi_0, X_t \rangle} - l \right) \Big|_{\mathbb{P}_{t,\mu}} \xrightarrow{d} Y$$

for some nontrivial finite random measure Y . According to [24, Theorem 16.16], it suffices to show that for each continuous function f with compact support in E ,

$$a_t \left(\frac{\langle f, X_t \rangle}{\langle \phi_0, X_t \rangle} - \langle f, \psi_0 \rangle_m \right) \Big|_{\mathbb{P}_{t,\mu}} \xrightarrow{d} \langle f, Y \rangle.$$

This is equivalent to finding a_t such that

$$a_t \frac{\langle \tilde{f}, X_t \rangle}{\langle \phi_0, X_t \rangle} \Big|_{\mathbb{P}_{t,\mu}} \xrightarrow{d} \langle f, Y \rangle,$$

where $\tilde{f} = f - \langle f, \psi_0 \rangle_m \phi_0$ satisfies $\langle \tilde{f}, \psi_0 \rangle_m = 0$. This is the reason that we consider only functions $f \in \mathcal{C}_2$ and $\langle f, \psi_0 \rangle_m = 0$ in the next theorem.

Define

$$(1.27) \quad \sigma_f^2 = \int_0^\infty \langle A(T_s f)^2, \psi_0 \rangle_m ds.$$

THEOREM 1.9. *Suppose that $f \in \mathcal{C}_2$ and $\langle f, \psi_0 \rangle_m = 0$, then we have $\sigma_f^2 < \infty$ and, for any non-zero $\mu \in \mathcal{M}_F(E)$,*

$$(1.28) \quad (t^{-1} \langle \phi_0, X_t \rangle, t^{-1/2} \langle f, X_t \rangle) \Big|_{\mathbb{P}_{t,\mu}} \xrightarrow{d} (W, G(f) \sqrt{W}),$$

where $G(f) \sim \mathcal{N}(0, \sigma_f^2)$ is a normal random variable and W is the random variable defined in Theorem 1.6. Moreover, W and $G(f)$ are independent.

Combining Theorems 1.6 and 1.9, we see that, when $\sigma_f^2 > 0$, the density of $G(f) \sqrt{W}$ is

$$d(x) = \frac{1}{\sqrt{2\nu\sigma_f^2}} \exp \left\{ -\frac{2|x|}{\sqrt{2\nu\sigma_f^2}} \right\}, \quad x \in \mathbb{R}.$$

As a consequence of Theorem 1.9, we immediately get the following result, which can be thought of as some sort of central limit theorem.

COROLLARY 1.10. *Suppose that $f \in \mathcal{C}_2$ and $\langle f, \psi_0 \rangle_m = 0$, then we have $\sigma_f^2 < \infty$ and, for any non-zero $\mu \in \mathcal{M}_F(E)$,*

$$(1.29) \quad \left(t^{-1} \langle \phi_0, X_t \rangle, \frac{\langle f, X_t \rangle}{\sqrt{\langle \phi_0, X_t \rangle}} \right) \Big|_{\mathbb{P}_{t,\mu}} \xrightarrow{d} (W, G(f)),$$

where $G(f) \sim \mathcal{N}(0, \sigma_f^2)$ is a normal random variable and W is the random variable defined in Theorem 1.6. Moreover, W and $G(f)$ are independent.

REMARK 1.11. Suppose that m is a probability measure, the spatial motion ξ is conservative (that is, $P_t 1 = 1$), and that the branching mechanism is spatial-independent with

$$(1.30) \quad \Psi(z) = bz^2 + \int_0^\infty (e^{-zy} - 1 + zy)n(dy),$$

where $b \geq 0$ and $\int_0^\infty z^2 n(dz) < \infty$. Then $T_t = P_t$, $\lambda_0 = 0$ and $\phi_0(x) = 1$. Thus, Assumption 1.3 is satisfied. The process $\{\|X_t\|, t \geq 0\}$ is a continuous state branching process with branching mechanism Ψ . We assume that Ψ satisfies the Grey condition:

$$(1.31) \quad \int_0^\infty \frac{1}{\Psi(z)} dz < \infty.$$

Then, for any $\mu \in \mathcal{M}_F(E)$,

$$\lim_{t \rightarrow \infty} t \mathbb{P}_\mu (\|X_t\| \neq 0) = 2A^{-1} \|\mu\|,$$

where $A = 2b + \int_0^\infty y^2 n(dy)$, and

$$t^{-1} \|X_t\| \Big|_{\mathbb{P}_{t,\mu}} \xrightarrow{d} W,$$

where W is an exponential random variable with parameter $2A^{-1}$. The proofs can be found in [36], [37]. It is easy to check that, under the assumptions above, Assumption 1.4 is satisfied, see the end of Section 2.2.

Suppose that the spatial motion ξ satisfies Assumption 1.1 and

ASSUMPTION 1.2'. There exists $t_0 > 0$ such that $a_{t_0}, \widehat{a}_{t_0} \in L^2(E, m)$.

Then using an argument similar to that in [47, Lemma 2.6(1)], we can get that, for $f \in L^2(E, m) \cap L^4(E, m)$,

$$\lim_{t \rightarrow \infty} \mathbb{V}ar_{\delta_x} \langle f, X_t \rangle = \sigma_f^2 < \infty.$$

Thus, using the same arguments as in the proofs of Theorem 1.6 and Theorem 1.9 below, we can get that Theorem 1.6 and Theorem 1.9 are also valid in this case for $f \in L^2(E, m) \cap L^4(E, m)$ and $\mu \in \mathcal{M}_F(E)$ with compact support. We will not give the detailed proof in this case.

Note that in this case we do not need Assumption 1.2. One can check that super inward Ornstein–Uhlenbeck processes satisfy Assumption 1.1 and Assumption 1.2', see [11, Examples 4.1]. Thus, Theorem 1.6 and Theorem 1.9

hold for super inward Ornstein–Uhlenbeck processes with spatial-independent branching mechanism Ψ given by (1.30).

1.4. Examples. In this subsection, we present a list of examples which satisfy Assumptions 1.1 and 1.2. For simplicity, we will not try to give the weakest possible conditions. The first six are examples where the processes are symmetric with respect to some measure.

EXAMPLE 1.12. Suppose that E is a connected open subset of \mathbb{R}^d with finite Lebesgue measure and that m denotes the Lebesgue measure on E . Let ξ be the subprocess in E of a diffusion process in \mathbb{R}^d corresponding to a uniformly elliptic divergence form second order differential operator. Then it is well known that ξ has a transition density $p(t, x, y)$ which is a strictly positive, continuous and symmetric function of (x, y) for any $t > 0$ and that there exists $c > 0$ such that

$$p(t, x, y) \leq ct^{d/2}, \quad (t, x, y) \in (0, \infty) \times E \times E.$$

Thus Assumption 1.1 is trivially satisfied. If E is a bounded Lipschitz connected open set, then it follows from [12] that the semigroup $\{P_t : t \geq 0\}$ of ξ is intrinsic ultracontractive and that the eigenfunction $\tilde{\phi}_0$ corresponding to the largest eigenvalue of the generator of $\{P_t : t \geq 0\}$ is bounded. Thus Assumption 1.2 is satisfied. Under much weaker regularity assumptions on E , Assumptions 1.1 and 1.2 are still satisfied. For some of these weaker regularity assumptions, one can see [6] and the references therein.

EXAMPLE 1.13. Suppose that E is the closure of a bounded connected C^2 open set in \mathbb{R}^d and that m denotes the Lebesgue measure on E . Let ξ be the reflecting Brownian motion in E . Then ξ has a transition density $p(t, x, y)$ which is a strictly positive, continuous and symmetric function of (x, y) for any $t > 0$ and that there exists $c > 0$ such that

$$p(t, x, y) \leq ct^{d/2}, \quad (t, x, y) \in (0, \infty) \times E \times E.$$

The largest eigenvalue of the generator of the semigroup $\{P_t : t \geq 0\}$ of ξ is $\tilde{\lambda}_0 = 0$ and the corresponding eigenfunction $\tilde{\phi}_0$ is a positive constant. Thus, Assumptions 1.1 and 1.2 are trivially satisfied.

EXAMPLE 1.14. Suppose that E is an open subset of \mathbb{R}^d with finite Lebesgue measure and that m denotes the Lebesgue measure on E . Let ξ be the subprocesses in E of any of the subordinate Brownian motions studied in [32], [33]. Then it is known (see [9], [10]) that ξ has a transition density $p(t, x, y)$ which is a strictly positive, continuous, bounded, symmetric function of (x, y) for any $t > 0$. Thus Assumption 1.1 is trivially satisfied. It follows from [29] that the semigroup $\{P_t : t \geq 0\}$ of ξ is intrinsic ultracontractive and that the eigenfunction $\tilde{\phi}_0$ corresponding to the largest eigenvalue of the generator of $\{P_t : t \geq 0\}$ is bounded. Thus, Assumption 1.2 is also satisfied.

EXAMPLE 1.15. Suppose $a > 2$ is a constant. Assume that $E = \mathbb{R}^d$ and m is the Lebesgue measure on \mathbb{R}^d . Let ξ be a Markov process on \mathbb{R}^d corresponding to the infinitesimal generator $\Delta - |x|^a$. Let $p(t, x, y)$ denote the transition density of ξ with respect to the Lebesgue measure on \mathbb{R}^d . It follows from [12, Theorem 6.1] and its proof that, for any $t > 0$, there exists $c_t > 0$ such that

$$p(t, x, y) \leq c_t \exp\left(-\frac{2}{2+a}|x|^{1+a/2}\right) \exp\left(-\frac{2}{2+a}|y|^{1+a/2}\right), \quad x, y \in \mathbb{R}^d,$$

that the eigenfunction $\tilde{\phi}_0$ corresponding to the largest eigenvalue of the generator of $\{P_t : t \geq 0\}$ of ξ is bounded and that $\{P_t : t \geq 0\}$ is intrinsically ultracontractive. Thus, Assumptions 1.1 and 1.2 are satisfied.

EXAMPLE 1.16. Assume that $E = \mathbb{R}^d$ and m is the Lebesgue measure on \mathbb{R}^d . Suppose that V is a nonnegative and locally bounded function on \mathbb{R}^d such that there exist $R > 0$ and $M \geq 1$ such that for all $|x| > R$,

$$M^{-1}(1 + V(x)) \leq V(y) \leq M(1 + V(x)), \quad y \in B(x, 1),$$

and that

$$\lim_{|x| \rightarrow \infty} \frac{V(x)}{\log |x|} = \infty.$$

Suppose $\beta \in (0, 2)$ is a constant. Let ξ be a Markov process on \mathbb{R}^d corresponding to the infinitesimal generator $-(-\Delta)^{\beta/2} - V(x)$. Let $p(t, x, y)$ denote the transition density of ξ with respect to the Lebesgue measure on \mathbb{R}^d . It follows from [23, Corollaries 3 and 4] that, for any $t > 0$, there exists $c_t > 0$ such that

$$p(t, x, y) \leq c_t \frac{1}{(1 + V(x))(1 + |x|)^{d+\beta}} \frac{1}{(1 + V(y))(1 + |y|)^{d+\beta}}, \quad x, y \in \mathbb{R}^d,$$

that the eigenfunction $\tilde{\phi}_0$ corresponding to the largest eigenvalue of the generator of $\{P_t : t \geq 0\}$ of ξ is bounded and that $\{P_t : t \geq 0\}$ is intrinsically ultracontractive. Thus, Assumptions 1.1 and 1.2 are satisfied.

EXAMPLE 1.17. Assume that $E = \mathbb{R}^d$ and m is the Lebesgue measure on \mathbb{R}^d . A nondecreasing function $L : [0, \infty) \rightarrow [0, \infty)$ is said to be in the class \mathbf{L} if $\lim_{t \rightarrow \infty} L(t) = \infty$ and there exists $c > 1$ such that

$$L(t + 1) \leq c(1 + L(t)), \quad t \geq 0.$$

Suppose that V is a nonnegative function on \mathbb{R}^d such that

$$\lim_{|x| \rightarrow \infty} \frac{V(x)}{|x|} = \infty$$

and that there exists a function $L \in \mathbf{L}$ such that there exists $C > 0$ such that

$$L(|x|) \leq V(x) \leq C(1 + L(|x|)), \quad x \in \mathbb{R}^d.$$

Suppose that $r > 0$ and $\beta \in (0, 2)$ are constants. Let ξ be a Markov process on \mathbb{R}^d corresponding to the infinitesimal generator $r - (-\Delta + r^{2/\beta})^{\beta/2} - V(x)$.

Let $p(t, x, y)$ denote the transition density of ξ with respect to the Lebesgue measure on \mathbb{R}^d . It follows from [35, Theorem 1.6] that, for any $t > 0$, there exists $c_t > 0$ such that for all $x, y \in \mathbb{R}^d$,

$$p(t, x, y) \leq c_t \frac{\exp(-r^{1/\beta}|x|)}{(1 + V(x))(1 + |x|)^{(d+\beta+1)/2}} \frac{\exp(-r^{1/\beta}|y|)}{(1 + V(y))(1 + |y|)^{(d+\beta+1)/2}},$$

that the eigenfunction $\tilde{\phi}_0$ corresponding to the largest eigenvalue of the generator of $\{P_t : t \geq 0\}$ of ξ is bounded and that $\{P_t : t \geq 0\}$ is intrinsically ultracontractive. Thus, Assumptions 1.1 and 1.2 are satisfied.

In the next five examples, the processes may not symmetric.

EXAMPLE 1.18. Suppose that $\beta \in (0, 2)$ and that $\xi^{(1)} = \{\xi_t^{(1)} : t \geq 0\}$ is a strictly β -stable process in \mathbb{R}^d . Suppose that, in the case $d \geq 2$, the spherical part η of the Lévy measure μ of $\xi^{(1)}$ satisfies the following assumption: there exist a positive function Φ on the unit sphere S in \mathbb{R}^d and $\kappa > 1$ such that

$$\Phi = \frac{d\eta}{d\sigma} \quad \text{and} \quad \kappa^{-1} \leq \Phi(z) \leq \kappa \quad \text{on } S,$$

where σ is the surface measure on S . In the case $d = 1$, we assume that the Lévy measure of $\xi^{(1)}$ is given by

$$\mu(dx) = c_1 x^{-1-\beta} 1_{\{x>0\}} + c_2 |x|^{-1-\beta} 1_{\{x<0\}}$$

with $c_1, c_2 > 0$. Suppose that E is an open set in \mathbb{R}^d of finite Lebesgue measure. Let ξ be the process in E obtained by killing $\xi^{(1)}$ upon exiting E . Then it follows from [30, Example 4.1] that ξ has a transition density $p(t, x, y)$ which is a strictly positive, bounded continuous function of (x, y) for any $t > 0$. Thus Assumption 1.1 is trivially satisfied. It follows also from [30, Example 4.1] that the semigroup $\{P_t : t \geq 0\}$ of ξ is intrinsic ultracontractive and that the eigenfunction $\tilde{\phi}_0$ corresponding to the largest eigenvalue of the generator of $\{P_t : t \geq 0\}$ is bounded. Thus, Assumption 1.2 is also satisfied.

EXAMPLE 1.19. Suppose that $\beta \in (0, 2)$ and that $\xi^{(2)} = \{\xi_t^{(2)} : t \geq 0\}$ is a truncated strictly β -stable process in \mathbb{R}^d , that is, $\xi^{(2)}$ is a Lévy process with Lévy measure given by

$$\tilde{\mu}(dx) = \mu(dx) 1_{\{|x|<1\}},$$

where μ is the Lévy measure of the process $\xi^{(1)}$ in the previous example. Suppose that E is a connected open set in \mathbb{R}^d of finite Lebesgue measure. Let ξ be the process in E obtained by killing $\xi^{(2)}$ upon exiting E . Then it follows from [30, Example 4.2 and Proposition 4.4] that ξ has a transition density $p(t, x, y)$ which is a strictly positive, bounded continuous function of (x, y) for any $t > 0$. Thus, Assumption 1.1 is trivially satisfied. It follows also from [30, Example 4.2 and Proposition 4.4] that the semigroup $\{P_t : t \geq 0\}$

of ξ is intrinsic ultracontractive and that the eigenfunction $\tilde{\phi}_0$ corresponding to the largest eigenvalue of the generator of $\{P_t : t \geq 0\}$ is bounded. Thus, Assumption 1.2 is also satisfied.

EXAMPLE 1.20. Suppose $\beta \in (0, 2)$, $\xi^{(1)} = \{\xi_t^{(1)} : t \geq 0\}$ is a strictly β -stable process in \mathbb{R}^d satisfying the assumptions in Example 1.18 and that $B = \{B_t : t \geq 0\}$ is an independent Brownian motion in \mathbb{R}^d . Let $\xi^{(3)}$ be the process defined by $\xi_t^{(3)} = \xi_t^{(1)} + B_t$. Suppose that E is an open set in \mathbb{R}^d of finite Lebesgue measure. Let ξ be the process in E obtained by killing $\xi^{(3)}$ upon exiting E . Then it follows from [30, Example 4.5 and Lemma 4.6] that ξ has a transition density $p(t, x, y)$ which is a strictly positive, bounded continuous function of (x, y) for any $t > 0$. Thus Assumption 1.1 is trivially satisfied. It follows also from [30, Example 4.5 and Lemma 4.6] that the semigroup $\{P_t : t \geq 0\}$ of ξ is intrinsic ultracontractive and that the eigenfunction $\tilde{\phi}_0$ corresponding to the largest eigenvalue of the generator of $\{P_t : t \geq 0\}$ is bounded. Thus, Assumption 1.2 is also satisfied.

EXAMPLE 1.21. Suppose $\beta \in (0, 2)$, $\xi^{(2)} = \{\xi_t^{(2)} : t \geq 0\}$ is a truncated strictly β -stable process in \mathbb{R}^d satisfying the assumptions in Example 1.19 and that $B = \{B_t : t \geq 0\}$ is an independent Brownian motion in \mathbb{R}^d . Let $\xi^{(4)}$ be the process defined by $\xi_t^{(4)} = \xi_t^{(2)} + B_t$. Suppose that E is a connected open set in \mathbb{R}^d of finite Lebesgue measure. Let ξ be the process in E obtained by killing $\xi^{(4)}$ upon exiting E . Then it follows from [30, Example 4.7 and Lemma 4.8] that ξ has a transition density $p(t, x, y)$ which is a strictly positive, bounded continuous function of (x, y) for any $t > 0$. Thus Assumption 1.1 is trivially satisfied. It follows also from [30, Example 4.7 and Lemma 4.8] that the semigroup $\{P_t : t \geq 0\}$ of ξ is intrinsic ultracontractive and that the eigenfunction $\tilde{\phi}_0$ corresponding to the largest eigenvalue of the generator of $\{P_t : t \geq 0\}$ is bounded. Thus Assumption 1.2 is also satisfied.

EXAMPLE 1.22. Suppose $d \geq 3$ and that $\mu = (\mu^1, \dots, \mu^d)$, where each μ^j is a signed measure on \mathbb{R}^d such that

$$\limsup_{r \rightarrow 0} \sup_{x \in \mathbb{R}^d} \int_{B(x, r)} \frac{|\mu^j|(dy)}{|x - y|^{d-1}} = 0.$$

Let $\xi^{(5)} = \{\xi_t^{(5)} : t \geq 0\}$ be a Brownian motion with drift μ in \mathbb{R}^d , see [26]. Suppose that E is a bounded connected open set in \mathbb{R}^d and that $K > 0$ is a constant such that $E \subset B(0, K/2)$. Put $B = B(0, K)$. Let G_B be the Green function of $\xi^{(5)}$ in B and define $H(x) := \int_B G_B(x, y) dy$. Then H is a strictly positive continuous function on B . Let ξ be the process obtained by killing $\xi^{(5)}$ upon exiting E . Let m be the measure on E defined by $m(dx) = H(x) dx$. Then it follows from [55, Example 4.6] or [27], [29] that ξ has a transition density $p(t, x, y)$ with respect to m and that $p(t, x, y)$ is a strictly positive,

bounded continuous function of (x, y) for any $t > 0$. Thus Assumption 1.1 is trivially satisfied. It follows also from [55, Example 4.6] or [27], [29] that the semigroup $\{P_t : t \geq 0\}$ of ξ is intrinsic ultracontractive and that the eigenfunction $\tilde{\phi}_0$ corresponding to the largest eigenvalue of the generator of $\{P_t : t \geq 0\}$ is bounded. Thus, Assumption 1.2 is also satisfied.

EXAMPLE 1.23. Suppose $d \geq 2$, $\beta \in (1, 2)$, and that $\mu = (\mu^1, \dots, \mu^d)$, where each μ^j is a signed measure on \mathbb{R}^d such that

$$\limsup_{r \rightarrow 0} \sup_{x \in \mathbb{R}^d} \int_{B(x,r)} \frac{|\mu^j|(dy)}{|x-y|^{d-\beta+1}} = 0.$$

Let $\xi^{(6)} = \{\xi_t^{(6)} : t \geq 0\}$ be an β -stable process with drift μ in \mathbb{R}^d , see [31]. Suppose that E is a bounded open set in \mathbb{R}^d and suppose $K > 0$ is such that $D \subset B(0, K/2)$. Put $B = B(0, K)$. Let G_B be the Green function of $\xi^{(6)}$ in B and define $H(x) := \int_B G_B(x, y) dy$. Then H is a strictly positive continuous function on B . Let ξ be the process obtained by killing $\xi^{(6)}$ upon exiting D . Let m be the measure on E defined by $m(dx) = H(x) dx$. Then it follows from [55, Example 4.7] or [8] that ξ has a transition density $p(t, x, y)$ with respect to m and that $p(t, x, y)$ is a strictly positive, bounded continuous function of (x, y) for any $t > 0$. Thus, Assumption 1.1 is trivially satisfied. It follows also from [55, Example 4.7] or [8] that the semigroup $\{P_t : t \geq 0\}$ of ξ is intrinsic ultracontractive and that the eigenfunction $\tilde{\phi}_0$ corresponding to the largest eigenvalue of the generator of $\{P_t : t \geq 0\}$ is bounded. Thus, Assumption 1.2 is also satisfied.

2. Preliminaries

2.1. **Density of $\{T_t : t \geq 0\}$.** In this subsection, we show that, under Assumption 1.1, the semigroup $\{T_t : t \geq 0\}$ has a strictly positive density $q(t, x, y)$ and, for any $t > 0$, $q(t, x, y)$ is continuous in (x, y) .

LEMMA 2.1. *Suppose that Assumption 1.1 holds. The semigroup $\{T_t : t \geq 0\}$ has a density $q(t, x, y)$ such that*

$$(2.1) \quad e^{-Kt} p(t, x, y) \leq q(t, x, y) \leq e^{Kt} p(t, x, y), \quad (t, x, y) \in (0, \infty) \times E \times E.$$

Furthermore, for any $t > 0$, $q(t, x, y)$ is a continuous function of (x, y) on $E \times E$.

Proof. For any $(t, x, y) \in (0, \infty) \times E \times E$, define

$$I_0(t, x, y) := p(t, x, y),$$

$$I_n(t, x, y) := \int_0^t \int_E p(s, x, z) I_{n-1}(t-s, z, y) \alpha(z) m(dz) ds, \quad n \geq 1.$$

Using arguments similar to those in Section 1.2 of [45], we easily get that the function

$$(2.2) \quad q(t, x, y) := \sum_{n=0}^{\infty} I_n(t, x, y), \quad (t, x, y) \in (0, \infty) \times E \times E$$

is well defined and $q(t, x, y)$ is the density of T_t satisfying (2.1). We omit the details.

We now prove the continuity of $q(t, x, y)$ in $(x, y) \in E \times E$ for each fixed $t > 0$. As in Section 1.2 of [45], it suffices to show that, for any $0 < \varepsilon < t/2$,

$$\int_{\varepsilon}^{t-\varepsilon} \int_E p(s, x, z)p(t-s, z, y)\alpha(z)m(dz) ds$$

is continuous on $E \times E$. By (1.14), we get that

$$\begin{aligned} & p(s, x, z)p(t-s, z, y)|\alpha(z)| \\ & \leq K a_{\varepsilon/2}(x)^{1/2} \widehat{a}_{\varepsilon/2}(y)^{1/2} \widehat{a}_{s-\varepsilon/2}(z)^{1/2} a_{t-s-\varepsilon/2}(z)^{1/2}. \end{aligned}$$

We claim that the function $t \rightarrow \int_E a_t(x)m(dx)$ is decreasing. Using this claim and Hölder's inequality, we get that

$$\begin{aligned} & \int_{\varepsilon}^{t-\varepsilon} \int_E \widehat{a}_{s-\varepsilon/2}(z)^{1/2} a_{t-s-\varepsilon/2}(z)^{1/2} m(dz) ds \\ & \leq \int_{\varepsilon}^{t-\varepsilon} \left(\int_E \widehat{a}_{s-\varepsilon/2}(z) m(dz) \right)^{1/2} \left(\int_E a_{t-s-\varepsilon/2}(z) m(dz) \right)^{1/2} ds \\ & = \int_{\varepsilon}^{t-\varepsilon} \left(\int_E a_{s-\varepsilon/2}(z) m(dz) \right)^{1/2} \left(\int_E a_{t-s-\varepsilon/2}(z) m(dz) \right)^{1/2} ds \\ & \leq t \int_E a_{\varepsilon/2}(z) m(dz). \end{aligned}$$

The equality above follows from the fact $\int_E \widehat{a}_t(z)m(dz) = \int_E a_t(z)m(dz)$. Thus, by Assumption 1.1(ii) and the dominated convergence theorem, we get that the function

$$(x, y) \mapsto \int_{\varepsilon}^{t-\varepsilon} \int_E p(s, x, z)p(t-s, z, y)\alpha(z)m(dz) ds$$

is continuous.

Now, we prove the claim that the function $t \rightarrow \int_E a_t(x)m(dx)$ is decreasing. In fact, by Fubini's theorem and Hölder's inequality, we get

$$\begin{aligned} a_{t+s}(x) &= \int_E p(t+s, x, y) \int_E p(t, x, z)p(s, z, y)m(dz)m(dy) \\ &= \int_E p(t, x, z) \int_E p(t+s, x, y)p(s, z, y)m(dy)m(dz) \\ &\leq a_{t+s}(x)^{1/2} \int_E p(t, x, z)a_s(z)^{1/2}m(dz), \end{aligned}$$

which implies that

$$(2.3) \quad a_{t+s}(x) \leq \left(\int_E p(t, x, z) a_s(z)^{1/2} m(dz) \right)^2 \leq \int_E p(t, x, z) a_s(z) m(dz).$$

Thus, by Fubini’s theorem and Assumption 1.1(i), we get that

$$(2.4) \quad \int_E a_{t+s}(x) m(dx) \leq \int_E a_s(z) \int_E p(t, x, z) m(dx) m(dz) \leq \int_E a_s(z) m(dz).$$

We have now finished the proof of our claim. □

2.2. Extinction and non-extinction of $\{X_t : t \geq 0\}$. In this subsection, we will give some sufficient conditions for Assumption 1.4, see Lemma 2.3 below. In the case when the function $a(x)$ in (1.6) is identically zero, this lemma follows from [13, Lemma 11.5.1]. Here we provide a proof for completeness.

Let $\tilde{\Psi}(x, z)$ be a function on $E_\partial \times (0, \infty)$ with the form:

$$(2.5) \quad \begin{aligned} \tilde{\Psi}(x, z) = & -\tilde{a}(x)z + \tilde{b}(x)z^2 \\ & + \int_{(0, +\infty)} (e^{-zy} - 1 + zy)\tilde{n}(x, dy), \quad x \in E_\partial, z \geq 0, \end{aligned}$$

where $\tilde{a} \in \mathcal{B}_b(E_\partial)$, $\tilde{b} \in \mathcal{B}_b^+(E_\partial)$ and \tilde{n} is a kernel from E_∂ to $(0, \infty)$ satisfying

$$(2.6) \quad \int_{(0, +\infty)} (y \wedge y^2)\tilde{n}(x, dy) < \infty.$$

The following Lemma 2.2 is similar to [39, Corollary 5.18]. Recall that, unless explicitly mentioned otherwise, every function f on E is automatically extended to E_∂ by setting $f(\partial) = 0$. The function g in the lemma below may not satisfy $g(\partial) = 0$.

LEMMA 2.2. *Suppose that $\Psi(x, z) \geq \tilde{\Psi}(x, z)$ for all $x \in E$ and $z \geq 0$. Assume that f and g are bounded nonnegative measurable functions on E_∂ such that $f(\partial) = 0$ and $f(x) \leq g(x)$ for all $x \in E_\partial$. If $v_g(t, x)$ is the unique locally bounded non-negative solution to the equation:*

$$v_g(t, x) = -\Pi_x \int_0^t \tilde{\Psi}(\xi_s, v_g(t-s, \xi_s)) ds + \Pi_x g(\xi_t), \quad x \in E_\partial, t \geq 0,$$

then $v_g(t, x) \geq u_f(t, x)$ for all $t \geq 0$ and $x \in E$, where u_f is the unique locally bounded non-negative solution to (1.9).

Proof. Recall that $u_f(t, \partial) = 0$ and

$$u_f(t, x) = - \int_0^t \Pi_x(\Psi(\xi_s, u_f(t-s, \xi_s))) ds + \Pi_x(f(\xi_t)), \quad x \in E_\partial.$$

Define another branching mechanism $\Psi_1(x, z)$ as follows:

$$\Psi_1(x, z) = \begin{cases} \tilde{\Psi}(x, z), & x \in E; \\ 0, & x = \partial. \end{cases}$$

Put $g_1(x) = g(x)1_E(x)$, for $x \in E_\partial$. Then, for all $x \in E_\partial$, $\Psi_1(x, z) \leq \Psi(x, z)$ and $f(x) \leq g_1(x)$. Let $u_{g_1}^1(t, x)$ be the unique locally bounded non-negative solution to the equation:

$$u_{g_1}^1(t, x) = - \int_0^t \Pi_x(\Psi_1(\xi_s, u_{g_1}^1(t-s, \xi_s))) ds + \Pi_x(g_1(\xi_t)), \quad x \in E_\partial, t \geq 0.$$

It follows from [39, Corollary 5.18] that

$$(2.7) \quad u_f(t, x) \leq u_{g_1}^1(t, x), \quad x \in E, t \geq 0.$$

By [39, Proposition 2.20], we have $u_{g_1}^1(t, \partial) \leq \Pi_\partial[e^{\int_0^t \alpha(\xi_s) ds} g_1(\xi_t)] = 0$, here we used the fact that $g_1(\partial) = 0$. Therefore $u_{g_1}^1(t, \partial) = 0$. Since $\tilde{\Psi}(x, 0) = 0$, we have that

$$\begin{aligned} \Pi_x(\Psi_1(\xi_s, u_{g_1}^1(t-s, \xi_s))) &= \Pi_x(\tilde{\Psi}(\xi_s, u_{g_1}^1(t-s, \xi_s)); s < \zeta) \\ &= \Pi_x(\tilde{\Psi}(\xi_s, u_{g_1}^1(t-s, \xi_s))), \end{aligned}$$

which implies that $u_{g_1}^1(t, x)$ is the unique locally bounded non-negative solution to the equation:

$$u_{g_1}^1(t, x) = - \int_0^t \Pi_x(\tilde{\Psi}(\xi_s, u_{g_1}^1(t-s, \xi_s))) ds + \Pi_x(g_1(\xi_t)), \quad x \in E_\partial, t \geq 0.$$

Since $g_1(x) \leq g(x)$, for all $x \in E_\partial$, by [39, Corollary 5.18], we have

$$(2.8) \quad u_{g_1}^1(t, x) \leq v_g(t, x), \quad x \in E, t \geq 0.$$

Combining (2.7) and (2.8), we arrive at the desired assertion of this lemma. □

LEMMA 2.3. *Suppose that $\tilde{\Psi}(z) \leq \inf_{x \in E} \Psi(x, z)$, and $\tilde{\Psi}(z)$ can be written in the form*

$$\tilde{\Psi}(z) = \tilde{a}z + \tilde{b}z^2 + \int_0^\infty (e^{-zy} - 1 + zy)\tilde{n}(dy)$$

with $\tilde{a} \in \mathbb{R}$, $\tilde{b} \geq 0$ and \tilde{n} is a measure on $(0, \infty)$ satisfying $\int_0^\infty (y \wedge y^2)\tilde{n}(dy) < \infty$. If $\tilde{\Psi}(\infty) = \infty$ and $\tilde{\Psi}(z)$ satisfies

$$(2.9) \quad \int^\infty \frac{1}{\tilde{\Psi}(z)} dz < \infty,$$

then, for any $t > 0$, $\inf_{x \in E} q_t(x) > 0$.

Proof. Let \tilde{X} be a continuous state branching process with branching mechanism $\tilde{\Psi}$. Let $\tilde{\mathbb{P}}$ be the law of \tilde{X} with $\tilde{X}_0 = 1$. Define

$$u_\theta(t, x) = -\log \mathbb{P}_{\delta_x} e^{-\theta \|X_t\|}, \quad v_\theta(t) = -\log \tilde{\mathbb{P}} e^{-\theta \tilde{X}_t}.$$

It is easy to see that $u_\theta(t, \partial) = 0$ and, for $x \in E$ and $t > 0$,

$$u_\theta(t, x) = -\Pi_x \int_0^t \Psi(\xi_s, u_f(t-s, \xi_s)) ds + \theta \Pi_x(t < \zeta)$$

and

$$v_\theta(t) = -\int_0^t \tilde{\Psi}(v_\theta(s)) ds + \theta.$$

Applying Lemma 2.2 with $\tilde{\Psi}(x, z) = \tilde{\Psi}(z)$, $x \in E_\partial, z \geq 0$ and $g(x) = \theta$, $x \in E_\partial$, we get that, for all $t > 0$, $x \in E$ and $\theta > 0$, $u_\theta(t, x) \leq v_\theta(t)$. Letting $\theta \rightarrow \infty$, we get $-\log \mathbb{P}_{\delta_x}(\|X_t\| = 0) \leq -\log \tilde{\mathbb{P}}(\tilde{X}_t = 0)$. It is well known that, under the conditions of this lemma, $\tilde{\mathbb{P}}(\tilde{X}_t = 0) > 0$. Thus $\inf_{x \in E} q_t(x) = \inf_{x \in E} \mathbb{P}_{\delta_x}(\|X_t\| = 0) \geq \tilde{\mathbb{P}}(\tilde{X}_t = 0) > 0$. \square

It was proved in [50] that (2.9) is equivalent to $\int^\infty \frac{1}{\tilde{\Phi}(z)} dz < \infty$, where $\tilde{\Phi}(z) := \tilde{\Psi}(z) - \tilde{a}z$. Lemma 2.3 says that if the spatially dependent branching mechanism $\Psi(x, z)$ is dominated from below by a spatially independent branching mechanism $\tilde{\Psi}(z)$ satisfying $\tilde{\Psi}(\infty) = \infty$ and (2.9), then Assumption 1.4 holds. In particular, when Ψ does not depend on the spatial variable x and satisfies $\Psi(\infty) = \infty$ and the condition $\int^\infty \frac{1}{\Psi(\lambda)} d\lambda < \infty$, Assumption 1.4 holds. If $\tilde{b} := \inf_{x \in E} b(x)\beta(x) > 0$, then $\Psi(x, z) \geq -Kz + \tilde{b}z^2$, where K is the constant given in (1.11). In this case, we can take $\tilde{\Psi}(z) := -Kz + \tilde{b}z^2$ and it is clear that $\tilde{\Psi}(z)$ satisfies (2.9).

2.3. Estimates on moments. In the remainder of this paper, we will use the following notation: for two positive functions f and g on E , $f(x) \lesssim g(x)$ for $x \in E$ means that there exists a constant $c > 0$ such that $f(x) \leq cg(x)$ for all $x \in E$. Throughout this paper, c is a constant whose value may vary from line to line.

By (1.19) and the assumption that $\lambda_0 = 0$, we have, for any $(t, x, y) \in (1, \infty) \times E \times E$,

$$(2.10) \quad |q(t, x, y) - \phi_0(x)\psi_0(y)| \leq ce^{-\gamma t} \phi_0(x)\psi_0(y).$$

It follows that, if $f \in \mathcal{C}_1$, we have, for $(t, x) \in (1, \infty) \times E$,

$$(2.11) \quad |T_t f(x) - \langle f, \psi_0 \rangle_m \phi_0(x)| \leq ce^{-\gamma t} \langle |f|, \psi_0 \rangle_m \phi_0(x)$$

and

$$(2.12) \quad |T_t f(x)| \leq (1+c) \langle |f|, \psi_0 \rangle_m \phi_0(x).$$

Recall the second moment formula of the superprocess $\{X_t : t \geq 0\}$ (see, for example, [39, Corollary 2.39]): for $f \in \mathcal{B}_b(E)$, we have for any $t > 0$,

$$(2.13) \quad \mathbb{P}_\mu \langle f, X_t \rangle^2 = (\mathbb{P}_\mu \langle f, X_t \rangle)^2 + \int_E \int_0^t T_s [A(T_{t-s}f)^2](x) ds \mu(dx).$$

Thus,

$$(2.14) \quad \text{Var}_\mu \langle f, X_t \rangle = \langle \text{Var}_{\delta_x} \langle f, X_t \rangle, \mu \rangle = \int_E \int_0^t T_s [A(T_{t-s}f)^2](x) ds \mu(dx),$$

where Var_μ stands for the variance under \mathbb{P}_μ . For any $f \in \mathcal{C}_2$ and $x \in E$, applying the Cauchy–Schwarz inequality, we have $(T_{t-s}f)^2(x) \leq e^{K(t-s)} T_{t-s}(f^2)(x)$, which implies that

$$(2.15) \quad \int_0^t T_s [A(T_{t-s}f)^2](x) ds \leq e^{Kt} T_t(f^2)(x) < \infty.$$

Thus, using a routine limit argument, one can easily check that (2.13) and (2.14) also hold for $f \in \mathcal{C}_2$.

LEMMA 2.4. *Assume that $f \in \mathcal{C}_2$. If $\langle f, \psi_0 \rangle_m = 0$, then for $(t, x) \in (2, \infty) \times E$, we have*

$$(2.16) \quad |\text{Var}_{\delta_x} \langle f, X_t \rangle - \sigma_f^2 \phi_0(x)| \lesssim e^{-\gamma t} \phi_0(x),$$

where σ_f^2 is defined in (1.27). Therefore, for $(t, x) \in (2, \infty) \times E$, we have

$$(2.17) \quad \text{Var}_{\delta_x} \langle f, X_t \rangle \lesssim \phi_0(x).$$

Proof. First, we show that $\sigma_f^2 < \infty$. For $s \leq 1$, $|T_s f(x)|^2 \leq e^{Ks} T_s(f^2)(x)$. Hence, for $s \leq 1$,

$$(2.18) \quad \langle A(T_s f)^2, \psi_0 \rangle_m \leq K e^{sK} \langle T_s(f^2), \psi_0 \rangle_m = K e^{sK} \langle f^2, \psi_0 \rangle_m.$$

For $s > 1$, by (2.11), $|T_s f(x)| \lesssim e^{-\gamma s} \langle |f|, \psi_0 \rangle_m \phi_0(x)$. Hence, for $s > 1$,

$$(2.19) \quad \langle A(T_s f)^2, \psi_0 \rangle_m \lesssim e^{-2\gamma s}.$$

Therefore, combining (2.18) and (2.19) we have that

$$\sigma_f^2 = \int_0^\infty \langle A(T_s f)^2, \psi_0 \rangle_m ds \lesssim \int_0^1 e^{sK} ds + \int_1^\infty e^{-2\gamma s} ds < \infty.$$

By (2.14), for $t > 2$, we have

$$(2.20) \quad \begin{aligned} & |\text{Var}_{\delta_x} \langle f, X_t \rangle - \sigma_f^2 \phi_0(x)| \\ & \leq \int_0^{t-1} |T_{t-s} [A(T_s f)^2](x) - \langle A(T_s f)^2, \psi_0 \rangle_m \phi_0(x)| ds \\ & \quad + \int_{t-1}^t T_{t-s} [A(T_s f)^2](x) ds + \int_{t-1}^\infty \langle A(T_s f)^2, \psi_0 \rangle_m ds \phi_0(x) \\ & =: V_1(t, x) + V_2(t, x) + V_3(t, x). \end{aligned}$$

First, we consider $V_1(t, x)$. By (2.11), for $t - s > 1$, we have

$$|T_{t-s}[A(T_s f)^2](x) - \langle A(T_s f)^2, \psi_0 \rangle_m \phi_0(x)| \lesssim e^{-\gamma(t-s)} \langle A(T_s f)^2, \psi_0 \rangle_m \phi_0(x).$$

Therefore, by (2.18) and (2.19), we have, for $(t, x) \in (2, \infty) \times E$,

$$(2.21) \quad V_1(t, x) \lesssim \int_1^t e^{-\gamma(t+s)} ds \phi_0(x) + \int_0^1 e^{-\gamma(t-s)} ds \phi_0(x) \lesssim e^{-\gamma t} \phi_0(x).$$

For $V_2(t, x)$, by (2.11), for $s > t - 1 > 1$, $|T_s f(x)| \lesssim e^{-\gamma s} \phi_0(x)$. Thus,

$$(2.22) \quad V_2(t, x) \lesssim \int_{t-1}^t e^{-2\gamma s} T_{t-s}[\phi_0^2](x) ds = e^{-2\gamma t} \int_0^1 e^{2\gamma s} T_s[\phi_0^2](x) ds.$$

By Hölder's inequality, we have

$$\phi_0^2(x) = (T_1 \phi_0(x))^2 \leq e^K T_1(\phi_0^2)(x).$$

Thus by (2.22) and (2.12), for $(t, x) \in (2, \infty) \times E$, we have

$$(2.23) \quad V_2(t, x) \lesssim e^{-2\gamma t} \int_0^1 T_{s+1}(\phi_0^2)(x) ds \lesssim e^{-2\gamma t} \phi_0(x).$$

For $V_3(t, x)$, by (2.11), for $s > t - 1 > 1$, $|T_s f(x)| \lesssim e^{-\gamma s} \phi_0(x)$. Thus,

$$(2.24) \quad V_3(t, x) \lesssim \int_{t-1}^\infty e^{-2\gamma s} ds \langle \phi_0^2, \psi_0 \rangle_m \phi_0(x) \lesssim e^{-2\gamma t} \phi_0(x).$$

It follows from (2.21), (2.23) and (2.24) that, for $(t, x) \in (2, \infty) \times E$,

$$|\text{Var}_{\delta_x} \langle f, X_t \rangle - \sigma_f^2 \phi_0(x)| \lesssim e^{-\gamma t} \phi_0(x).$$

Now (2.17) follows immediately. □

2.4. Excursion measures of $\{X_t : t \geq 0\}$. We use \mathbb{D} to denote the space of $\mathcal{M}_F(E)$ -valued right continuous functions $t \mapsto \omega_t$ on $(0, \infty)$ having zero as a trap. We use $(\mathcal{A}, \mathcal{A}_t)$ to denote the natural σ -algebras on \mathbb{D} generated by the coordinate process.

Under Assumption 1.4, it is known (see [39, Chapter 8]) that one can associate with $\{\mathbb{P}_{\delta_x} : x \in E\}$ a family of σ -finite measures $\{\mathbb{N}_x : x \in E\}$ defined on $(\mathbb{D}, \mathcal{A})$ such that $\mathbb{N}_x(\{0\}) = 0$, and

$$\mathbb{N}_x(1 - e^{-\langle f, \omega_t \rangle}) = -\log \mathbb{P}_{\delta_x}(e^{-\langle f, X_t \rangle}), \quad f \in \mathcal{B}_b^+(E), t \geq 0.$$

For further information on excursion measures of superprocesses, we refer the reader to [16], [17], [38].

For any $\mu \in \mathcal{M}_F(E)$, let $N(d\omega)$ be a Poisson random measure on the space \mathbb{D} with intensity $\int_E \mathbb{N}_x(d\omega) \mu(dx)$, in a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \mathbf{P}_\mu)$. We define another process $\{\Lambda_t : t \geq 0\}$ by $\Lambda_0 = \mu$ and

$$\Lambda_t := \int_{\mathbb{D}} \omega_t N(d\omega), \quad t > 0.$$

Let $\tilde{\mathcal{F}}_t$ be the σ -algebra generated by the random variables $\{N(A) : A \in \mathcal{A}_t\}$. Then $\{\Lambda, (\tilde{\mathcal{F}}_t)_{t \geq 0}, \mathbf{P}_\mu\}$ has the same law as $\{X, (\mathcal{G}_t)_{t \geq 0}, \mathbb{P}_\mu\}$, see [39, Theorem 8.24].

Now we list some properties of \mathbb{N}_x . The proofs are similar to those of [16, Corollary 1.2, Proposition 1.1].

PROPOSITION 2.5. *If $\mathbb{P}_{\delta_x}|\langle f, X_t \rangle| < \infty$, then*

$$(2.25) \quad \mathbb{N}_x \langle f, \omega_t \rangle = \mathbb{P}_{\delta_x} \langle f, X_t \rangle.$$

If $\mathbb{P}_{\delta_x} \langle f, X_t \rangle^2 < \infty$, then

$$(2.26) \quad \mathbb{N}_x \langle f, \omega_t \rangle^2 = \text{Var}_{\delta_x} \langle f, X_t \rangle.$$

For $f \in \mathcal{C}_1$, by (2.12), $T_t f$ is bounded and in \mathcal{C}_1 . It follows from Proposition 2.5 that, for any $f \in \mathcal{C}_1$,

$$\int_E \int_D \langle |f|, \omega_s \rangle \mathbb{N}_x(d\omega) X_t(dx) < \infty, \quad \mathbb{P}_\mu\text{-a.s.}$$

Now, by the Markov property of X , we get that for any $f \in \mathcal{C}_1$,

$$(2.27) \quad \begin{aligned} & \mathbb{P}_\mu \left[\exp\{i\theta \langle f, X_{t+s} \rangle\} \middle| X_t \right] \\ &= \mathbb{P}_{X_t} \left[\exp\{i\theta \langle f, X_s \rangle\} \right] = \mathbf{P}_{X_t} \left[\exp\{i\theta \langle f, \Lambda_s \rangle\} \right] \\ &= \exp \left\{ \int_E \int_{\mathbb{D}} (e^{i\theta \langle f, \omega_s \rangle} - 1) \mathbb{N}_x(d\omega) X_t(dx) \right\}. \end{aligned}$$

3. Proofs of main results

In this section, we will prove our main theorems.

3.1. Proofs of Theorems 1.5 and 1.6. For $x \in E$ and $z > 0$, define

$$(3.1) \quad r(x, z) = \Psi(x, z) + \alpha(x)z$$

and

$$(3.2) \quad r^{(2)}(x, z) = \Psi(x, z) + \alpha(x)z - \frac{1}{2}A(x)z^2.$$

LEMMA 3.1. *For any $x \in E$ and $z > 0$,*

$$(3.3) \quad 0 \leq r(x, z) \leq Kz^2/2$$

and

$$(3.4) \quad |r^{(2)}(x, z)| \leq e(x, z)z^2,$$

where

$$(3.5) \quad e(x, z) = \beta(x) \int_0^\infty y^2 \left(1 \wedge \frac{1}{6}yz \right) n(x, dy).$$

Proof. It is easy to see that

$$(3.6) \quad r(x, z) = \beta(x) \left(b(x)z^2 + \int_0^\infty (e^{-zy} - 1 + zy)n(x, dy) \right)$$

and

$$r^{(2)}(x, z) = \beta(x) \int_0^\infty \left(e^{-zy} - 1 + zy - \frac{1}{2}y^2z^2 \right) n(x, dy).$$

It follows from Taylor's expansion that, for $\theta > 0$,

$$(3.7) \quad 0 < e^{-\theta} - 1 + \theta \leq \frac{1}{2}\theta^2$$

and

$$(3.8) \quad \left| e^{-\theta} - 1 + \theta - \frac{1}{2}\theta^2 \right| \leq \frac{1}{6}\theta^3.$$

By (3.7), we also have $|e^{-\theta} - 1 + \theta - \frac{1}{2}\theta^2| \leq \theta^2$. Thus, we have

$$(3.9) \quad \left| e^{-\theta} - 1 + \theta - \frac{1}{2}\theta^2 \right| \leq \theta^2 \left(1 \wedge \frac{1}{6}\theta \right).$$

Therefore, by (3.7) and (3.9), we have

$$0 < r(x, z) \leq \beta(x) \left(b(x) + \frac{1}{2} \int_0^\infty y^2 n(x, dy) \right) z^2 \leq Kz^2/2$$

and

$$r^{(2)}(x, z) \leq \beta(x) \int_0^\infty y^2 \left(1 \wedge \frac{1}{6}yz \right) n(x, dy) z^2.$$

The proof is now complete. □

Recall that

$$u_f(t, x) := -\log \mathbb{P}_{\delta_x} e^{-\langle f, X_t \rangle}.$$

LEMMA 3.2. *If $f \in \mathcal{C}_1^+$, then $0 \leq u_f(t, x) < \infty$ for all $t \geq 0, x \in E$, and the function R_f defined by*

$$(3.10) \quad R_f(t, x) := T_t f(x) - u_f(t, x)$$

satisfies

$$(3.11) \quad R_f(t, x) = \int_0^t T_s [r(\cdot, u_f(t-s, \cdot))](x) ds, \quad t \geq 0, x \in E.$$

Moreover,

$$(3.12) \quad 0 \leq R_f(t, x) \leq e^{Kt} T_t (f^2)(x), \quad t \geq 0, x \in E.$$

Proof. First, we assume that $f \in \mathcal{B}_b^+$. Recall that $u_f(t, x) = -\log \mathbb{P}_{\delta_x} e^{-\langle f, X_t \rangle}$ satisfies

$$u_f(t, x) + \Pi_x \int_0^t \Psi(\xi_s, u_f(t-s, \xi_s)) ds = \Pi_x(f(\xi_t)), \quad t \geq 0, x \in E.$$

It follows from [39, Theorem 2.23] that $u_f(t, x)$ also satisfies

$$(3.13) \quad u_f(t, x) = -\int_0^t T_s[r(\cdot, u_f(t-s, \cdot))](x) ds + T_t f(x), \quad t \geq 0, x \in E.$$

Thus, we get (3.11) immediately.

For general $f \in C_1^+$, we have $T_t f(x) < \infty$. Let $f_n(x) = f(x) \wedge n \in \mathcal{B}_b^+$. Since (3.11) holds for f_n , applying the monotone convergence theorem, we get that (3.11) also holds for f . Therefore, by (3.3), $R_f(t, x) \geq 0$, which means $u_f(t, x) \leq T_t f(x) < \infty$. Recall that, as a consequence of the Cauchy–Schwarz inequality, we have $(T_{t-s} f)^2(y) \leq e^{K(t-s)} T_{t-s}(f^2)(y)$. Combining this with (3.3), we get

$$\begin{aligned} 0 \leq R_f(t, x) &\leq \frac{K}{2} \int_0^t T_s [(u_f(t-s))^2](x) ds \\ &\leq \frac{K}{2} \int_0^t T_s [(T_{t-s} f)^2](x) ds \leq e^{Kt} T_t(f^2)(x). \end{aligned} \quad \square$$

Recall that $q_t(x) = \mathbb{P}_{\delta_x}(\|X_t\| = 0)$.

LEMMA 3.3.

$$(3.14) \quad \liminf_{t \rightarrow \infty} \inf_{x \in E} q_t(x) = 1.$$

Proof. For $\theta > 0$, let

$$u_\theta(t, x) := -\log \mathbb{P}_{\delta_x} e^{-\langle \theta, X_t \rangle}.$$

By the Markov property of X ,

$$(3.15) \quad \begin{aligned} q_{t+s}(x) &= \lim_{\theta \rightarrow \infty} \mathbb{P}_{\delta_x}(e^{-\theta \|X_{t+s}\|}) \\ &= \lim_{\theta \rightarrow \infty} \mathbb{P}_{\delta_x}(e^{-\langle u_\theta(s), X_t \rangle}) = \mathbb{P}_{\delta_x}(e^{-\langle -\log q_s, X_t \rangle}). \end{aligned}$$

Since $q_t(x)$ is increasing in t , $q(x) := \lim_{t \rightarrow \infty} q_t(x)$ exists. Put $w(x) = -\log q(x)$. Letting $s \rightarrow \infty$ in (3.15), we get $q(x) = \mathbb{P}_{\delta_x}(e^{-\langle w, X_t \rangle})$, which implies, for $t > 0$,

$$(3.16) \quad w(x) = u_w(t, x), \quad x \in E.$$

By Assumption 1.4, for $s > t_0$,

$$\|w\|_\infty \leq \|-\log q_s\|_\infty \leq \|-\log q_{t_0}\|_\infty = -\log\left(\inf_{x \in E} q_{t_0}(x)\right) < \infty,$$

which implies $w \in \mathcal{C}_1^+$, and $-\log q_s \in \mathcal{C}_1^+$. Thus, by (3.10), (3.11) and (3.16), we have

$$(3.17) \quad w(x) = T_t(w)(x) - \int_0^t T_s(r(\cdot, w(\cdot)))(x) ds, \quad x \in E.$$

By (2.11), we have $\lim_{t \rightarrow \infty} T_t(w)(x) = \langle w, \psi_0 \rangle_m \phi_0(x)$.

If $\langle r(\cdot, w(\cdot)), \psi_0 \rangle_m > 0$, then

$$\lim_{t \rightarrow \infty} T_t(r(\cdot, w(\cdot)))(x) = \langle r(\cdot, w(\cdot)), \psi_0 \rangle_m \phi_0(x) > 0, \quad \text{for any } x \in E,$$

which implies

$$\lim_{t \rightarrow \infty} \int_0^t T_s[r(\cdot, w(\cdot))](x) ds = \infty, \quad \text{for any } x \in E.$$

Thus, by (3.17), we get

$$0 \leq w(x) = \lim_{t \rightarrow \infty} (T_t(w))(x) - \int_0^t T_s[r(\cdot, w(\cdot))](x) ds = -\infty,$$

which is a contradiction. Therefore, $r(x, w(x)) = 0$, a.e.-m. Then, by (3.17), we get, for all $x \in E$,

$$(3.18) \quad w(x) = \langle w, \psi_0 \rangle_m \phi_0(x),$$

which implies that $w \equiv 0$ on E or $w(x) > 0$ for any $x \in E$. Since $r(x, w(x)) = 0$, a.e.-m, by (3.6), we obtain $w \equiv 0$ on E . For $s > t_0$, by (3.15) and Lemma 3.2, we get

$$\begin{aligned} -\log q_{2+s}(x) &= u_{-\log q_s}(2, x) \leq T_2(-\log q_s)(x) \\ &\leq (1 + c) \langle -\log q_s, \psi_0 \rangle_m \|\phi_0\|_\infty, \end{aligned}$$

where in the last inequality we used (2.11). Since $-\log q_s(x) \rightarrow 0$, by the dominated convergence theorem, we get

$$\lim_{s \rightarrow \infty} \langle -\log q_s, \psi_0 \rangle_m = 0.$$

Now (3.14) follows immediately. □

LEMMA 3.4. For any $f \in \mathcal{C}_1^+$, there exists a function $h_f(t, x)$ such that

$$(3.19) \quad u_f(t, x) = (1 + h_f(t, x)) \langle u_f(t, \cdot), \psi_0 \rangle_m \phi_0(x).$$

Furthermore,

$$(3.20) \quad \lim_{t \rightarrow \infty} \|h_f(t)\|_\infty = 0 \quad \text{uniformly in } f \in \mathcal{C}_1^+.$$

Proof. For any $f \in \mathcal{C}_1^+$, we have $u_f(t, x) \leq T_t f(x) < \infty$ and $\langle u_f(t, \cdot), \psi_0 \rangle_m \leq \langle T_t f, \psi_0 \rangle_m = \langle f, \psi_0 \rangle_m < \infty$. So $u_f(t, x) \in \mathcal{C}_1^+$. If $m(f > 0) = 0$, then $T_t f(x) = 0$ for all $t > 0$ and $x \in E$, which implies $u_f(t, x) = 0$ and $\langle u_f(t, \cdot), \psi_0 \rangle_m = 0$. In this case, we define $h_f(t, x) = 0$. If $m(f > 0) > 0$, then $T_t f(x) > 0$ for all $t > 0$

and $x \in E$, which implies $\mathbb{P}_{\delta_x}(\langle f, X_t \rangle = 0) < 1$. Thus, we have $u_f(t, x) > 0$ and $\langle u_f(t, \cdot), \psi_0 \rangle_m > 0$. Define

$$h_f(t, x) = \frac{u_f(t, x) - \langle u_f(t, \cdot), \psi_0 \rangle_m \phi_0(x)}{\langle u_f(t, \cdot), \psi_0 \rangle_m \phi_0(x)}.$$

We only need to prove that $\|h_f(t, \cdot)\|_\infty \rightarrow 0$ uniformly in $f \in \mathcal{C}_1^+ \setminus \{0\}$ as $t \rightarrow \infty$. Since $\mathbb{P}_\mu(e^{-\langle f, X_t \rangle}) \geq \mathbb{P}_\mu(\|X_t\| = 0)$, we get that

$$(3.21) \quad \|u_f(t, \cdot)\|_\infty \leq \|-\log q_t\|_\infty \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

By the Markov property of X we have

$$(3.22) \quad u_f(t, x) = -\log \mathbb{P}_{\delta_x} e^{-\langle u_f(t-s, \cdot), X_s \rangle} = u_{u_f(t-s)}(s, x), \quad t \geq s > 0, x \in E,$$

where in the subscript on the right-hand side, $u_f(t-s)$ stands for the function $x \rightarrow u_f(t-s, x)$. In the remainder of this proof, we keep this convention. By (3.10), we have

$$(3.23) \quad u_f(t, x) = T_s(u_f(t-s, \cdot))(x) - R_{u_f(t-s)}(s, x).$$

Thus,

$$(3.24) \quad \langle u_f(t, \cdot), \psi_0 \rangle_m = \langle u_f(t-s, \cdot), \psi_0 \rangle_m - \langle R_{u_f(t-s)}(s, \cdot), \psi_0 \rangle_m.$$

Therefore, by (2.11), (2.12) and (3.12), we have, for $1 < s < t$ and $x \in E$,

$$\begin{aligned} & |u_f(t, x) - \langle u_f(t, \cdot), \psi_0 \rangle_m \phi_0(x)| \\ & \leq |T_s(u_f(t-s, \cdot))(x) - \langle u_f(t-s, \cdot), \psi_0 \rangle_m \phi_0(x)| \\ & \quad + |R_{u_f(t-s)}(s, x)| + |\langle R_{u_f(t-s)}(s, \cdot), \psi_0 \rangle_m \phi_0(x)| \\ & \leq ce^{-\gamma s} \langle u_f(t-s, \cdot), \psi_0 \rangle_m \phi_0(x) \\ & \quad + e^{Ks} T_s(u_f^2(t-s, \cdot))(x) + e^{Ks} \langle u_f^2(t-s, \cdot), \psi_0 \rangle_m \phi_0(x) \\ & \leq ce^{-\gamma s} \langle u_f(t-s, \cdot), \psi_0 \rangle_m \phi_0(x) + (2+c)e^{Ks} \langle u_f^2(t-s, \cdot), \psi_0 \rangle_m \phi_0(x) \\ & \leq [ce^{-\gamma s} + (2+c)e^{Ks} \|-\log q_{t-s}\|_\infty] \langle u_f(t-s, \cdot), \psi_0 \rangle_m \phi_0(x), \end{aligned}$$

where in the last inequality we used (3.21) and c is the constant in (2.10).

By Lemma 3.2 and (3.23), we get

$$(3.25) \quad \begin{aligned} & T_s(u_f(t-s, \cdot))(x) \\ & \geq u_f(t, x) \geq T_s(u_f(t-s, \cdot))(x) - e^{Ks} T_s(u_f^2(t-s, \cdot))(x) \\ & \geq T_s(u_f(t-s, \cdot))(x) - e^{Ks} \|-\log q_{t-s}\|_\infty T_s(u_f(t-s, \cdot))(x). \end{aligned}$$

Thus, we have

$$(3.26) \quad \begin{aligned} \langle u_f(t-s, \cdot), \psi_0 \rangle_m & \geq \langle u_f(t, \cdot), \psi_0 \rangle_m \\ & \geq (1 - e^{Ks} \|-\log q_{t-s}\|_\infty) \langle u_f(t-s, \cdot), \psi_0 \rangle_m. \end{aligned}$$

For any $s > 1$, $(1 - e^{Ks}\|-\log q_{t-s}\|_\infty) > 0$ when t is large enough. Therefore, as $t \rightarrow \infty$,

$$\|h_f(t, \cdot)\|_\infty \leq \frac{ce^{-\gamma s} + (2 + c)e^{Ks}\|-\log q_{t-s}\|_\infty}{1 - e^{Ks}\|-\log q_{t-s}\|_\infty} \rightarrow ce^{-\gamma s}.$$

Now, letting $s \rightarrow \infty$, we get $\|h_f(t, \cdot)\|_\infty \rightarrow 0$ uniformly in $f \in C_1^+ \setminus \{0\}$ as $t \rightarrow \infty$. □

LEMMA 3.5. *For any $\delta > 0$,*

$$(3.27) \quad \lim_{n \rightarrow \infty} \frac{1}{n\delta} \left(\frac{1}{\langle u_f(n\delta, \cdot), \psi_0 \rangle_m} - \frac{1}{\langle f, \psi_0 \rangle_m} \right) = \nu$$

uniformly in $f \in C_1^+ \setminus \{0\}$. Here ν is defined in (1.22).

Proof. In this proof, we sometimes use $u_f(t)$ to denote the function $x \rightarrow u_f(t, x)$. Since f is non-negative and $m(f > 0) > 0$, we have $u_f(t, x) > 0$ for all $t > 0$ and $x \in E$. Consequently, we have $\langle u_f(t), \psi_0 \rangle_m > 0$. It is clear that $u_f(0) = f$. First note that

$$\begin{aligned} & \frac{1}{n\delta} \left(\frac{1}{\langle u_f(n\delta), \psi_0 \rangle_m} - \frac{1}{\langle f, \psi_0 \rangle_m} \right) \\ &= \frac{1}{n\delta} \sum_{k=0}^{n-1} \left(\frac{1}{\langle u_f((k+1)\delta), \psi_0 \rangle_m} - \frac{1}{\langle u_f(k\delta), \psi_0 \rangle_m} \right) \\ &= \frac{1}{n\delta} \sum_{k=0}^{n-1} \left(\frac{\langle u_f(k\delta), \psi_0 \rangle_m - \langle u_f((k+1)\delta), \psi_0 \rangle_m}{\langle u_f((k+1)\delta), \psi_0 \rangle_m \langle u_f(k\delta), \psi_0 \rangle_m} \right). \end{aligned}$$

Recall the identity (3.22) and the definition of $r^{(2)}(x, z)$ given in (3.2). Using (3.24) with $t = (k+1)\delta$ and $s = \delta$, we get

$$\begin{aligned} & \langle u_f(k\delta), \psi_0 \rangle_m - \langle u_f((k+1)\delta), \psi_0 \rangle_m \\ &= \langle R_{u_f(k\delta)}(\delta, \cdot), \psi_0 \rangle_m \\ &= \int_0^\delta \langle r(\cdot, u_f(k\delta + s, \cdot)), \psi_0 \rangle_m ds \\ &= \frac{1}{2} \int_0^\delta \langle A(u_f(k\delta + s))^2, \psi_0 \rangle_m ds \\ & \quad + \int_0^\delta \langle r^{(2)}(\cdot, u_f(k\delta + s, \cdot)), \psi_0 \rangle_m ds \\ &=: I_1 + I_2. \end{aligned}$$

By (3.19) and (3.26), we have, for $s \in [0, \delta]$,

$$\begin{aligned} & |u_f(t+s, x) - \langle u_f(t), \psi_0 \rangle_m \phi_0(x)| \\ & \leq |u_f(t+s, x) - \langle u_f(t+s), \psi_0 \rangle_m \phi_0(x)| \end{aligned}$$

$$\begin{aligned}
 & + |\langle u_f(t), \psi_0 \rangle_m - \langle u_f(t+s), \psi_0 \rangle_m| \phi_0(x) \\
 & \leq \|h_f(t+s)\|_\infty \langle u_f(t+s), \psi_0 \rangle_m \phi_0(x) \\
 & \quad + e^{Ks} \|-\log q_t\|_\infty \langle u_f(t), \psi_0 \rangle_m \phi_0(x) \\
 & \leq (\|h_f(t+s)\|_\infty + e^{Ks} \|-\log q_t\|_\infty) \langle u_f(t), \psi_0 \rangle_m \phi_0(x) \\
 (3.28) \quad & \leq c_f(t) \langle u_f(t), \psi_0 \rangle_m \phi_0(x),
 \end{aligned}$$

where $c_f(t) = \sup_{0 \leq s \leq \delta} (\|h_f(t+s)\|_\infty + e^{Ks} \|-\log q_t\|_\infty)$. By (3.20) and Lemma 3.3, we get $c_f(t) \rightarrow 0$, as $t \rightarrow \infty$, uniformly in $f \in \mathcal{C}_1^+$. Thus, by (3.28) we have for $s \in [0, \delta]$,

$$(3.29) \quad \frac{|u_f(t+s, x)^2 - \langle u_f(t), \psi_0 \rangle_m^2 (\phi_0(x))^2|}{\langle u_f(t), \psi_0 \rangle_m^2} \leq (2 + c_f(t)) c_f(t) (\phi_0(x))^2.$$

Therefore, we have,

$$\begin{aligned}
 \left| \frac{I_1}{\langle u_f(k\delta), \psi_0 \rangle_m^2} - \delta\nu \right| &= \frac{|\int_0^\delta \langle A((u_f(k\delta + s))^2 - \langle u_f(k\delta), \psi_0 \rangle_m^2 \phi_0^2), \psi_0 \rangle_m ds|}{2\langle u_f(k\delta), \psi_0 \rangle_m^2} \\
 &\leq \frac{1}{2} \langle A\phi_0^2, \psi_0 \rangle_m \delta (2 + c_f(k\delta)) c_f(k\delta) \rightarrow 0, \quad \text{as } k \rightarrow \infty,
 \end{aligned}$$

uniformly in $f \in \mathcal{C}_1^+ \setminus \{0\}$. By (3.26), we have

$$0 \leq 1 - \frac{\langle u_f((k+1)\delta), \psi_0 \rangle_m}{\langle u_f(k\delta), \psi_0 \rangle_m} \leq e^{K\delta} \|-\log q_{k\delta}\|_\infty,$$

which implies that

$$(3.30) \quad \frac{\langle u_f(k\delta), \psi_0 \rangle_m}{\langle u_f((k+1)\delta), \psi_0 \rangle_m} \rightarrow 1, \quad \text{as } k \rightarrow \infty,$$

uniformly in $f \in \mathcal{C}_1^+ \setminus \{0\}$. It follows that

$$(3.31) \quad \lim_{k \rightarrow \infty} \frac{I_1}{\langle u_f(k\delta), \psi_0 \rangle_m \langle u_f((k+1)\delta), \psi_0 \rangle_m} = \delta\nu$$

uniformly in $f \in \mathcal{C}_1^+ \setminus \{0\}$.

For I_2 , by (3.4) and (3.28), we have

$$\begin{aligned}
 \frac{\langle r^{(2)}(\cdot, u_f(k\delta + s, \cdot)), \psi_0 \rangle_m}{\langle u_f(k\delta), \psi_0 \rangle_m^2} &\leq \frac{\langle e(\cdot, u_f(k\delta + s, \cdot)) u_f(k\delta + s)^2, \psi_0 \rangle_m}{\langle u_f(k\delta), \psi_0 \rangle_m^2} \\
 &\leq (1 + c_f(k\delta))^2 \langle e(\cdot, u_f(k\delta + s, \cdot)) \phi_0^2, \psi_0 \rangle_m \\
 &\leq (1 + c_f(k\delta))^2 \langle e(\cdot, \|-\log q_{k\delta}\|_\infty) \phi_0^2, \psi_0 \rangle_m,
 \end{aligned}$$

here the last inequality follows from $\|u_f(k\delta + u)\|_\infty \leq \|-\log q_{k\delta+u}\|_\infty \leq \|-\log q_{k\delta}\|_\infty$ and the fact $z \rightarrow e(x, z)$ is increasing. It is easy to see that

the function $e(x, z) \downarrow 0$ as $z \downarrow 0$. Thus, as $k \rightarrow \infty$,

$$\frac{I_2}{\langle u_f(k\delta), \psi_0 \rangle_m^2} \leq \delta(1 + c_f(k\delta))^2 \langle e(\cdot, \|\cdot\| - \log q_{k\delta}) \phi_0^2, \psi_0 \rangle_m \rightarrow 0$$

uniformly in $f \in \mathcal{C}_1^+ \setminus \{0\}$. By (3.30), we have

$$(3.32) \quad \lim_{k \rightarrow \infty} \frac{I_2}{\langle u_f(k\delta), \psi_0 \rangle_m \langle u_f((k+1)\delta), \psi_0 \rangle_m} = 0$$

uniformly in $f \in \mathcal{C}_1^+ \setminus \{0\}$. Using (3.31) and (3.32), we get,

$$\lim_{k \rightarrow \infty} \frac{\langle u_f(k\delta), \psi_0 \rangle_m - \langle u_f((k+1)\delta), \psi_0 \rangle_m}{\langle u_f((k+1)\delta), \psi_0 \rangle_m \langle u_f(k\delta), \psi_0 \rangle_m} = \delta\nu$$

uniformly in $f \in \mathcal{C}_1^+ \setminus \{0\}$. Now, (3.27) follows immediately. □

Proof of Theorem 1.5. For $t > 0$, we have

$$(3.33) \quad \mathbb{P}_\mu(\|X_t\| \neq 0) = \lim_{\theta \rightarrow \infty} (1 - \exp\{-\langle u_\theta(t), \mu \rangle\}).$$

Using Lemma 3.5 with $\delta = 1$, we have

$$(3.34) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \left(\frac{1}{\langle u_\theta(n), \psi_0 \rangle_m} - \frac{1}{\theta \langle 1, \psi_0 \rangle_m} \right) = \nu$$

uniformly in $\theta > 0$. For $\theta > 1$, it holds that

$$(3.35) \quad \frac{1}{n} \frac{1}{\theta \langle 1, \psi_0 \rangle_m} \leq \frac{1}{n} \frac{1}{\langle 1, \psi_0 \rangle_m} \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

uniformly in $\theta > 1$. It follows from (3.34) and (3.35) that

$$(3.36) \quad \lim_{n \rightarrow \infty} n \langle u_\theta(n), \psi_0 \rangle_m = \nu^{-1}$$

uniformly in $\theta > 1$. By (3.19) and (3.20), we have, as $n \rightarrow \infty$, for any $\mu(E) \leq M$,

$$\begin{aligned} & n |\langle u_\theta(n), \mu \rangle - \langle u_\theta(n), \psi_0 \rangle_m \langle \phi_0, \mu \rangle| \\ &= n \langle u_\theta(n), \psi_0 \rangle_m |\langle h_\theta(n) \phi_0, \mu \rangle| \\ &\leq M \|h_\theta(n)\|_\infty \phi_0 \|n \langle u_\theta(n), \psi_0 \rangle_m\| \rightarrow 0, \end{aligned}$$

uniformly in $\theta > 1$. Thus,

$$(3.37) \quad \lim_{n \rightarrow \infty} n \langle u_\theta(n), \mu \rangle = \nu^{-1} \langle \phi_0, \mu \rangle$$

uniformly in $\theta > 1$ and μ with $\mu(E) \leq M$. By (3.21), we have $\langle u_\theta(n), \mu \rangle \leq \langle -\log q_n, \mu \rangle \leq \|-\log q_n\|_\infty \|\mu\| \rightarrow 0$, as $n \rightarrow \infty$, uniformly in $\theta > 0$ and μ with $\mu(E) \leq M$. Therefore, it follows from (3.37) that

$$\lim_{n \rightarrow \infty} n(1 - \exp\{-\langle u_\theta(n), \mu \rangle\}) = \nu^{-1} \langle \phi_0, \mu \rangle$$

uniformly in $\theta > 1$ and μ with $\mu(E) \leq M$. Hence by (3.33), we have

$$\lim_{n \rightarrow \infty} n\mathbb{P}_\mu(\|X_n\| \neq 0) = \nu^{-1} \langle \phi_0, \mu \rangle,$$

uniformly in μ with $\mu(E) \leq M$. Since $\mathbb{P}_\mu(\|X_t\| \neq 0)$ is decreasing in t , we have

$$[t]\mathbb{P}_\mu(\|X_{([t]+1)}\| \neq 0) \leq t\mathbb{P}_\mu(\|X_t\| \neq 0) \leq ([t] + 1)\mathbb{P}_\mu(\|X_{[t]}\| \neq 0).$$

Now (1.23) follows immediately. □

Now we are ready to prove Theorem 1.6.

Proof of Theorem 1.6. First, we consider the special case when $f(x) = \phi_0(x)$. We only need to show that, for any $\lambda > 0$,

$$(3.38) \quad \mathbb{P}_\mu(\exp\{-\lambda t^{-1} \langle \phi_0, X_t \rangle\} \|X_t\| \neq 0) \rightarrow \frac{1}{\lambda\nu + 1}, \quad \text{as } t \rightarrow \infty.$$

Note that

$$\begin{aligned} & \mathbb{P}_\mu(\exp\{-\lambda t^{-1} \langle \phi_0, X_t \rangle\} \|X_t\| \neq 0) \\ &= \frac{\mathbb{P}_\mu(\exp\{-\lambda t^{-1} \langle \phi_0, X_t \rangle\}) - \mathbb{P}_\mu(\|X_t\| = 0)}{\mathbb{P}_\mu(\|X_t\| \neq 0)} \\ &= 1 - \frac{1 - \mathbb{P}_\mu(\exp\{-\lambda t^{-1} \langle \phi_0, X_t \rangle\})}{\mathbb{P}_\mu(\|X_t\| \neq 0)}. \end{aligned}$$

By Theorem 1.5, to prove (3.38), it suffices to show that, as $t \rightarrow \infty$,

$$(3.39) \quad \begin{aligned} & t(1 - \mathbb{P}_\mu(\exp\{-\lambda t^{-1} \langle \phi_0, X_t \rangle\})) \\ &= t(1 - \exp\{-\langle u_{\lambda t^{-1} \phi_0}(t), \mu \rangle\}) \rightarrow \frac{\lambda}{\lambda\nu + 1} \langle \phi_0, \mu \rangle. \end{aligned}$$

Since X_t is right continuous and ϕ_0 is a bounded continuous function, $t \rightarrow \mathbb{P}_\mu(\exp\{-\lambda t^{-1} \langle \phi_0, X_t \rangle\})$ is a right continuous function. By the Croft–Kingman lemma (see, for example, [2, Section 6.5]), it suffices to show that, for every $\delta > 0$, (3.39) holds for every sequence $n\delta$ as $n \rightarrow \infty$. For this, it is enough to prove that for any $\delta > 0$, as $n \rightarrow \infty$,

$$(3.40) \quad n\delta \langle u_{\lambda(n\delta)^{-1} \phi_0}(n\delta), \mu \rangle \rightarrow \frac{\lambda}{\lambda\nu + 1} \langle \phi_0, \mu \rangle.$$

By Lemma 3.5, we have

$$\begin{aligned} & \lim_{n \rightarrow \infty} \frac{1}{(n\delta) \langle u_{\lambda(n\delta)^{-1} \phi_0}(n\delta), \psi_0 \rangle_m} \\ &= \lim_{n \rightarrow \infty} \frac{1}{n\delta} \left(\frac{1}{\langle u_{\lambda(n\delta)^{-1} \phi_0}(n\delta), \psi_0 \rangle_m} - \frac{1}{\langle \lambda(n\delta)^{-1} \phi_0, \psi_0 \rangle_m} \right) + \frac{1}{\lambda} \\ &= \nu + \lambda^{-1}, \end{aligned}$$

which implies that

$$(3.41) \quad (n\delta)\langle u_{\lambda(n\delta)^{-1}\phi_0}(n\delta), \psi_0 \rangle_m \rightarrow \frac{\lambda}{\lambda\nu + 1}, \quad \text{as } n \rightarrow \infty.$$

Using Lemma 3.4 and (3.41), we get that, as $n \rightarrow \infty$,

$$(3.42) \quad n\delta \left| \langle u_{\lambda(n\delta)^{-1}\phi_0}(n\delta), \mu \rangle - \langle u_{\lambda(n\delta)^{-1}\phi_0}(n\delta), \psi_0 \rangle_m \langle \phi_0, \mu \rangle \right| \\ \leq n\delta \left\| h_{\lambda(n\delta)^{-1}\phi_0}(n\delta) \right\|_\infty \langle u_{\lambda(n\delta)^{-1}\phi_0}(n\delta), \psi_0 \rangle_m \langle \phi_0, \mu \rangle \rightarrow 0.$$

Now (3.40) follows easily from (3.41) and (3.42).

For a general f , let

$$(3.43) \quad \tilde{f}(x) = f(x) - \langle f, \psi_0 \rangle_m \phi_0(x).$$

Then, $\langle \tilde{f}, \psi_0 \rangle_m = 0$. It is clear that

$$(3.44) \quad \mathbb{P}_\mu((t^{-1}\langle \tilde{f}, X_t \rangle)^2 \mid \|X_t\| \neq 0) = \frac{\mathbb{P}_\mu(\langle \tilde{f}, X_t \rangle^2)}{t^2 \mathbb{P}_\mu(\|X_t\| \neq 0)}.$$

By the branching property and (2.17), we have,

$$\sup_{t>2} \text{Var}_\mu \langle \tilde{f}, X_t \rangle = \sup_{t>2} \langle \text{Var}_\delta \langle \tilde{f}, X_t \rangle, \mu \rangle < \infty.$$

It follows from (2.12) that

$$\sup_{t>1} |\mathbb{P}_\mu \langle \tilde{f}, X_t \rangle| = \sup_{t>1} |\langle T_t \tilde{f}, \mu \rangle| < \infty.$$

Combining the last two displays, we get that $\sup_{t>2} \mathbb{P}_\mu(\langle \tilde{f}, X_t \rangle)^2 < \infty$. Thus by (1.23) and (3.44), we get that as $t \rightarrow \infty$,

$$\mathbb{P}_\mu((t^{-1}\langle \tilde{f}, X_t \rangle)^2 \mid \|X_t\| \neq 0) \rightarrow 0,$$

which implies that, for any $\varepsilon > 0$,

$$(3.45) \quad \lim_{t \rightarrow \infty} \mathbb{P}_{t,\mu}(|t^{-1}\langle \tilde{f}, X_t \rangle| \geq \varepsilon) = 0.$$

Thus, by (3.43), we have

$$t^{-1}\langle f, X_t \rangle \Big|_{\mathbb{P}_{t,\mu}} \xrightarrow{d} \langle f, \psi_0 \rangle_m W. \quad \square$$

Proof of Corollary 1.8. Recall that for $f \in \mathcal{C}_2$, \tilde{f} was defined in (3.43). Thus

$$\frac{\langle f, X_t \rangle}{\langle \phi_0, X_t \rangle} - \langle f, \psi_0 \rangle_m = \frac{\langle \tilde{f}, X_t \rangle}{\langle \phi_0, X_t \rangle}.$$

For any $\varepsilon > 0$ and $\delta > 0$, by (3.45) and (1.24), we have,

$$\mathbb{P}_\mu \left(\frac{|\langle \tilde{f}, X_t \rangle|}{\langle \phi_0, X_t \rangle} > \varepsilon \mid \|X_t\| \neq 0 \right) \\ \leq \mathbb{P}_\mu(t^{-1}|\langle \tilde{f}, X_t \rangle| > \delta \mid \|X_t\| \neq 0) + \mathbb{P}_\mu(t^{-1}\langle \phi_0, X_t \rangle < \delta/\varepsilon \mid \|X_t\| \neq 0) \\ \rightarrow 0 + P(W < \delta/\varepsilon), \quad \text{as } t \rightarrow \infty.$$

Letting $\delta \rightarrow 0$, we get that

$$\lim_{t \rightarrow \infty} \mathbb{P}_{t,\mu} \left(\frac{|\langle \tilde{f}, X_t \rangle|}{\langle \phi_0, X_t \rangle} > \varepsilon \right) = 0,$$

which implies (1.26). All real-valued continuous functions with compact support in E belong to \mathcal{C}_2 . Thus, by (1.26), we have that for any real-valued continuous function f with compact support,

$$(3.46) \quad \frac{\langle f, X_t \rangle}{\langle \phi_0, X_t \rangle} \Big|_{\mathbb{P}_{t,\mu}} \xrightarrow{d} l(f) \quad (= \langle f, \psi_0 \rangle_m).$$

Hence by [24, Theorem 16.16], we get that

$$\frac{X_t}{\langle \phi_0, X_t \rangle} \Big|_{\mathbb{P}_{t,\mu}} \xrightarrow{d} l.$$

Since $\nu \rightarrow \rho(\nu, l) \wedge 1$ is a bounded continuous function on the space of Radon measures on E equipped with the vague topology, we have

$$\lim_{t \rightarrow \infty} \mathbb{P}_{t,\mu} \left[\rho \left(\frac{X_t}{\langle \phi_0, X_t \rangle}, l \right) \wedge 1 \right] = 0,$$

from which the last assertion of the corollary follows immediately. □

3.2. Proof of Theorem 1.9. In this subsection, we give the proof of Theorem 1.9. We prove a simple lemma first.

LEMMA 3.6. *Suppose that \mathcal{V} is an index set and $\{F_v : v \in \mathcal{V}\}$ is a family of uniformly bounded random variables, that is, there is a constant M such that $|F_v| \leq M$ for all $v \in \mathcal{V}$, then any $s > 0$,*

$$(3.47) \quad \lim_{t \rightarrow \infty} \sup_{v \in \mathcal{V}} |\mathbb{P}_{t+s,\mu}(F_v) - \mathbb{P}_{t,\mu}(F_v)| = 0.$$

Proof. By Theorem 1.5, we have

$$(3.48) \quad \lim_{t \rightarrow \infty} \frac{\mathbb{P}_\mu(\|X_t\| \neq 0)}{\mathbb{P}_\mu(\|X_{t+s}\| \neq 0)} = 1.$$

By the definition of $\mathbb{P}_{t,\mu}$, we have

$$\begin{aligned} & \mathbb{P}_{t+s,\mu}(F_v) \\ &= \mathbb{P}_{t,\mu}(F_v, \|X_{t+s}\| \neq 0) \frac{\mathbb{P}_\mu(\|X_t\| \neq 0)}{\mathbb{P}_\mu(\|X_{t+s}\| \neq 0)} \\ &= \mathbb{P}_{t,\mu}(F_v) \frac{\mathbb{P}_\mu(\|X_t\| \neq 0)}{\mathbb{P}_\mu(\|X_{t+s}\| \neq 0)} - \mathbb{P}_{t,\mu}(F_v, \|X_{t+s}\| = 0) \frac{\mathbb{P}_\mu(\|X_t\| \neq 0)}{\mathbb{P}_\mu(\|X_{t+s}\| \neq 0)}. \end{aligned}$$

Thus, as $t \rightarrow \infty$,

$$\begin{aligned} & |\mathbb{P}_{t+s,\mu}(F_v) - \mathbb{P}_{t,\mu}(F_v)| \\ & \leq M \left| \frac{\mathbb{P}_\mu(\|X_t\| \neq 0)}{\mathbb{P}_\mu(\|X_{t+s}\| \neq 0)} - 1 \right| + M \mathbb{P}_{t,\mu}(\|X_{t+s}\| = 0) \frac{\mathbb{P}_\mu(\|X_t\| \neq 0)}{\mathbb{P}_\mu(\|X_{t+s}\| \neq 0)} \end{aligned}$$

$$\begin{aligned}
 &= M \left| \frac{\mathbb{P}_\mu(\|X_t\| \neq 0)}{\mathbb{P}_\mu(\|X_{t+s}\| \neq 0)} - 1 \right| + M \frac{\mathbb{P}_\mu(\|X_{t+s}\| = 0, \|X_t\| \neq 0)}{\mathbb{P}_\mu(\|X_t\| \neq 0)} \frac{\mathbb{P}_\mu(\|X_t\| \neq 0)}{\mathbb{P}_\mu(\|X_{t+s}\| \neq 0)} \\
 &= M \left| \frac{\mathbb{P}_\mu(\|X_t\| \neq 0)}{\mathbb{P}_\mu(\|X_{t+s}\| \neq 0)} - 1 \right| + M \frac{\mathbb{P}_\mu(\|X_t\| \neq 0) - \mathbb{P}_\mu(\|X_{t+s}\| \neq 0)}{\mathbb{P}_\mu(\|X_{t+s}\| \neq 0)} \\
 &= 2M \left| \frac{\mathbb{P}_\mu(\|X_t\| \neq 0)}{\mathbb{P}_\mu(\|X_{t+s}\| \neq 0)} - 1 \right| \rightarrow 0. \quad \square
 \end{aligned}$$

We now recall some facts about weak convergence which will be used later. For $f : \mathbb{R}^d \rightarrow \mathbb{R}$, let $\|f\|_L := \sup_{x \neq y} |f(x) - f(y)|/\|x - y\|$ and $\|f\|_{BL} := \|f\|_\infty + \|f\|_L$. For any probability measures ν_1 and ν_2 on \mathbb{R}^d , define

$$\beta(\nu_1, \nu_2) := \sup \left\{ \left| \int f d\nu_1 - \int f d\nu_2 \right| : \|f\|_{BL} \leq 1 \right\}.$$

Then β is a metric. It follows from [14, Theorem 11.3.3] that the topology generated by β is equivalent to the weak convergence topology. From the definition, we can easily see that, if ν_1 and ν_2 are the distributions of two \mathbb{R}^d -valued random variables X and Y respectively, then

$$(3.49) \quad \beta(\nu_1, \nu_2) \leq E\|X - Y\| \leq \sqrt{E\|X - Y\|^2}.$$

The following simple fact will be used several times later in this section:

$$(3.50) \quad \left| e^{ix} - \sum_{m=0}^n \frac{(ix)^m}{m!} \right| \leq \min \left(\frac{|x|^{n+1}}{(n+1)!}, \frac{2|x|^n}{n!} \right).$$

Now we are ready to prove Theorem 1.9.

Proof of Theorem 1.9. Define an \mathbb{R}^2 -valued random variable:

$$U_1(t) := (t^{-1}\langle \phi_0, X_t \rangle, t^{-1/2}\langle f, X_t \rangle).$$

We need to prove that, conditioning on $\|X_t\| \neq 0$, $U_1(t)$ converges to $(W, G(f)\sqrt{W})$ in distribution as $t \rightarrow \infty$, which is equivalent to proving that, when one lets t tend to ∞ first and then lets s tend to ∞ ,

$$(3.51) \quad U_1(t+s)|_{\mathbb{P}_{t+s, \mu}} \xrightarrow{d} (W, G(f)\sqrt{W}).$$

Before we prove (3.51), we first give the main idea of the proof. In Theorem 1.6, we have proved that the first component of $U_1(t)$ converges to W . So the key is the second component. If we condition on X_t , the mean of $\langle f, X_{s+t} \rangle$ is $\langle T_s f, X_t \rangle$. Let us consider the centered random variable $\langle f, X_{s+t} \rangle - \langle T_s f, X_t \rangle$. For fixed $s > 0$, as $t \rightarrow \infty$, since the ‘infinitesimal particles’ evolve independently after time t , it is reasonable to expect that, conditioning on X_t and $\|X_t\| \neq 0$, $(\langle f, X_{s+t} \rangle - \langle T_s f, X_t \rangle) / \sqrt{\langle \text{Var} \langle f, X_{s+t} \rangle | X_t \rangle}$ converges in distribution to a standard normal random variable. Note that $\text{Var}(\langle f, X_{s+t} \rangle | X_t) = \langle \text{Var}_\delta \langle f, X_s \rangle, X_t \rangle$. By Theorem 1.6, we have

$t^{-1} \text{Var}(\langle f, X_{s+t} \rangle | X_t) \xrightarrow{d} \langle \text{Var}_\delta \langle f, X_s \rangle, \psi_0 \rangle_m W$, as $t \rightarrow \infty$. We may thus conclude that $t^{-1/2}(\langle f, X_{s+t} \rangle - \langle T_s f, X_t \rangle) \xrightarrow{d} \sqrt{W} G_s$, where $G_s \sim \mathcal{N}(0, \sigma_f^2(s))$ with $\sigma_f^2(s) = \langle \text{Var}_\delta \langle f, X_s \rangle, \psi_0 \rangle_m$ and W is the random variable defined in Theorem 1.6.

The above analysis suggests that we should first consider another \mathbb{R}^2 -valued random variable $U_2(s, t)$ defined by

$$U_2(s, t) = (t^{-1} \langle \phi_0, X_t \rangle, t^{-1/2} (\langle f, X_{s+t} \rangle - \langle T_s f, X_t \rangle)), \quad s, t > 2.$$

We claim that,

$$(3.52) \quad U_2(s, t) |_{\mathbb{P}_{t,\mu}} \xrightarrow{d} (W, \sqrt{W} G_s), \quad \text{as } t \rightarrow \infty.$$

We will leave the proof of (3.52) to the end of the proof of this theorem.

Define

$$U_3(s, t) := ((t+s)^{-1} \langle \phi_0, X_t \rangle, (t+s)^{-1/2} (\langle f, X_{s+t} \rangle - \langle T_s f, X_t \rangle)).$$

By (3.52), we have

$$(3.53) \quad U_3(s, t) |_{\mathbb{P}_{t,\mu}} \xrightarrow{d} (W, \sqrt{W} G_s),$$

as $t \rightarrow \infty$. It follows from (2.11) and (1.23) that, as $t \rightarrow \infty$,

$$\begin{aligned} (t+s)^{-2} \mathbb{P}_{t,\mu} (\langle \phi_0, X_{t+s} \rangle - \langle \phi_0, X_t \rangle)^2 &= \frac{\mathbb{P}_\mu (\langle \phi_0, X_{t+s} \rangle - \langle \phi_0, X_t \rangle)^2}{(t+s)^2 \mathbb{P}_\mu (\|X_t\| \neq 0)} \\ &= \frac{\mathbb{P}_\mu (\langle \text{Var}_\delta \langle \phi_0, X_s \rangle, X_t \rangle)}{(t+s)^2 \mathbb{P}_\mu (\|X_t\| \neq 0)} \rightarrow 0. \end{aligned}$$

If we put

$$U_4(s, t) := ((t+s)^{-1} \langle \phi_0, X_{t+s} \rangle, (t+s)^{-1/2} (\langle f, X_{s+t} \rangle - \langle T_s f, X_t \rangle)),$$

then, as $t \rightarrow \infty$,

$$(3.54) \quad U_4(s, t) |_{\mathbb{P}_{t,\mu}} \xrightarrow{d} (W, \sqrt{W} G_s).$$

By Lemma 3.6, we have

$$\begin{aligned} &\lim_{t \rightarrow \infty} \mathbb{P}_{t+s,\mu} (\exp\{i\theta_1 (t+s)^{-1} \langle \phi_0, X_{t+s} \rangle \\ &\quad + i\theta_2 (t+s)^{-1/2} (\langle f, X_{s+t} \rangle - \langle T_s f, X_t \rangle)\}) \\ &= \lim_{t \rightarrow \infty} \mathbb{P}_{t,\mu} (\exp\{i\theta_1 (t+s)^{-1} \langle \phi_0, X_{t+s} \rangle \\ &\quad + i\theta_2 (t+s)^{-1/2} (\langle f, X_{s+t} \rangle - \langle T_s f, X_t \rangle)\}) \\ &= P(\exp\{i\theta_1 W + i\theta_2 \sqrt{W} G_s\}). \end{aligned}$$

Thus, we have

$$(3.55) \quad U_4(s, t) |_{\mathbb{P}_{t+s,\mu}} \xrightarrow{d} (W, \sqrt{W} G_s), \quad \text{as } t \rightarrow \infty.$$

Now, we deal with $J_2(t, s) := \frac{\langle T_s f, X_t \rangle}{(t+s)^{1/2}}$. We claim that

$$(3.56) \quad \lim_{s \rightarrow \infty} \limsup_{t \rightarrow \infty} \mathbb{P}_{t+s, \delta_x} (|J_2(t, s)|^2) = 0.$$

By (2.11), we have that $\mathbb{P}_\mu \langle T_s f, X_t \rangle = \langle T_{t+s} f, \mu \rangle \rightarrow 0$, as $t \rightarrow \infty$. Thus by (1.23) and (2.16), we have

$$(3.57) \quad \begin{aligned} & \limsup_{t \rightarrow \infty} \mathbb{P}_{t+s, \delta_x} (|J_2(t, s)|^2) \\ &= \limsup_{t \rightarrow \infty} \frac{\mathbb{P}_\mu (\langle T_s f, X_t \rangle^2, \|X_{t+s}\| \neq 0)}{(t+s) \mathbb{P}_\mu (\|X_{t+s}\| \neq 0)} \\ &\leq \limsup_{t \rightarrow \infty} \frac{\mathbb{P}_\mu (\langle T_s f, X_t \rangle^2)}{(t+s) \mathbb{P}_\mu (\|X_{t+s}\| \neq 0)} = \nu \sigma_{(T_s f)}^2. \end{aligned}$$

It follows from (1.27) that, as $s \rightarrow \infty$,

$$\sigma_{(T_s f)}^2 = \int_s^\infty \langle A(T_u f)^2, \psi_0 \rangle_m du \rightarrow 0.$$

Now (3.56) follows immediately.

By (2.16), we have $\lim_{s \rightarrow \infty} \text{Var}_{\delta_x} \langle f, X_s \rangle = \sigma_f^2 \phi_1(x)$, thus $\lim_{s \rightarrow \infty} \sigma_f^2(s) = \sigma_f^2$. Hence,

$$(3.58) \quad \lim_{s \rightarrow \infty} \beta(G_s, G(f)) = 0.$$

Let $\mathcal{D}(s+t)$ and $\tilde{\mathcal{D}}(s, t)$ be the distributions of $U_1(s+t)$ and $U_4(s, t)$ under $\mathbb{P}_{t+s, \mu}$ respectively, and let $\hat{\mathcal{D}}(s)$ and \mathcal{D} be the distributions of $(W, \sqrt{W}G_s)$ and $(W, \sqrt{W}G(f))$, respectively. Then, using (3.49), we have

$$(3.59) \quad \begin{aligned} & \limsup_{t \rightarrow \infty} \beta(\mathcal{D}(s+t), \mathcal{D}) \\ &\leq \limsup_{t \rightarrow \infty} [\beta(\mathcal{D}(s+t), \tilde{\mathcal{D}}(s, t)) + \beta(\tilde{\mathcal{D}}(s, t), \hat{\mathcal{D}}(s)) + \beta(\hat{\mathcal{D}}(s), \mathcal{D})] \\ &\leq \limsup_{t \rightarrow \infty} \sqrt{\mathbb{P}_{t+s, \mu} ((t+s)^{-1} \langle T_s f, X_t \rangle^2)} + 0 + \beta(\hat{\mathcal{D}}(s), \mathcal{D}). \end{aligned}$$

Then we have

$$\begin{aligned} \limsup_{t \rightarrow \infty} \beta(\mathcal{D}(t), \mathcal{D}) &= \limsup_{t \rightarrow \infty} \beta(\mathcal{D}(s+t), \mathcal{D}) \\ &\leq \limsup_{t \rightarrow \infty} \sqrt{\mathbb{P}_{t+s, \mu} (J_2(s, t)^2)} + \beta(\hat{\mathcal{D}}(s), \mathcal{D}). \end{aligned}$$

Letting $s \rightarrow \infty$, by (3.56) and (3.58), we get

$$\limsup_{t \rightarrow \infty} \beta(\mathcal{D}(t), \mathcal{D}) = 0,$$

which implies the result of theorem.

Now we prove (3.52).

Denote the characteristic function of $U_2(s, t)$ under $\mathbb{P}_{t, \mu}$ by $\kappa_1(\theta_1, \theta_2, s, t)$:

$$\begin{aligned}
 (3.60) \quad \kappa_1(\theta_1, \theta_2, s, t) &= \mathbb{P}_{t, \mu}(\exp\{i\theta_1 t^{-1}\langle \phi_0, X_t \rangle + i\theta_2 t^{-1/2}(\langle f, X_{s+t} \rangle - \langle T_s f, X_t \rangle)\}) \\
 &= \mathbb{P}_{t, \mu}\left(\exp\left\{i\theta_1 t^{-1}\langle \phi_0, X_t \rangle\right.\right. \\
 &\quad \left.\left.+ \int_E \int_{\mathbb{D}} (e^{i\theta_2 t^{-1/2}\langle f, \omega_s \rangle} - 1 - i\theta_2 t^{-1/2}\langle f, \omega_s \rangle) \mathbb{N}_x(d\omega) X_t(dx)\right\}\right),
 \end{aligned}$$

where in the last equality we used the Markov property of X , (2.27) and (2.25). Define

$$J_s(\theta, x) := \int_{\mathbb{D}} (\exp\{i\theta f, \omega_s\} - 1 - i\theta\langle f, \omega_s \rangle) \mathbb{N}_x(d\omega)$$

and

$$I_s(\theta, x) := \int_{\mathbb{D}} \left(\exp\{i\theta f, \omega_s\} - 1 - i\theta\langle f, \omega_s \rangle + \frac{1}{2}\theta^2\langle f, \omega_s \rangle^2\right) \mathbb{N}_x(d\omega).$$

Let $V_s(x) = \mathbb{V}ar_{\delta_x}\langle f, X_s \rangle \in \mathcal{C}_2^+$. Then, by (2.26), we have

$$\begin{aligned}
 J_s(\theta, x) &= -\frac{1}{2}\theta^2 V_s(x) + I_s(\theta, x) \\
 &= -\frac{1}{2}\theta^2 \langle V_s, \psi_0 \rangle_m \phi_0(x) - \frac{1}{2}\theta^2 \tilde{V}_s(x) + I_s(\theta, x),
 \end{aligned}$$

where $\tilde{V}_s = V_s - \langle V_s, \psi_0 \rangle_m \phi_0(x) \in \mathcal{C}_2$. Thus, we have

$$\begin{aligned}
 (3.61) \quad &i\theta_1 t^{-1}\langle \phi_0, X_t \rangle + \langle J_s(t^{-1/2}\theta_2, \cdot), X_t \rangle \\
 &= \left(i\theta_1 - \frac{1}{2}\theta_2^2 \langle V_s, \psi_0 \rangle_m\right) t^{-1}\langle \phi_0, X_t \rangle \\
 &\quad - \frac{1}{2}\theta_2^2 t^{-1}\langle \tilde{V}_s, X_t \rangle + \langle I_s(t^{-1/2}\theta_2, \cdot), X_t \rangle.
 \end{aligned}$$

By (3.45), we know that, for any $\varepsilon > 0$,

$$(3.62) \quad \lim_{t \rightarrow \infty} \mathbb{P}_{t, \mu}(|t^{-1}\langle \tilde{V}_s, X_t \rangle| \geq \varepsilon) = 0.$$

By (3.50), we have

$$(3.63) \quad |I_s(t^{-1/2}\theta_2, x)| \leq \theta_2^2 t^{-1} \mathbb{N}_x\left(\langle f, \omega_s \rangle^2 \left(\frac{t^{-1/2}\theta_2 \langle f, \omega_s \rangle}{6} \wedge 1\right)\right).$$

Let

$$h(x, s, t) = \mathbb{N}_x\left(\langle f, \omega_s \rangle^2 \left(\frac{t^{-1/2}\theta_2 \langle f, \omega_s \rangle}{6} \wedge 1\right)\right).$$

We note that $h(x, s, t) \downarrow 0$ as $t \uparrow \infty$. By (2.17), we have

$$h(x, s, t) \leq \mathbb{N}_x(\langle f, X_s \rangle^2) = \mathbb{V}ar_{\delta_x}\langle f, X_s \rangle \lesssim \phi_0(x) \in \mathcal{C}_2.$$

Thus, by (1.23) and (2.11), we have, for any $u < t$,

$$\begin{aligned} t^{-1}\mathbb{P}_{t,\mu}\langle h(\cdot, s, t), X_t \rangle &\leq t^{-1}\mathbb{P}_{t,\mu}\langle h(\cdot, s, u), X_t \rangle \\ &= \frac{\mathbb{P}_\mu\langle h(\cdot, s, u), X_t \rangle}{t\mathbb{P}_\mu(\|X_t\| \neq 0)} \rightarrow \nu\langle h(\cdot, s, u), \psi_0 \rangle_m, \end{aligned}$$

as $t \rightarrow \infty$. Letting $u \rightarrow \infty$, we get $\langle h(\cdot, s, u), \psi_0 \rangle_m \rightarrow 0$. Thus, by (3.63), we get that

$$\lim_{t \rightarrow \infty} \mathbb{P}_{t,\mu} \left| \langle I_s(t^{-1/2}\theta_2, \cdot), X_t \rangle \right| = 0,$$

which implies that, for any $\varepsilon > 0$,

$$(3.64) \quad \lim_{t \rightarrow \infty} \mathbb{P}_{t,\mu} \left(\left| \langle I_s(t^{-1/2}\theta_2, \cdot), X_t \rangle \right| \geq \varepsilon \right) = 0.$$

Thus, by (3.62), (3.64) and (3.61), we get

$$i\theta_1 t^{-1} \langle \phi_0, X_t \rangle + \langle J_s(t^{-1/2}\theta_2, \cdot), X_t \rangle \Big|_{\mathbb{P}_{t,\mu}} \xrightarrow{d} \left(i\theta_1 - \frac{1}{2}\theta_2^2 \langle V_s, \psi_0 \rangle_m \right) W.$$

Since the real part of $J_s(t^{-1/2}\theta_2, x)$ is non-positive, we have

$$\left| \exp \left\{ i\theta_1 t^{-1} \langle \phi_0, X_t \rangle + \langle J_s(t^{-1/2}\theta_2, \cdot), X_t \rangle \right\} \right| \leq 1.$$

Therefore, by (3.60) and the dominated convergence theorem, we get

$$\lim_{t \rightarrow \infty} \kappa_1(\theta_1, \theta_2, s, t) = P \left(\exp \left\{ \left(i\theta_1 - \frac{1}{2}\theta_2^2 \langle V_s, \psi_0 \rangle_m \right) W \right\} \right),$$

which implies our claim (3.52). \square

REFERENCES

- [1] R. Adamczak and P. Miłoś, *CLT for Ornstein–Uhlenbeck branching particle system*, Electron. J. Probab. **20** (2015), Art. ID 42. MR 3339862
- [2] W. J. Anderson, *Continuous time Markov chains*, Springer-Verlag, New York, 1991. MR 1118840
- [3] S. Asmussen and H. Hering, *Branching processes*, Birkhäuser, Boston, MA, 1983. MR 0701538
- [4] K. Athreya and P. Ney, *Branching processes*, Springer-Verlag, New York, 1972. MR 0373040
- [5] K. Athreya and P. Ney, *Functionals of critical multitype branching processes*, Ann. Probab. **2** (1974), 339–343. MR 0356264
- [6] R. Banuelos, *Intrinsic ultracontractivity and eigenfunction estimates for Schrödinger operators*, J. Funct. Anal. **100** (1991), 181–206. MR 1124298
- [7] N. Champagnat and S. Roelly, *Limit theorems for conditioned multitype Dawson–Watanabe processes and Feller diffusions*, Electron. J. Probab. **13** (2008), 777–810. MR 2399296
- [8] Z.-Q. Chen, P. Kim and R. Song, *Dirichlet heat kernel estimates for fractional Laplacian with gradient perturbation*, Ann. Probab. **40** (2012), 2483–2538. MR 3050510
- [9] Z.-Q. Chen, P. Kim and R. Song, *Dirichlet heat kernel estimates for rotationally symmetric Levy processes*, Proc. Lond. Math. Soc. (3) **109** (2014), 90–120. MR 3237737

- [10] Z.-Q. Chen, P. Kim and R. Song, *Dirichlet heat kernel estimates for subordinate Brownian motions with Gaussian components*, J. Reine Angew. Math. **711** (2016), 111–138.
- [11] Z.-Q. Chen, Y.-X. Ren, R. Song and R. Zhang, *Strong law of large numbers for supercritical superprocesses under second moment condition*, Front. Math. China **10** (2015), 807–838. MR 3352888
- [12] E. B. Davies and B. Simon, *Ultracontractivity and the heat kernel for Schrödinger operators and Dirichlet Laplacians*, J. Funct. Anal. **59** (1984), 335–395. MR 0766493
- [13] D. A. Dawson, *Measure-valued Markov processes*, Springer-Verlag, Berlin, 1993. MR 1242575
- [14] R. M. Dudley, *Real analysis and probability*, Cambridge University Press, Cambridge, 2002. MR 1932358
- [15] E. B. Dynkin, *Superprocesses and partial differential equations*, Ann. Probab. **21** (1993), 1185–1262. MR 1235414
- [16] E. B. Dynkin and S. E. Kuznetsov, *N-measure for branching exit Markov system and their applications to differential equations*, Probab. Theory Related Fields **130** (2004), 135–150. MR 2092876
- [17] N. El Karoui and S. Roelly, *Propriétés de martingales, explosion et représentation de Lévy–Khintchine d’une classe de processus de branchement à valeurs mesures*, Stochastic Process. Appl. **38** (1991), 239–266. MR 1119983
- [18] J. Engländer, Y.-X. Ren and R. Song, *Weak extinction versus global exponential growth of total mass for superdiffusions corresponding to the operator $Lu + \beta u - ku^2$* , Ann. Inst. Henri Poincaré Probab. Stat. **52** (2016), 448–482. MR 3449310
- [19] S. N. Evans and E. Perkins, *Measure-valued Markov branching processes conditioned on non-extinction*, Israel J. Math. **71** (1990), 329–337. MR 1088825
- [20] M. I. Goldstein and F. M. Hoppe, *Critical multitype branching processes with infinite variance*, J. Math. Anal. Appl. **65** (1978), 675–686. MR 0510478
- [21] T. E. Harris, *The theory of branching processes*, Springer, Berlin, 1963. MR 0163361
- [22] A. Joffe and F. Spitzer, *On multitype branching processes with $\rho \leq 1$* , J. Math. Anal. Appl. **19** (1967), 409–430. MR 0212895
- [23] K. Kaleta and T. Kulczycki, *Intrinsic ultracontractivity for Schrödinger operators based on fractional Laplacians*, Potential Anal. **33** (2010), 313–339. MR 2726901
- [24] O. Kallenberg, *Foundations of modern probability*, 2nd ed., Springer, New York, 2002. MR 1876169
- [25] H. Kesten, P. Ney and F. Spitzer, *The Galton–Watson process with mean one and finite variance*, Theory Probab. Appl. **11** (1966), 513–540.
- [26] P. Kim and R. Song, *Two-sided estimates on the density of Brownian motion with singular drift*, Illinois J. Math. **50** (2006), 635–688. MR 2247841
- [27] P. Kim and R. Song, *On dual processes of non-symmetric diffusions with measure-valued drifts*, Stochastic Process. Appl. **118** (2008), 790–817. MR 2411521
- [28] P. Kim and R. Song, *Intrinsic ultracontractivity of non-symmetric diffusion semi-groups in bounded domains*, Tohoku Math. J. (2) **60** (2008), 527–547. MR 2487824
- [29] P. Kim and R. Song, *Intrinsic ultracontractivity of non-symmetric diffusions with measure-valued drifts and potentials*, Ann. Probab. **36** (2008), 1904–1945. MR 2440927
- [30] P. Kim and R. Song, *Intrinsic ultracontractivity for non-symmetric Lévy processes*, Forum Math. **21** (2009), 43–66. MR 2494884
- [31] P. Kim and R. Song, *Stable process with singular drift*, Stochastic Process. Appl. **124** (2014), 2479–2516. MR 3192504
- [32] K. Kim, R. Song and Z. Vondracek, *Two-sided Green function estimates for killed subordinate Brownian motions*, Proc. Lond. Math. Soc. (3) **104** (2012), 927–958. MR 2928332

- [33] K. Kim, R. Song and Z. Vondracek, *Potential theory of subordinate Brownian motions with Gaussian components*, Stochastic Process. Appl. **123** (2013), 764–795. MR 3005005
- [34] A. Kolmogorov, *Zur Lösung einer biologischen Aufgabe*, Izvestiya nauchno-issledovatel'skogo instituta matematiki i mekhaniki pri Tomskom Gosudarstvennom Universitete **2** (1938), 1–6.
- [35] T. Kulczycki and B. Siudeja, *Intrinsic ultracontractivity of the Feynman–Kac semigroup for relativistic stable processes*, Trans. Amer. Math. Soc. **358** (2006), 5025–5057. MR 2231884
- [36] A. Lambert, *Quasi-stationary distributions and the continuous-state branching process conditioned to be never extinct*, Electron. J. Probab. **12** (2007), 420–446. MR 2299923
- [37] Z. Li, *Asymptotic behavior of continuous time and state branching process*, J. Austral. Math. Soc. Ser. A **68** (2000), 68–84. MR 1727226
- [38] Z. Li, *Skew convolution semigroups and related immigration processes*, Theory Probab. Appl. **46** (2003), 274–296.
- [39] Z. Li, *Measure-valued branching Markov processes*, Springer, Heidelberg, 2011. MR 2760602
- [40] R. Lyons, R. Pemantle and Y. Peres, *Conceptual proofs of $L \log L$ criteria for mean behavior of branching processes*, Ann. Probab. **23** (1995), 1125–1138. MR 1349164
- [41] P. Miłoś, *Spatial CLT for the supercritical Ornstein–Uhlenbeck superprocess*, preprint, 2012; available at [arXiv:1203.6661](https://arxiv.org/abs/1203.6661).
- [42] P. Ney, *Critical multi-type degenerate branching processes*, technical report, University of Wisconsin.
- [43] A. G. Pakes, *Critical Markov branching process limit theorems allowing infinite variance*, Adv. in Appl. Probab. **42** (2010), 460–488. MR 2675112
- [44] Y.-X. Ren, R. Song and R. Zhang, *Central limit theorems for super Ornstein–Uhlenbeck processes*, Acta Appl. Math. **130** (2014), 9–49. MR 3180938
- [45] Y.-X. Ren, R. Song and R. Zhang, *Central limit theorems for supercritical branching Markov processes*, J. Funct. Anal. **266** (2014), 1716–1756. MR 3146834
- [46] Y.-X. Ren, R. Song and R. Zhang, *Central limit theorems for supercritical superprocesses*, Stochastic Process. Appl. **125** (2015), 428–457. MR 3293289
- [47] Y.-X. Ren, R. Song and R. Zhang, *Central Limit Theorems for Supercritical Branching Non-symmetric Markov Processes*, to appear in Ann. Probab.; available at [arXiv:1404.0116](https://arxiv.org/abs/1404.0116).
- [48] H. H. Schaefer, *Banach lattices and positive operators*, Springer, New York, 1974. MR 0423039
- [49] E. Seneta, *The Galton–Watson process with mean one*, J. Appl. Probab. **4** (1967), 489–495. MR 0228075
- [50] Y.-C. Sheu, *Lifetime and compactness of range for super-Brownian motion with a general branching mechanism*, Stochastic Process. Appl. **70** (1997), 129–141. MR 1472962
- [51] R. S. Slack, *A branching process with mean one and possibly infinite variance*, Z. Wahrsch. Verw. Gebiete **9** (1968), 139–145. MR 0228077
- [52] V. A. Vatutin, *Limit theorems for critical multitype Markov branching processes with infinite second moments*, Mat. Sb. **103** (1977), 253–264. MR 0443115
- [53] V. A. Vatutin, *Linear functionals for critical multitype Galton–Watson branching processes*, J. Math. Sci. (N. Y.) **99** (2000), 1502–1509. MR 1766080
- [54] A. M. Yaglom, *Certain limit theorems of the theory of branching random processes*, Doklady Akad. Nauk SSSR (N.S.) **56** (1947), 795–798. MR 0022045
- [55] J. Zhang, S. Li and R. Song, *Quasi-stationarity and quasi-ergodicity of general Markov processes*, Sci. China Math. **57** (2014), 2013–2024. MR 3247530

YAN-XIA REN, LMAM SCHOOL OF MATHEMATICAL SCIENCES & CENTER FOR STATISTICAL SCIENCE, PEKING UNIVERSITY, BEIJING, 100871, P.R. CHINA

E-mail address: yxren@math.pku.edu.cn

RENMING SONG, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, URBANA, IL 61801, USA

E-mail address: rsong@math.uiuc.edu

RUI ZHANG, SCHOOL OF MATHEMATICAL SCIENCES, CAPITAL NORMAL UNIVERSITY, BEIJING, 100048, P.R. CHINA

E-mail address: zhangrui27@cnu.edu.cn