Topological Methods in Nonlinear Analysis Volume 53, No. 2, 2019, 575–601 DOI: 10.12775/TMNA.2019.012

© 2019 Juliusz Schauder Centre for Nonlinear Studies Nicolaus Copernicus University in Toruń

REIDEMEISTER SPECTRA FOR SOLVMANIFOLDS IN LOW DIMENSIONS

KAREL DEKIMPE — SAM TERTOOY — IRIS VAN DEN BUSSCHE

ABSTRACT. The Reidemeister number of an endomorphism of a group is the number of twisted conjugacy classes determined by that endomorphism. The collection of all Reidemeister numbers of all automorphisms of a group G is called the Reidemeister spectrum of G. In this paper, we determine the Reidemeister spectra of all fundamental groups of solvmanifolds up to Hirsch length 4.

1. Introduction

Let G be a group and $\varphi \colon G \to G$ an endomorphism. Consider the following equivalence relation on G:

 $x \sim_{\varphi} y$ if and only if $\exists z \in G : x = zy\varphi(z)^{-1}$.

The equivalence classes under \sim_{φ} are the *Reidemeister classes of* φ or the φ twisted conjugacy classes. We denote the set of these equivalence classes by $\mathcal{R}(\varphi)$. The number of equivalence classes is called the *Reidemeister number of* φ and is denoted by $R(\varphi)$. If $\mathcal{R}(\varphi)$ is infinite, we write $R(\varphi) = \infty$. Subsequently, the *Reidemeister spectrum of* G is defined as

$$\operatorname{Spec}_{\mathbf{R}}(G) := \{ R(\varphi) \mid \varphi \in \operatorname{Aut}(G) \}.$$

²⁰¹⁰ Mathematics Subject Classification. 20E36, 20F16.

 $Key\ words\ and\ phrases.$ Twisted conjugacy; Reidemeister number; polycyclic group; solvmanifold.

If $\operatorname{Spec}_{\mathrm{R}}(G) = \mathbb{N} \cup \{\infty\}$, the Reidemeister spectrum of G is said to be *full*. If $\operatorname{Spec}_{\mathrm{R}}(G) = \{\infty\}$, we say that G has the R_{∞} property.

Reidemeister numbers of morphisms of groups correspond to Reidemeister numbers of self-maps on topological spaces that play a crucial role in the Nielsen theory of fixed points and periodic points of these maps. We refer to [13], [14] and [15] for details about it. The study of groups having (or not having) the R_{∞} property was initiated by A. Fel'shtyn and R. Hill in the mid 90's and is now a very active research topic (see [3]–[9], [11], [19] for some papers in this area). In [7] it was conjectured that finitely generated, residually finite groups without the R_{∞} property must be virtually solvable. Having this in mind and also the topological meaning of a Reidemeister number (as being related to the study of fixed points), we will focus on a special class of solvable groups, namely the fundamental groups of solvmanifolds. In fact, for the rest of this paper, we will study these groups up to dimension 4. It is well known [18] that a group E is the fundamental group of a compact solvmanifold if and only if E is an extension of the form

$$(1.1) 1 \longrightarrow N \longrightarrow E \xrightarrow{p} \mathbb{Z}^k \longrightarrow 1$$

where N is a finitely generated torsion-free nilpotent group and k is a nonnegative integer (E = N when k = 0). The dimension of the corresponding compact solvmanifold is the same as the Hirsch length of the group E.

The aim of this paper is to determine the Reidemeister spectrum of all such E of Hirsch length at most 4. It turns out that many of these groups satisfy the R_{∞} property. The groups that do not satisfy the R_{∞} property nor have full spectrum (abelian groups), have either Reidemeister spectrum $\{2, \infty\}, \{4, \infty\}, \{8, \infty\}, 2\mathbb{N} \cup \{\infty\}, 4\mathbb{N} \cup \{\infty\}, 6\mathbb{N} \cup \{\infty\}$ or $8\mathbb{N} \cup \{\infty\}$.

This paper is organised in five sections. We begin the paper by recalling some formulas to compute Reidemeister numbers. In Section 3, we first focus on the special case where the group E is nilpotent. In Section 4, we consider the groups E of Hirsch length 3; in Section 5, we consider the groups E of Hirsch length 4.

2. Preliminaries

When determining the Reidemeister spectrum of the group E, an extension of N by \mathbb{Z}^k , it will be convenient that the subgroup N is characteristic. In this situation, we can easily determine Reidemeister numbers using the following *addition formula*:

LEMMA 2.1 ([8, Lemma 2.1]). Let E be an extension of some group G by \mathbb{Z}^k . Let s be any (set-theoretic) section, so we have the exact sequence with $p \circ s$ the identity on \mathbb{Z}^k :

$$1 \longrightarrow G \longrightarrow E \xrightarrow{p} \mathbb{Z}^k \longrightarrow 1.$$

Define the function $\alpha \colon \mathbb{Z}^k \to \operatorname{Aut}(G)$ by $\alpha(z)(g) = s(z) g s(z)^{-1}$. Let φ be an endomorphism of E such that $\varphi(G) \subseteq G$. Write $\varphi|_G = \varphi'$ and let $\overline{\varphi}$ denote the induced endomorphism on the quotient \mathbb{Z}^k . Suppose $\{z_i \mid i \in \mathcal{I}\}$ is a complete set of representatives for the Reidemeister classes of $\overline{\varphi}$. Then

$$R(\varphi) = \sum_{i \in \mathcal{I}} R(\alpha(z_i) \circ \varphi').$$

In particular, $R(\varphi) = \infty$ whenever $R(\overline{\varphi}) = \infty$ or $R(\varphi') = \infty$. In general, for any automorphism $\varphi \colon G \to G$ inducing $\overline{\varphi} \colon G/H \to G/H$, it always holds that if $R(\overline{\varphi}) = \infty$, then $R(\varphi) = \infty$ as well. Indeed, the map $\widehat{\pi} \colon \mathcal{R}(\varphi) \to \mathcal{R}(\overline{\varphi})$ defined by $[x]_{\varphi} \mapsto [\overline{x}]_{\overline{\varphi}}$ is well-defined and surjective. Hence, if H is a characteristic subgroup of G and G/H has the R_{∞} property, then G has the R_{∞} property as well.

Since the groups E we are interested in in this paper are built by repeated extensions of the groups \mathbb{Z}^n , the following well-known formula is pivotal:

LEMMA 2.2. Suppose $\varphi \colon \mathbb{Z}^n \to \mathbb{Z}^n$ is multiplication by $M \in \operatorname{GL}_n(\mathbb{Z})$. Then the Reidemeister classes of φ are the cosets of ker (I - M) in \mathbb{Z}^n , that is, $\mathcal{R}(\varphi) = \mathbb{Z}^n / \operatorname{ker} (I - M)$. Moreover, $R(\varphi) = |\det(I - M)|$ if this is non-zero, and $R(\varphi) = \infty$ otherwise.

In the sequel, we will also use R(M) to denote the Reidemeister number $R(\varphi)$ of the automorphism φ which is multiplication by M.

If the subgroup N in (1.1) is not characteristic, we can sometimes compute Reidemeister numbers using the *averaging formula*. We briefly explain the setting of this formula. We refer to [2] for a more general introduction and to [1], [20], [22] for the proofs of the results mentioned.

Consider the group of affine transformations $\operatorname{Aff}(\mathbb{R}^n) = \mathbb{R}^n \rtimes \operatorname{GL}_n(\mathbb{R})$ on \mathbb{R}^n with multiplication $(a, A) \cdot (b, B) = (a + Ab, AB)$. An *n*-dimensional Bieberbach group is a torsion-free cocompact discrete subgroup of $\mathbb{R}^n \rtimes C$, where C is a maximal compact subgroup of $\operatorname{GL}_n(\mathbb{R})$. Equivalently, an *n*-dimensional Bieberbach group Γ is a torsion-free subgroup of $\operatorname{Aff}(\mathbb{R}^n)$ such that its subgroup of pure translations $\Gamma \cap \mathbb{R}^n$ has finite index in Γ and is a uniform lattice of \mathbb{R}^n . A lattice of \mathbb{R}^n is a discrete and cocompact subgroup of \mathbb{R}^n and hence is isomorphic to \mathbb{Z}^n . From the geometric point of view, one mostly chooses C to be the orthogonal group O(n). Then Γ is a subgroup of the group of Euclidean motions and the quotient manifold $\Gamma \setminus \mathbb{R}^n$ inherits the flat metric structure from Euclidean space. From the algebraic point of view however, it is often easier to adopt another point of view. After an inner conjugation of $\operatorname{Aff}(\mathbb{R}^n)$, we may also assume that $\Gamma \cap \mathbb{R}^n$ is not only isomorphic to \mathbb{Z}^n , but really coincides with \mathbb{Z}^n . From now on, we will always assume that this is the case. The condition $\Gamma \cap \mathbb{R}^n = \mathbb{Z}^n$ implies that

- any element $(a, A) \in \Gamma$ has linear part A in $\operatorname{GL}_n(\mathbb{Z})$,
- any two elements $(a, A), (b, A) \in \Gamma$ are equal modulo \mathbb{Z}^n .

Hence $F := \{A \in \operatorname{GL}_n(\mathbb{Z}) \mid \exists a \in \mathbb{R}^n : (a, A) \in \Gamma\} \cong \Gamma/\mathbb{Z}^n$. Note that F is finite by definition. We call F the holonomy group of Γ .

Let φ be an automorphism of Γ . The second Bieberbach Theorem says that φ must be conjugation with some element $\operatorname{Aff}(\mathbb{R}^n)$, that is, there exists $(m, M) \in \operatorname{Aff}(\mathbb{R}^n)$ such that $\varphi(\gamma) = (m, M)\gamma(m, M)^{-1}$ for all $\gamma \in \Gamma$. In particular, $\varphi((z, I)) = (Mz, I)$ for all $z \in \mathbb{Z}^n$, hence φ restricts to the automorphism $M \in \operatorname{GL}_n(\mathbb{Z})$ on \mathbb{Z}^n . In [16] (see also [12]) we can find the following result:

LEMMA 2.3. Let Γ be a Bieberbach group with holonomy F. Let $\varphi \colon \Gamma \to \Gamma$ be an automorphism defined by $\varphi(\gamma) = (m, M)\gamma(m, M)^{-1}$. Then

$$R(\varphi) = \frac{1}{|F|} \sum_{A \in F} R(AM).$$

3. The nilpotent case

We start with the special case where E is itself nilpotent, corresponding to the subclass of nilmanifolds. In this case, already much is known, so we quickly review the Reidemeister spectra of the finitely generated torsion-free nilpotent groups of Hirsch length at most 4. We present the results in order of increasing nilpotency degree.

3.1. Nilpotency degree 1. The abelian groups we have to consider are the groups \mathbb{Z} , \mathbb{Z}^2 , \mathbb{Z}^3 and \mathbb{Z}^4 . It is easy to see that $\operatorname{Spec}_{\mathrm{R}}(\mathbb{Z}) = \{2, \infty\}$ and $\operatorname{Spec}_{\mathrm{R}}(\mathbb{Z}^n) = \mathbb{N} \cup \{\infty\}$ for $n \geq 2$. See for instance [19].

3.2. Nilpotency degree 2. The finitely generated 2-step nilpotent groups of Hirsch length at most 4 are of the form H_n and $H_n \times \mathbb{Z}$, where $H_n := \langle x, y, z | [z, x] = 1, [z, y] = 1, [y, x] = z^n \rangle$ for all $n \in \mathbb{N}$. Roman'kov showed [19, Section 3] that $\operatorname{Spec}_{\mathbb{R}}(H_n) = 2\mathbb{N} \cup \{\infty\}$ for n = 1, but his argument goes through for general n.

We next compute the Reidemeister spectrum of $H_n \times \mathbb{Z}$. Take a generator u of \mathbb{Z} . Let φ be an automorphism of $H_n \times \mathbb{Z}$, and let φ' and $\overline{\varphi}$ denote the induced automorphism on the center $\langle z, u \rangle$ and on the quotient $\langle \overline{x}, \overline{y} \rangle$, respectively. Further, denote by $M \in \operatorname{GL}_2(\mathbb{Z})$ the matrix representing $\overline{\varphi}$ relative to the basis $\{\overline{x}, \overline{y}\}$. It is easy to check that $\varphi(z)^n = [\varphi(y), \varphi(x)] = z^{n \det(M)}$. Hence

 $\varphi(z) = z^{\det(M)}$. So φ' is represented by the matrix

$$N := \begin{pmatrix} \det(M) & r \\ 0 & \varepsilon \end{pmatrix} \in \operatorname{GL}_2(\mathbb{Z})$$

for some $\varepsilon \in \{\pm 1\}$ and $r \in \mathbb{Z}$. If $\det(M) = 1$ or $\varepsilon = 1$, then $R(\varphi') = \infty$, and if $\det(M) = \varepsilon = -1$, then $R(\varphi') = 4$. Moreover, the addition formula (Lemma 2.1) simplifies to $R(\varphi) = R(\varphi')R(\overline{\varphi})$. Hence $R(\varphi) \in 4\mathbb{N} \cup \{\infty\}$.

Conversely, take $m \in \mathbb{N}$ and consider the automorphism

$$\varphi_m \colon H_n \times \mathbb{Z} \to H_n \times \mathbb{Z}, \qquad x^a y^b z^c u^d \mapsto y^a (xy^m)^b z^{-c} u^{-d}.$$

Note that φ_m induces the automorphisms N := -I and $M := \begin{pmatrix} 0 & 1 \\ 1 & m \end{pmatrix}$ on $\langle z, u \rangle$ and on $\langle \overline{x}, \overline{y} \rangle$, respectively. Hence $R(\varphi_m) = R(N)R(M) = 4m$. We conclude that $\operatorname{Spec}_{\mathbf{R}}(H_n \times \mathbb{Z}) = 4\mathbb{N} \cup \{\infty\}$ for all $n \in \mathbb{N}$.

3.3. Nilpotency degree 3. The 3-step nilpotent groups necessarily have Hirsch length 4. Gonçalves and Wong treated in [9, Example 5.2] an example of such a group and showed that this group has the R_{∞} property. But in fact, using an analogous argument, one can show that all finitely generated torsionfree nilpotent groups of class 3 and Hirsch length 4 have the R_{∞} property.

4. The 3-dimensional case

In this section, we determine the Reidemeister spectrum of all fundamental groups of solvmanifolds of dimension at most 3. In the sequel, we denote by $\mathbb{Z}^l \rtimes_A \mathbb{Z}, A \in \operatorname{GL}_l(\mathbb{Z})$, the semidirect product in which the generator of \mathbb{Z} is acting via A on \mathbb{Z}^l . More generally, we will use the notation $G \rtimes_{\psi} \mathbb{Z}$ for a semidirect product where the action is determined by an automorphism $\psi \in \operatorname{Aut}(G)$.

Recall that the groups E we are interested in fit in an exact sequence

$$1 \longrightarrow N \longrightarrow E \longrightarrow \mathbb{Z}^k \longrightarrow 1$$

with N finitely generated torsion-free nilpotent. Then $E \cong \mathbb{Z}$ if h(E) = 1 and either $E \cong \mathbb{Z}^2$ or $E \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}$ if h(E) = 2.

When h(E) = 3, either $E \cong \mathbb{Z}^3$ or $E \cong H_n$ if k = 0, and $E \cong \mathbb{Z}^2 \rtimes \mathbb{Z}$ if k = 1. Moreover, if h(E) = 3 and k = 2, the following lemma says that E can be viewed as a semidirect product $\mathbb{Z}^2 \rtimes \mathbb{Z}$ (so as an extension with $N \cong \mathbb{Z}^2$ and k = 1) as well:

LEMMA 4.1. Let E be an extension of \mathbb{Z} by \mathbb{Z}^k of Hirsch length k + 1. Then $E \cong N \rtimes \mathbb{Z}$ for some finitely generated torsion-free nilpotent group N of Hirsch length k.

PROOF. By assumption, the group E fits in the exact sequence

 $1 \longrightarrow \mathbb{Z} \longrightarrow E \stackrel{p}{\longrightarrow} \mathbb{Z}^k \longrightarrow 1.$

Take generators x_1, \ldots, x_k of \mathbb{Z}^k . The action of any element of \mathbb{Z}^k on \mathbb{Z} is either multiplication by 1 or -1, so we may assume that all x_2, \ldots, x_k act trivially on \mathbb{Z} , that is, \mathbb{Z} is contained in the center of $p^{-1}(\langle x_2, \ldots, x_k \rangle)$. Moreover, the quotient $p^{-1}(\langle x_2, \ldots, x_k \rangle)/\mathbb{Z} \cong \langle x_2, \ldots, x_k \rangle$ is abelian, hence $p^{-1}(\langle x_2, \ldots, x_k \rangle)$ is a finitely generated, torsion-free, 2-step nilpotent group. Since $E/p^{-1}(\langle x_2, \ldots, x_k \rangle) \cong \mathbb{Z}$, we conclude that $E \cong p^{-1}(\langle x_2, \ldots, x_k \rangle) \rtimes \mathbb{Z}$, as desired. \Box

Gonçalves and Wong already showed that $\mathbb{Z} \rtimes_{-1} \mathbb{Z}$ has the R_{∞} property [9, Theorem 2.2], so it remains to study the Reidemeister spectrum of $\mathbb{Z}^2 \rtimes \mathbb{Z}$.

We start by elaborating Lemma 2.1.

LEMMA 4.2. Suppose the subgroup \mathbb{Z}^n is characteristic in $\mathbb{Z}^n \rtimes_A \mathbb{Z}$. Then $\operatorname{Spec}_{\mathrm{R}}(\mathbb{Z}^n \rtimes_A \mathbb{Z}) = \{\infty\} \cup \{R(M) + R(AM) \mid M \in \operatorname{GL}_n(\mathbb{Z}), MA = A^{-1}M\}.$

PROOF. Take a generator t of the quotient \mathbb{Z} . Let φ be an automorphism of $\mathbb{Z}^n \rtimes_A \mathbb{Z}$ and suppose $\varphi|_{\mathbb{Z}^n}$ is multiplication with $M \in \operatorname{GL}_n(\mathbb{Z})$. Note that $R(\varphi)$ is infinite if φ induces the identity on the quotient \mathbb{Z} . If not, φ induces the automorphism $\overline{\varphi} = -\operatorname{Id}_{\mathbb{Z}}$ on \mathbb{Z} . Using the representatives 1 and t for $\mathcal{R}(\overline{\varphi})$, the addition formula implies that $R(\varphi) = R(M) + R(AM)$.

Moreover, one easily verifies that there exists an automorphism φ of $\mathbb{Z}^n \rtimes_A \mathbb{Z}$ inducing $M \in \operatorname{GL}_n(\mathbb{Z})$ on \mathbb{Z}^n and $-\operatorname{Id}_{\mathbb{Z}}$ on \mathbb{Z} if and only if $MA = A^{-1}M$. \Box

The following lemma asserts that for most A, the subgroup \mathbb{Z}^n is characteristic and Lemma 4.2 applies. It is an effortless generalization of [10, Lemma 2.1].

LEMMA 4.3. If A does not have 1 as eigenvalue, the subgroup \mathbb{Z}^n of $\mathbb{Z}^n \rtimes_A \mathbb{Z}$ is characteristic.

We distinguish cases based on the eigenvalues of A.

4.1. The matrix A has no eigenvalue 1 or -1. In [8], Gonçalves and Wong investigated the R_{∞} property of this group. They found that $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ does not have the R_{∞} property if and only if $A \in \mathrm{SL}_2(\mathbb{Z})$ and there exists M in $\mathrm{SL}_2(\mathbb{Z})$ satisfying $\mathrm{Tr}(M) = 0$ and $MA = A^{-1}M$. Writing

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$
 and $M = \begin{pmatrix} m & n \\ p & -m \end{pmatrix}$,

they showed that $MA = A^{-1}M$ is equivalent to (a - d)m + bp + cn = 0. Hence $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ does not have the R_{∞} property if and only if the system

(4.1)
$$\begin{cases} -m^2 - np = 1, \\ (a - d)m + bp + cn = 0, \end{cases}$$

has a solution (m, n, p) in \mathbb{Z}^3 . Furthermore, they showed that both R(M) and R(AM) equal $1 + \det(M) = 2$, hence:

PROPOSITION 4.4 ([8]). Let $A \in GL_2(\mathbb{Z})$ have eigenvalues different from ± 1 . Then $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ has Reidemeister spectrum $\{\infty\}$ or $\{4,\infty\}$. Moreover, if $\det(A) = -1$, then $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ has the R_∞ property.

Actually, if A has complex eigenvalues, system (4.1) has no solutions, so we moreover have:

PROPOSITION 4.5. Let $A \in \operatorname{GL}_2(\mathbb{Z})$ have non-real complex eigenvalues. Then $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ has the R_{∞} property.

PROOF. Note that the eigenvalues of A are the roots of the polynomial

$$\det(xI - A) = x^2 - \operatorname{Tr}(A)x + 1.$$

When A has complex eigenvalues, we therefore must have $Tr(A)^2 - 4 < 0$.

Suppose, for a contradiction, that $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ does not have the R_{∞} property, that is, the system (4.1) has an integral solution (m, n, p). Note that $c \neq 0$, for otherwise A would have eigenvalues ± 1 . Substituting the second equation in the top one gives

$$-m^2 + \frac{a-d}{c}mp + \frac{b}{c}p^2 = 1.$$

Note that $p \neq 0$. Denoting $y := m/p \in \mathbb{Q}$, this relation rewrites to

$$-y^{2} + \frac{a-d}{c}y + \frac{b}{c} - \frac{1}{p^{2}} = 0.$$

In particular, the discriminant $((a - d)/c)^2 + 4(b/c - 1/p^2)$ must be positive. Using $ad - bc = \det(A) = 1$, this simplifies to

$$0 \le \left(\frac{a-d}{c}\right)^2 + 4\left(\frac{b}{c} - \frac{1}{p^2}\right) = \frac{a^2 + d^2 - 2ad}{c^2} + \frac{4b}{c} - \frac{4}{p^2}$$
$$= \frac{(a^2 + d^2 - 2ad + 4bc)p^2 - 4c^2}{c^2p^2} = \frac{(\operatorname{Tr}(A)^2 - 4)p^2 - 4c^2}{c^2p^2}.$$

Since $Tr(A)^2 - 4 < 0$, we have reached a contradiction.

In general, determining which A allow an integral solution to (4.1) is hard and we will not pursue this further. Instead, we proceed with the remaining case where A has 1 or -1 as eigenvalue. Note that this implies that both eigenvalues are ± 1 .

4.2. The matrix A has eigenvalues 1 or -1. Let ε , $\delta \in \{\pm 1\}$ be the eigenvalues of A. Choose an eigenvector $v \in \mathbb{Q}^2$ corresponding to ε . Clearing denominators if necessary, we may assume that $v \in \mathbb{Z}^2$ and that v extends to a basis $\{v, w\}$ of \mathbb{Z}^2 . So, for some integer r, we can write

$$A = \begin{pmatrix} \varepsilon & r \\ 0 & \delta \end{pmatrix}.$$

Let $W_{\varepsilon} := \{z \in \mathbb{Z}^2 \mid A(z) = \varepsilon z\}$ denote the eigenspace of ε . We make the following easy observations, the proof of which we leave to the reader:

LEMMA 4.6. Let $A \in GL_n(\mathbb{Z})$.

- (a) If A does not have 1 as eigenvalue, the eigenspace W₋₁ is characteristic in Zⁿ ⋊_A Z.
- (b) The group $\mathbb{Z}^n \rtimes_A \mathbb{Z}$ has center $W_1 \times \langle t^d \rangle$, where d is the order of A if this order is finite. If A has infinite order, we set d := 0.

Swapping ε and δ if necessary, we have the following three cases.

Case 1. $\varepsilon = \delta = 1$. One easily sees that in this case, $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ is nilpotent. Specifically, $\mathbb{Z}^2 \rtimes_A \mathbb{Z} \cong \mathbb{Z}^3$ if A = I and $\mathbb{Z}^2 \rtimes_A \mathbb{Z} \cong H_n$ for some $n \in \mathbb{N}$ if $A \neq I$.

Case 2. $\varepsilon = \delta = -1$. We further distinguish the cases $r \neq 0$ and r = 0.

PROPOSITION 4.7. Let $A \neq -I$ in $\operatorname{GL}_2(\mathbb{Z})$ have repeated eigenvalue -1. Then $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ has the R_{∞} property.

PROOF. The eigenspace of -1 equals $W_{-1} = \langle v \rangle$. Moreover, the quotient $(\mathbb{Z}^2 \rtimes_A \mathbb{Z})/W_{-1} \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}$ has the R_{∞} property [9, Theorem 2.2]. Hence $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ has the R_{∞} property as well.

PROPOSITION 4.8. The group $\mathbb{Z}^n \rtimes_{-I} \mathbb{Z}$ has Reidemeister spectrum $2\mathbb{N} \cup \{\infty\}$ for all $n \geq 2$.

PROOF. Note that any $M \in \operatorname{GL}_n(\mathbb{Z})$ satisfies (-I)M(-I) = M. Consequently, Lemma 4.2 implies that

$$\operatorname{Spec}_{\mathbf{R}}(\mathbb{Z}^n \rtimes_{-I} \mathbb{Z}) = \{\infty\} \cup \{R(M) + R(-M) \mid M \in \operatorname{GL}_n(\mathbb{Z})\}.$$

Recall that R(M) + R(-M), if finite, equals $|\det(I - M)| + |\det(I + M)|$. Since $-1 \equiv 1 \mod 2$, surely $R(M) + R(-M) \in 2\mathbb{N} \cup \{\infty\}$.

Conversely, for m in $\mathbb{N} \cup \{0\}$, consider the $(n \times n)$ -matrix

$$M_m := \begin{pmatrix} 0 & \cdots & 0 & 1 \\ \hline & & & 0 \\ & & & \vdots \\ & I_{n-1} & & 0 \\ & & & m \end{pmatrix}.$$

One easily verifies that $|\det(I - M_m)| + |\det(I + M_m)|$ equals 2m if n is even and 2(m+1) if n is odd.

Case 3. $\varepsilon = 1, \delta = -1.$

PROPOSITION 4.9. Let A in $\operatorname{GL}_2(\mathbb{Z})$ have eigenvalues 1 and -1. Then $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ has the R_{∞} property.

PROOF. The center of $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ equals $Z = \langle v \rangle \times \langle t^2 \rangle$. The quotient

$$(\mathbb{Z}^2 \rtimes_A \mathbb{Z})/Z \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$$

satisfies the R_{∞} property [9, Proposition 2.3], hence so does $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$.

4.3 .	Conclusion.	We summarise	our findings	in the	following	table:
--------------	-------------	--------------	--------------	--------	-----------	--------

eigenvalues	Α	$\operatorname{Spec}_{\mathrm{R}}(\mathbb{Z}^2 \rtimes_A \mathbb{Z})$
-1, -1	$A = -I$ $A \neq -I$	$2\mathbb{N} \cup \{\infty\}$ $\{\infty\}$
1, -1		$\{\infty\}$
$\mathrm{real} \neq \pm 1$	det(A) = 1 $det(A) = -1$	$\{\infty\} \text{ or } \{4,\infty\}$ $\{\infty\}$
non-real		$\{\infty\}$

5. The 4-dimensional case

In this section, we consider extensions E as in (1.1) where h(E) = 4. Again, the special case k = 0 corresponds to the nilpotent groups of Section 3, and by Lemma 4.1, all other groups will appear in the situations k = 1 and k = 2.

In case k = 1, the sequence splits, hence $E \cong \mathbb{Z}^3 \rtimes \mathbb{Z}$ or $E \cong H_n \rtimes \mathbb{Z}$.

In case k = 2, the group E is an extension of \mathbb{Z}^2 by \mathbb{Z}^2 :

 $1 \longrightarrow \mathbb{Z}^2 \longrightarrow E \stackrel{p}{\longrightarrow} \mathbb{Z}^2 \longrightarrow 1.$

Let $\alpha \colon \mathbb{Z}^2 \to \operatorname{GL}_2(\mathbb{Z})$ denote the induced action of the quotient \mathbb{Z}^2 on \mathbb{Z}^2 . Take generators x, y of this quotient, write $\alpha(x) = A$ and $\alpha(y) = B$. As x and ycommute, the matrices A and B commute as well. This will severely limit the possible values of A and B. We need the following lemma:

LEMMA 5.1. Let $M \neq \pm I$ in $\operatorname{GL}_2(\mathbb{Z})$ have finite order. Then the centraliser of M in $\operatorname{GL}_2(\mathbb{Z})$ is the subgroup $\langle -I, M \rangle$.

PROOF. This lemma is easily checked by a case-by-case study, using the fact that there are, up to conjugacy, only six finite cyclic subgroups in $\text{GL}_2(\mathbb{Z})$ (see e.g. [17, p. 179]).

Using this, we will show that there are essentially three possibilities for A and B:

LEMMA 5.2. We can choose x and y such that one of the following conditions holds:

(a) B = I,

- (b) B = -I and A has infinite order,
- (c) B = -I and $A \neq \pm I$ has order 2.

PROOF. Note that $\alpha \colon \mathbb{Z}^2 \to \operatorname{GL}_2(\mathbb{Z})$ is not injective, as $\operatorname{PSL}_2(\mathbb{Z}) \cong \mathbb{Z}_2 * \mathbb{Z}_3$ does not contain \mathbb{Z}^2 so neither does $\operatorname{GL}_2(\mathbb{Z})$. Thus either $\mathbb{Z}^2/\operatorname{ker}(\alpha)$ is finite (Case 1) or $\mathbb{Z}^2/\operatorname{ker}(\alpha) \cong \mathbb{Z} \oplus \mathbb{Z}_p$ for some $p \in \mathbb{N}$ (Case 2).

Case 1. If $\mathbb{Z}^2/\ker(\alpha)$ is finite, both A and B have finite order. If A = I or B = I, we are in situation (a). Moreover, if both A = -I and B = -I, replacing y by xy results in situation (a) as well. Hence, it remains to prove the lemma when A or B is different from $\pm I$. We may assume $A \neq \pm I$. By Lemma 5.1, there exists $k \in \mathbb{Z}$ such that $B = \pm A^k$. Replacing y by $x^{-k}y$ if necessary, we can further reduce to $B = \pm I$. In case B = I, we are back in situation (a), so assume B = -I. If A happens to have order 2, we are in situation (c). Otherwise, A has order 3, 4 or 6. If A has order 3, the order of -A is 6. Replacing x by xy if necessary, we are left with the situation where A has order $d \in \{4, 6\}$. However, then $A^{d/2} = -I$ and replacing y by $x^{d/2}y$ delivers situation (a).

Case 2. Suppose $\mathbb{Z}^2/\ker(\alpha) \cong \mathbb{Z} \oplus \mathbb{Z}_p$. Take $u \in \mathbb{Z}^2$ such that \overline{u} generates \mathbb{Z}_p . If we choose x, y such that $y^k = u$ for some k in \mathbb{Z} , then B has finite order, whereas A must have infinite order as $\mathbb{Z}^2/\ker(\alpha)$ is infinite. By Lemma 5.1, this only happens when $B = \pm I$, corresponding to situations (a) and (b).

So, we may assume that x, y are generators as in Lemma 5.2 above. Choose a section $s: \mathbb{Z}^2 \to E$ and write u = s(x) and t = s(y). Then $E \cong (\mathbb{Z}^2 \rtimes_B \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}$ with action

$$\psi(zt^{l}) = uzt^{l}u^{-1} = uzu^{-1}(utu^{-1})^{l} = A(z)(n_{0}t)^{l}, \quad z \in \mathbb{Z}^{2}, \ l \in \mathbb{Z},$$

setting $n_0 := utu^{-1}t^{-1} \in \mathbb{Z}^2$. Thus either E is isomorphic to $\mathbb{Z}^3 \rtimes \mathbb{Z}$, or E is isomorphic to $(\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}$ where $\psi|_{\mathbb{Z}^2} = A$ has infinite order or $A \neq \pm I$ and $A^2 = I$.

Hence, it remains to study the Reidemeister spectrum of the groups $\mathbb{Z}^3 \rtimes \mathbb{Z}$, $H_n \rtimes \mathbb{Z}$ and $(\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}$.

5.1. The semidirect product $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$. In this subsection, we determine the Reidemeister spectrum of the group $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$. We distinguish cases based on the eigenvalues of A.

5.1.1. The matrix A does not have eigenvalue 1. According to Lemma 4.2, we must determine which M satisfy $MA = A^{-1}M$. If any such M exists, A is conjugate to its inverse. The following lemma says that this only happens if A has eigenvalue -1, as we assume that 1 is not an eigenvalue of A.

LEMMA 5.3. Let $n \in \mathbb{N}$ be odd, and $A \in GL_n(\mathbb{Z})$. Suppose A is conjugate to A^{-1} . Then A has eigenvalue det(A).

PROOF. Set $\delta = \det(A)$ and write $\lambda_1, \ldots, \lambda_n$ the eigenvalues of A. Let p_A and $p_{A^{-1}}$ denote the characteristic polynomial of A and A^{-1} , respectively.

Since A and A^{-1} are conjugate, their characteristic polynomials coincide, hence

$$p_{A}(x) = p_{A^{-1}}(x) = \prod_{i=1}^{n} (x - \lambda_{i}^{-1}) = \frac{x^{n}}{\lambda_{1} \dots \lambda_{n}} \prod_{i=1}^{n} (\lambda_{i} - x^{-1})$$
$$= (-1)^{n} \delta x^{n} \prod_{i=1}^{n} (x^{-1} - \lambda_{i}) = -\delta x^{n} p_{A}(x^{-1}).$$

Writing $p_A(x) = \sum_{i=0}^n a_i x^i$, this implies that $a_i = -\delta a_{n-i}$, for $x^n p_A(x^{-1})$ equals the reciprocal polynomial $\sum_{i=0}^n a_{n-i} x^i$ of p_A . It easily follows that

$$p_{A}(\delta) = \sum_{i=0}^{n} a_{i}\delta^{i} = \sum_{i=0}^{(n-1)/2} (a_{i}\delta^{i} + a_{n-i}\delta^{n-i}) = \sum_{i=0}^{(n-1)/2} \delta^{i}(a_{i} + \delta a_{n-i}) = 0,$$

that is, A has eigenvalue δ .

Subsequently, if neither 1 nor -1 are eigenvalues of A, Lemma 4.2 immediately implies the following:

PROPOSITION 5.4. Let $A \in GL_3(\mathbb{Z})$ have eigenvalues different from 1 and -1. Then $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has the R_{∞} property.

So it remains to study A that do have eigenvalue -1. Recall from Proposition 4.8 that $\operatorname{Spec}_{\mathbb{R}}(\mathbb{Z}^3 \rtimes_A \mathbb{Z}) = 2\mathbb{N} \cup \{\infty\}$ if A = -I. If $A \neq -I$, we have:

PROPOSITION 5.5. Let $A \in GL_3(\mathbb{Z})$ be different from -I and suppose that A has eigenvalue -1, but not eigenvalue 1. Then $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has the R_{∞} property.

PROOF. Let $\{u,v,w\}$ be a basis of \mathbb{Z}^3 transforming A into a matrix of the form

$$\begin{pmatrix} -1 & C \\ 0 & A' \end{pmatrix}$$

for some $A' \in GL_2(\mathbb{Z}), C \in \mathbb{Z}^{1 \times 2}$. Suppose first that -1 is not an eigenvalue of A'. Let φ be an automorphism inducing

$$M := \begin{pmatrix} m & N \\ P & Q \end{pmatrix} \in \mathrm{GL}_3(\mathbb{Z})$$

on \mathbb{Z}^3 for some $m \in \mathbb{Z}$, $N \in \mathbb{Z}^{1 \times 2}$, $P \in \mathbb{Z}^{2 \times 1}$ and $Q \in \mathbb{Z}^{2 \times 2}$. Since W_{-1} is characteristic, we must have P = 0, implying $m \in \{\pm 1\}$. As a result, either M or AM has eigenvalue 1, or equivalently, either R(M) or R(AM) is infinite. Hence $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has the R_{∞} property.

If A' does have eigenvalue -1, then -1 is the only eigenvalue of A'. If $W_{-1} =$ $\langle u \rangle$, it is easy to check that $A' \neq -I$. The quotient $\mathbb{Z}^3 \rtimes_A \mathbb{Z}/W_{-1} \cong \mathbb{Z}^2 \rtimes_{A'} \mathbb{Z}$ has the R_{∞} property by Lemma 4.7, hence so has $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$. If $W_{-1} \neq \langle u \rangle$, then $\mathbb{Z}^3 \rtimes_A \mathbb{Z}/W_{-1} \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}$ has the R_∞ property as well.

5.1.2. The matrix A has eigenvalue 1. We present our results in decreasing order of the algebraic multiplicity of the eigenvalue 1.

If 1 is the only eigenvalue of A, the group $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ is nilpotent. If the eigenvalue 1 has algebraic multiplicity two, we have the following:

PROPOSITION 5.6. Suppose 1 is an eigenvalue of $A \in GL_3(\mathbb{Z})$ of algebraic multiplicity two. Then $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has the R_{∞} property.

PROOF. If A has eigenvalue 1 of algebraic multiplicity two, the remaining eigenvalue must be -1. Accordingly, let $\{u, v, w\}$ be a basis of \mathbb{Z}^3 transforming A into a matrix of the form

$$\begin{pmatrix} 1 & r & s \\ 0 & 1 & n \\ 0 & 0 & -1 \end{pmatrix}$$

for some $r, s, n \in \mathbb{Z}$. If r is zero, A has order two, so the center of $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ is generated by u, v and t^2 . Moreover, the quotient $\mathbb{Z}^3 \rtimes_A \mathbb{Z}/\langle u, v, t^2 \rangle \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$ has the R_{∞} property, hence so has $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$.

Similarly, if r is nonzero, the order of A is infinite. Thus $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has center $\langle u \rangle$ and $\mathbb{Z}^3 \rtimes_A \mathbb{Z}/\langle u \rangle$ has the R_∞ property by Lemma 4.9. We conclude that $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has the R_∞ property as well, as desired.

If the eigenvalue 1 is not repeated, let $\{x, y, z\}$ be a basis of \mathbb{Z}^3 such that A takes the form

$$(5.1) \qquad \qquad \begin{pmatrix} 1 & r & s \\ 0 & \\ 0 & A' \end{pmatrix}$$

for some $r, s \in \mathbb{Z}$ and $A' \in GL_2(\mathbb{Z})$ not having eigenvalue 1. Thus A' has either repeated eigenvalue -1 or real or complex eigenvalues. This leaves the following possibilities:

- (i) A' = -I, (ii) $A' = \begin{pmatrix} -1 & n \\ 0 & -1 \end{pmatrix}$ for some $n \neq 0$,
- (iii) A' has real eigenvalues not equal to ± 1 ,
- (iv) A' has non-real complex eigenvalues.

In cases (ii) and (iii), the matrix A has infinite order, hence x generates the center of $E = \mathbb{Z}^3 \rtimes_A \mathbb{Z}$. Moreover, in $E/\langle x \rangle \cong \mathbb{Z}^2 \rtimes_{A'} \mathbb{Z}$, the subgroup $\mathbb{Z}^2 = \langle \overline{y}, \overline{z} \rangle$ is characteristic by Lemma 4.3, hence \mathbb{Z}^3 is characteristic in E. Consequently, we can apply Lemma 4.2 to compute the Reidemeister spectrum of E. To this end.

take $M \in GL_3(\mathbb{Z})$ satisfying $MA = A^{-1}M$. Since $\langle x \rangle = Z(E)$ is characteristic, we can write

$$M = \begin{pmatrix} m & N \\ 0 & Q \end{pmatrix}$$

for some $m \in \{\pm 1\}$, $N \in \mathbb{Z}^{1 \times 2}$ and $Q \in \mathbb{Z}^{2 \times 2}$. If m = 1, clearly $R(M) = \infty$. So assume m = -1. Imposing AMA = M explicitly gives

$$\begin{pmatrix} -1 & -C + NA' + CQA' \\ 0 & A'QA' \end{pmatrix} = \begin{pmatrix} -1 & N \\ 0 & Q \end{pmatrix},$$

hence A'QA' = Q. Moreover, it is easy to see that

$$R(M) + R(AM) = 2(R(Q) + R(A'Q)).$$

Thus $\operatorname{Spec}_{R}(\mathbb{Z}^{3} \rtimes_{A} \mathbb{Z}) \subseteq 2\operatorname{Spec}_{R}(\mathbb{Z}^{2} \rtimes_{A'} \mathbb{Z})$. Therefore, Lemmas 4.7 and 4.4 immediately imply the following results.

PROPOSITION 5.7. Suppose A is conjugate to a matrix of the form (5.1) where $A' = \begin{pmatrix} -1 & n \\ 0 & -1 \end{pmatrix}$ for some $n \neq 0$. Then $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has the R_{∞} property.

PROPOSITION 5.8. Suppose A is conjugate to a matrix of the form (5.1), where A' has real eigenvalues $\neq \pm 1$. Then $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has Reidemeister spectrum $\{\infty\}$ or $\{8,\infty\}$. Moreover, if $\mathbb{Z}^2 \rtimes_{A'} \mathbb{Z}$ has the R_{∞} property, so has $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$.

REMARK 5.9. In general, the converse statement to Proposition 5.8 is not true: there exists A such that $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has the R_{∞} property, but $\mathbb{Z}^2 \rtimes_{A'} \mathbb{Z}$ does not. Indeed, consider

$$A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 5 & 2 \\ 0 & 2 & 1 \end{pmatrix}, \text{ so that } A' := \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$$

In the above argument, $\operatorname{Spec}_{R}(\mathbb{Z}^{3} \rtimes_{A} \mathbb{Z}) = \{8, \infty\}$ precisely when there exists Q in $\operatorname{SL}_{2}(\mathbb{Z})$ satisfying A'QA' = Q and

(5.2)
$$-N = C(I - QA')(I - A')^{-1} \in \mathbb{Z}^{1 \times 2},$$

where C = (0, 1). A straightforward calculation shows that if $Q \in SL_2(\mathbb{Z})$ satisfies A'QA' = Q, all entries of I - QA' are odd. At the same time, it is easy to check that

$$(I - A')^{-1} = \frac{1}{2} \begin{pmatrix} 0 & -1 \\ -1 & 2 \end{pmatrix}$$

Subsequently, condition (5.2) never holds. At the same time, $Q' = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ satisfies AQ'A = Q' and R(Q') + R(AQ') = 4. By Lemma 4.2, we conclude that the group $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has the R_{∞} property, while $\mathbb{Z}^2 \rtimes_{A'} \mathbb{Z}$ does not. In cases (i) and (iv), the matrix A has finite order, say order d. In this situation, we can use the averaging formula, Lemma 2.3, to compute Reidemeister numbers on $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$. Indeed, setting $\widetilde{A} := \begin{pmatrix} 1 & 0 \\ 0 & A \end{pmatrix} \in \mathrm{GL}_4(\mathbb{Z})$, the embedding

$$i: \mathbb{Z}^3 \rtimes_A \mathbb{Z} \to \operatorname{Aff}(\mathbb{R}^4), \qquad x^a y^b z^c t^n \mapsto \left(\begin{pmatrix} n/d \\ a \\ b \\ c \end{pmatrix}, \widetilde{A}^n \right)$$

identifies $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ with a 4-dimensional Bieberbach group with holonomy $\{I, \widetilde{A}, \ldots, \widetilde{A}^{d-1}\}$. Let φ be an automorphism of $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$. Let N denote the matrix representing φ on the center $\langle t^d, x \rangle$ and let M denote the matrix representing the induced automorphism $\overline{\varphi}$ on $\mathbb{Z}^2 = \langle \overline{y}, \overline{z} \rangle$. The restriction of φ to $\langle t^d, x, y, z \rangle \cong \mathbb{Z}^4$ is thus represented by a matrix of the form

$$\widetilde{M} := \begin{pmatrix} N & * \\ 0 & M \end{pmatrix} \in \mathrm{GL}_4(\mathbb{Z}).$$

Then, for all integers i, the matrix $\widetilde{A}^i \widetilde{M}$ has the form

$$\widetilde{A}^{i}\widetilde{M} = \begin{pmatrix} I & * \\ 0 & A'^{i} \end{pmatrix} \begin{pmatrix} N & * \\ 0 & M \end{pmatrix} = \begin{pmatrix} N & * \\ 0 & A'^{i}M \end{pmatrix},$$

showing $R(\widetilde{A}^i \widetilde{M}) = R(N)R(A'^i M)$. The averaging formula hence says that

$$R(\varphi) = \frac{1}{d} R(N) \sum_{i=0}^{d-1} R(A^{\prime i}M).$$

We start with case (i), so that A has order 2. Tahara showed [21, Proposition 2] that in this case, A is conjugate over $GL_3(\mathbb{Z})$ to the matrix

(5.3)
$$\begin{pmatrix} 1 & 0 & \delta \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

where δ is either 0 or 1.

PROPOSITION 5.10. Suppose A is of the form (5.3) where $\delta \in \{0, 1\}$.

- (a) If $\delta = 0$, the group $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has Reidemeister spectrum $2\mathbb{N} \cup \{\infty\}$.
- (b) If $\delta = 1$, the group $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has Reidemeister spectrum $4\mathbb{N} \cup \{\infty\}$.

PROOF. Write $E = \mathbb{Z}^3 \rtimes_A \mathbb{Z}$. Take an automorphism φ of E. The center of E is generated by x and t^2 , so we can write

$$\varphi(x)=x^at^{2b},\quad \varphi(y)=x^cy^dz^et^f,\quad \varphi(z)=x^gy^hz^it^j,\quad \varphi(t)=x^ky^lz^mt^n$$

for some a, \ldots, n in \mathbb{Z} . Moreover, f and j are even and n is odd as $\mathbb{Z}^3 \times \langle t^2 \rangle$ is characteristic in E.

The map φ is a morphism if and only if it respects the relations

[x, y] = 1, [x, z] = 1, [y, z] = 1, [x, t] = 1, $[y, t] = y^{-2}$, $[z, t] = x^{\delta} z^{-2}$. As both f and j are even, φ already respects the first four relations, so we only

need to examine the last two relations. Equating $\varphi(y)^{-2} = [\varphi(y), \varphi(t)]$ gives

$$\begin{aligned} x^{-2c}y^{-2d}z^{-2e}t^{-2f} &= t^{-f}z^{-e}y^{-d}x^{-c}t^{-n}z^{-m}y^{-l}x^{-k}x^{c}y^{d}z^{e}t^{f}x^{k}y^{l}z^{m}t^{n} \\ &= z^{-e}y^{-d}y^{-d}x^{\delta e}z^{-e}t^{-n}t^{n} = x^{\delta e}y^{-2d}z^{-2e}, \end{aligned}$$

so $\delta e = -2c$ and f = 0. Similarly, equating $\varphi(x)^{\delta}\varphi(z)^{-2} = [\varphi(z), \varphi(t)]$ gives $x^{\delta a - 2g}y^{-2h}z^{-2i}t^{2\delta b - 2j} = t^{-j}z^{-i}y^{-h}x^{-g}t^{-n}z^{-m}y^{-l}x^{-k}x^{g}y^{h}z^{i}t^{j}x^{k}y^{l}z^{m}t^{n}$ $= z^{-i}y^{-h}y^{-h}x^{\delta i}z^{-i}t^{-n}t^{n} = x^{\delta i}y^{-2h}z^{-2i}.$

hence $\delta i = \delta a - 2g$ and $\delta b = j$.

The averaging formula says that $R(\varphi) = R(N)(R(M) + R(-M))/2$ where

$$N := \begin{pmatrix} n & b \\ 2k + \delta m & a \end{pmatrix} \quad \text{and} \quad M := \begin{pmatrix} d & h \\ e & i \end{pmatrix}.$$

We distinguish two cases.

Case 1. Suppose first that $\delta = 1$. Then e = -2c is even, $a \equiv i \mod 2$ and b = j is even. As $R(M) = \pm 1$ is odd and e is even, both d and i (and thus also a) are odd. So b and e are even and n, a, d and i are odd. It follows that R(N) is either infinite or even. In addition, if R(M) + R(-M) is finite, then

$$RM + R(-M) = |1 + \det(M) - \operatorname{Tr}(M)| + |1 + \det(M) + \operatorname{Tr}(M)| \in 4\mathbb{N},$$

as $\operatorname{Tr}(M) = d + i$ is even. Hence $\operatorname{Spec}_{\mathbf{R}}(E) \subseteq 4\mathbb{N} \cup \{\infty\}$.

Conversely, for $\alpha \in \mathbb{N}$, setting

$$\begin{aligned} {}_{\alpha}(x) &= x^{1-2\alpha}t^{4\alpha}, \qquad \varphi_{\alpha}(y) &= x^{-1}y^{-1}z^{2}, \\ \varphi_{\alpha}(z) &= x^{1-\alpha}yz^{-1}t^{2\alpha}, \qquad \varphi_{\alpha}(t) &= zt^{-1} \end{aligned}$$

defines an automorphism φ_{α} of E inducing the automorphisms

$$\begin{pmatrix} -1 & 2\alpha \\ 1 & 1-2\alpha \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} -1 & 1 \\ 2 & -1 \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Z})$$

on $Z(E) = \langle t^2, x \rangle$ and $\mathbb{Z}^2 = \langle \overline{y}, \overline{z} \rangle$, respectively. It is easy to check that $R(\varphi_{\alpha}) = 4\alpha$, hence in this case, $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has Reidemeister spectrum $4\mathbb{N} \cup \{\infty\}$.

Case 2. If $\delta = 0$, then R(N) is either infinite or even as n is odd. Moreover, R(M) + R(-M), if finite, is always even, hence $R(\varphi) \in 2\mathbb{N} \cup \{\infty\}$.

Conversely, for $\alpha \in \mathbb{N}$, setting

$$\varphi_{\alpha}(x) = x^{1-2\alpha}t^2, \quad \varphi_{\alpha}(y) = z, \quad \varphi_{\alpha}(z) = yz, \quad \varphi_{\alpha}(t) = x^{\alpha}t^{-1}$$

defines an automorphism φ_{α} of E inducing

$$\begin{pmatrix} -1 & 1\\ 2\alpha & 1-2\alpha \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & 1\\ 1 & 1 \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Z})$$

on $Z(E) = \langle t^2, x \rangle$ and $\mathbb{Z}^2 = \langle \overline{y}, \overline{z} \rangle$, respectively. It is easy to check that $R(\varphi_{\alpha}) = 2\alpha$, which concludes the proof.

If A' has complex eigenvalues, the order d of A' is either 3, 4 or 6. Additionally, if $d \in \{4, 6\}$, then $A'^{d/2} = -I$. This will imply the following:

PROPOSITION 5.11. Suppose A is of the form (5.1), where A' has order 4 or 6. Then $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has the R_{∞} property.

PROOF. Write $E = \mathbb{Z}^3 \rtimes_A \mathbb{Z}$. Let φ be an automorphism and denote by M the matrix representing the induced automorphism $\overline{\varphi}$ on $E/\langle x,t\rangle = \langle \overline{y},\overline{z}\rangle$. By the averaging formula, the result will follow once we show that $R(A^kM) = \infty$ for some $k \in \mathbb{Z}$. Thereto, write $\varphi(t) = z_0 t^{\varepsilon}$ for some ε in $\{\pm 1\}$ and z_0 in \mathbb{Z}^3 . A simple calculation shows that $MA' = A'^{\varepsilon}M$. From Section 4.1, it immediately follows that $R(M) = \infty$ if $\varepsilon = -1$. So assume $\varepsilon = 1$. Lemma 5.1 shows that $M = \pm A'^k$ for some $k \in \mathbb{Z}$. Using $A'^{d/2} = -I$, we may assume that $M = A'^k$, showing $R(A'^{-k}M) = R(I) = \infty$ indeed.

So it remains to examine the situation where A' has order 3. Tahara [21, Proposition 3] showed that in this situation, A is conjugate over $GL_3(\mathbb{Z})$ to the matrix

(5.4)
$$\begin{pmatrix} 1 & 0 & \delta \\ 0 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix},$$

where δ is either 0 or 1.

PROPOSITION 5.12. Suppose A is of the form (5.4), where $\delta = 0$ or 1. Then $\mathbb{Z}^3 \rtimes_A \mathbb{Z}$ has Reidemeister spectrum $6\mathbb{N} \cup \{\infty\}$.

PROOF. Write $E = \mathbb{Z}^3 \rtimes_A \mathbb{Z}$. We continue to write

$$A' := \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}.$$

We first show that $\operatorname{Spec}_{\mathbb{R}}(E) \subseteq 6\mathbb{N} \cup \{\infty\}$. To this end, take an automorphism φ of E. Write $\overline{\varphi}$ the induced automorphism on $E/\langle x, t^3 \rangle$. If $\overline{\varphi}$ does not induce the identity on \mathbb{Z}_3 , we know from Section 4.1 that $R(\overline{\varphi}) = \infty$. Moreover, $\mathbb{Z}^3 \times \langle t^3 \rangle$ is characteristic in E. Therefore, if $R(\varphi)$ is finite, we can write

$$\varphi(x)=x^at^{3b},\quad \varphi(y)=x^cy^dz^et^{3f},\quad p(z)=x^gy^hz^it^{3j},\quad \varphi(t)=x^ky^lz^mt^n$$

for some a, \ldots, n in \mathbb{Z} , where $n \equiv 1 \mod 3$. Again, φ is a morphism if and only if it respects the relations

$$[y,t] = x^{\delta}y^{-2}z^{-1}, \qquad [z,t] = z^{-1}y.$$

Equating $\varphi(y)\varphi(z)^{-1} = [\varphi(z),\varphi(t)]$ gives

$$\begin{aligned} x^{c-g}y^{d-h}z^{e-i}t^{3f-3j} &= t^{-3j}z^{-i}y^{-h}x^{-g}t^{-n}z^{-m}y^{-l}x^{-k}x^{g}y^{h}z^{i}t^{3j}x^{k}y^{l}z^{m}t^{n} \\ &= z^{-i}y^{-h}x^{\delta h}y^{-h}z^{-h}y^{i}t^{-n}t^{n} = x^{\delta h}y^{-2h+i}z^{-h-i}, \end{aligned}$$

hence $c = \delta h + g$, d = -h + i, e = -h and f = j. Similarly, equating $\varphi(x)^{\delta}\varphi(y)^{-2}\varphi(z)^{-1} = [\varphi(y),\varphi(t)]$ gives

$$\begin{aligned} x^{\delta a - 2c - g} y^{-2d - h} z^{-2e - i} t^{3b\delta - 6f - 3j} \\ &= t^{-3f} z^{-e} y^{-d} x^{-c} t^{-n} z^{-m} y^{-l} x^{-k} x^{c} y^{d} z^{e} t^{3f} x^{k} y^{l} z^{m} t^{n} \\ &= z^{-e} y^{-d} x^{\delta d} y^{-d} z^{-d} y^{e} t^{-n} t^{n} = x^{\delta d} y^{-2d + e} z^{-d - e}, \end{aligned}$$

so we moreover have $\delta a = \delta d + 2c + g$ and $\delta b = 3f$. The averaging formula says that $R(\varphi) = (R(N)/3) \sum_{i=0}^{2} R(A'^{i}M)$, with

$$N := \begin{pmatrix} n & b \\ 3k + \delta(m+l) & a \end{pmatrix} \quad \text{and} \quad M := \begin{pmatrix} e+i & -e \\ e & i \end{pmatrix}.$$

Since M and A' commute, $M = \pm A'^s$ for some $s \in \mathbb{Z}$, hence $\sum_{i=0}^2 R(A'^i M)$ is either 6 or ∞ . Since $n \equiv 1 \mod 3$ and $\delta b = 3f$, we moreover have $R(N) \in 3\mathbb{N} \cup \{\infty\}$ as $\delta \in \{0, 1\}$. Thus $R(\varphi) \in 6\mathbb{N} \cup \{\infty\}$.

Conversely, for $\alpha \in \mathbb{N}$, setting

$$\begin{aligned} \varphi_{\alpha}(x) &= x^{3\alpha - 1} t^{3(\delta 3\alpha + (1 - \delta))}, \qquad \varphi_{\alpha}(y) &= x^{\delta \alpha} y^{-1} t^{3\delta \alpha}, \\ \varphi_{\alpha}(z) &= x^{\delta \alpha} z^{-1} t^{3\delta \alpha}, \qquad \qquad \varphi_{\alpha}(t) &= x^{(1 - \delta)\alpha} y t \end{aligned}$$

defines an automorphism φ_{α} of E inducing the automorphisms

$$N = \begin{pmatrix} 1 & \delta 3\alpha + (1-\delta) \\ \delta + (1-\delta)3\alpha & 3\alpha - 1 \end{pmatrix} \text{ and } M = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \in \operatorname{GL}_2(\mathbb{Z})$$

on $\langle t^3, x \rangle$ and $\langle \overline{y}, \overline{z} \rangle$, respectively. By construction

$$R(N) = 3\alpha$$
 and $\sum_{i=0}^{2} R(A'^{i}M) = 6$,

hence $R(\varphi) = 6\alpha$. This completes the proof.

A		$\operatorname{Spec}_{\mathrm{R}}(\mathbb{Z}^3 \rtimes_A \mathbb{Z})$
$1\notin\Lambda(A)$	$A = -I$ $A \neq -I$	$2\mathbb{N} \cup \{\infty\}$ $\{\infty\}$
$\begin{pmatrix} 1 & * & * \\ 0 & 1 & * \\ 0 & 0 & -1 \end{pmatrix}$, .	{∞} {∞}
$\begin{pmatrix} 1 & * & * \\ 0 & -1 & n \\ 0 & 0 & -1 \end{pmatrix}$	n eq 0	$\{\infty\}$
$\begin{pmatrix} 1 & 0 & \delta \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$	$\begin{split} \delta &= 0\\ \delta &= 1 \end{split}$	$2\mathbb{N} \cup \{\infty\}\\ 4\mathbb{N} \cup \{\infty\}$
$\begin{pmatrix} 1 & * & * \\ 0 & & \\ 0 & & A' \end{pmatrix}$	$\Lambda(A') \subseteq \mathbb{R} \setminus \{\pm 1\}:$ $\det(A') = 1$ $\det(A') = -1$	$\{\infty\} \text{ or } \{8,\infty\}$ $\{\infty\}$
$\begin{pmatrix} 1 & * & * \\ 0 & & \\ 0 & & A' \end{pmatrix}$	$A^{\prime 4} = I$ or $A^{\prime 6} = I$ $A^{\prime 3} = I$	$\{\infty\}$ $6\mathbb{N}\cup\{\infty\}$

5.1.3. Conclusion. We summarise the results of this subsection in the following table, where we use the notation $\Lambda(M) := \{\text{eigenvalues of } M\}.$

5.2. The semidirect product $(\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}$. In this subsection, we determine the Reidemeister spectrum of the group $(\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}$. Denoting u a generator of the outer \mathbb{Z} and t a generator of the inner \mathbb{Z} , the action ψ of u on $\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}$ is assumed to be of the form $\psi(zt^k) = A(z)(n_0t)^k$, where $n_0 \in \mathbb{Z}^2$ and A has either infinite order or $A \neq \pm I$ has order two.

We make the following observation.

LEMMA 5.13. The subgroup \mathbb{Z}^2 is characteristic in $(\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}$.

PROOF. Write $E = (\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}$. Let φ be an automorphism of E and take z in \mathbb{Z}^2 . The relation $z^2 = ztz^{-1}t^{-1}$ shows that z^2 belongs to [E, E], which is characteristic. Hence $\varphi(z)^2 \in [E, E]$. Moreover, $[E, E] \subseteq \mathbb{Z}^2$ as $E/\mathbb{Z}^2 \cong \mathbb{Z}^2$ is abelian. Hence $\varphi(z)^2 \in \mathbb{Z}^2$, showing $\varphi(z) \in \mathbb{Z}^2$ as well.

5.2.1. The matrix A has order 2. We start by applying Lemma 2.1 to the situation $A \neq \pm I$ and $A^2 = I$.

PROPOSITION 5.14. Let $E = (\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}$, where $A = \psi|_{\mathbb{Z}^2} \neq \pm I$ and $A^2 = I$. Then E has the R_{∞} property.

PROOF. As $A \neq \pm I$ and $A^2 = I$, there exists a basis $\{v, w\}$ of \mathbb{Z}^2 transforming A into the matrix

$$A = \begin{pmatrix} 1 & r \\ 0 & -1 \end{pmatrix}.$$

Let φ be an automorphism of E, and let $\overline{\varphi}$ denote the induced automorphism on the quotient $E/\mathbb{Z}^2 \cong \mathbb{Z}^2$. Write $\overline{\varphi}(\overline{t}) = \overline{t}^k \overline{u}^m$, $\overline{\varphi}(\overline{u}) = \overline{t}^l \overline{u}^n$ and $\varphi(v) = v^a w^c$, $\varphi(w) = v^b w^d$ so that $\overline{\varphi}$ and $\varphi|_{\mathbb{Z}^2}$ are represented by the matrices

$$K = \begin{pmatrix} k & l \\ m & n \end{pmatrix}$$
 and $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{GL}_2(\mathbb{Z})$

respectively. The action of the quotient E/\mathbb{Z}^2 on \mathbb{Z}^2 is given by $\alpha(\overline{t}^e \overline{u}^f) = (-I)^e A^f$ for all $e, f \in \mathbb{Z}$. It is easy to check that $M\alpha(\overline{z}) = \alpha(\overline{\varphi}(\overline{z}))M$ for all \overline{z} in E/\mathbb{Z}^2 . Applying the condition on \overline{t} gives $-M = (-I)^k A^m M$. Hence k is odd and m must be even, which forces n to be odd as $K \in \mathrm{GL}_2(\mathbb{Z})$. Applying the condition on \overline{u} gives $MA = (-I)^l A^n M$, so $MA = (-I)^l AM$ since n is odd. We distinguish two cases, based on the parity of l.

Case 1. Even *l*. In this case, *A* and *M* commute. Hence $M \in \{\pm I, \pm A\}$ by Lemma 5.1. At the same time, the Reidemeister classes $[1]_{\overline{\varphi}}, [\overline{t}]_{\overline{\varphi}}, [\overline{u}]_{\overline{\varphi}}, [\overline{t}\overline{u}]_{\overline{\varphi}}$ are all different, as

$$I - K = \begin{pmatrix} 1 - k & -l \\ -m & 1 - n \end{pmatrix} \in \begin{pmatrix} 2\mathbb{Z} & 2\mathbb{Z} \\ 2\mathbb{Z} & 2\mathbb{Z} \end{pmatrix}$$

The addition formula implies that $R(\varphi)$ is at least R(M) + R(-M) + R(AM) + R(-AM). As $M \in \{\pm I, \pm A\}$, one of these terms equals $R(I) = \infty$, showing $R(\varphi) = \infty$.

Case 2. Odd l. Now MA = -AM. Equating these matrices explicitly gives

$$\begin{pmatrix} a & -b + ar \\ c & -d + cr \end{pmatrix} = \begin{pmatrix} -a - cr & -b - dr \\ c & d \end{pmatrix}.$$

Note that r = 0 implies a = d = 0 and $r \neq 0$ implies a + d = 0. So in both cases M has trace zero. The same holds for AM, for this matrix also satisfies (AM)A = -A(AM). In addition, the Reidemeister classes $[1]_{\overline{\varphi}}$ and $[\overline{u}]_{\overline{\varphi}}$ are different, as

$$I - K = \begin{pmatrix} 1 - k & -l \\ -m & 1 - n \end{pmatrix} \in \begin{pmatrix} \mathbb{Z} & \mathbb{Z} \\ 2\mathbb{Z} & 2\mathbb{Z} \end{pmatrix}.$$

Hence $R(\varphi)$ is at least R(M) + R(AM). As M and AM both have trace zero, and $\det(A) = -1$, either $|\det(I - M)| = 1 + \det(M)$ or $|\det(I - AM)| = 1 + \det(AM)$ vanishes, thus, either R(M) or R(AM) is infinite. We conclude that $R(\varphi) = \infty$.

5.2.2. The matrix A has infinite order. Similarly, we can determine the Reidemeister spectrum if A has infinite order. We begin by elaborating Lemma 2.1.

LEMMA 5.15. Let $E = (\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}$, where $A = \psi_{|\mathbb{Z}^2}$ has infinite order. Let φ be an automorphism of E, write $\varphi_{|\mathbb{Z}^2} = M \in \operatorname{GL}_2(\mathbb{Z})$. Then $R(\varphi)$, if finite, equals

$$R(\varphi) = R(M) + R(-M) + R(AM) + R(-AM).$$

In addition, either $MA = A^{-1}M$ or $MA = -A^{-1}M$ and the group $\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}$ is characteristic in E.

PROOF. Let $\overline{\varphi}$ denote the induced automorphism on the quotient $E/\mathbb{Z}^2 \cong \mathbb{Z}^2$. Writing $\overline{\varphi}(\overline{t}) = \overline{t}^k \overline{u}^m$, $\overline{\varphi}(\overline{u}) = \overline{t}^l \overline{u}^n$, the automorphism $\overline{\varphi}$ is represented by the matrix

$$K = \begin{pmatrix} k & l \\ m & n \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Z}).$$

Note that $M\alpha(\overline{z}) = \alpha(\overline{\varphi}(\overline{z}))M$ for all \overline{z} in E/\mathbb{Z}^2 , where $\alpha \colon \mathbb{Z}^2 \to \operatorname{GL}_2(\mathbb{Z})$ is defined by $\overline{t}^e \overline{u}^f \mapsto (-I)^e A^f$. Taking $\overline{z} = \overline{t}$ gives $-M = (-I)^k A^m M$. Hence m must be zero, implying both $k, n \in \{\pm 1\}$. Note that this already shows that $\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}$ is characteristic in E. In case k = 1 or n = 1, we have $R(\overline{\varphi}) = \infty$, implying $R(\varphi) = \infty$ as well. So assume k = n = -1. Applying the condition on \overline{u} gives $MA = (-I)^l A^n M$, hence $MA = \pm A^{-1}M$. Moreover, the identity

$$I - K = \begin{pmatrix} 1 - k & -l \\ -m & 1 - n \end{pmatrix} = \begin{pmatrix} 2 & -l \\ 0 & 2 \end{pmatrix}$$

implies that the classes $[1]_{\overline{\varphi}}$, $[\overline{t}]_{\overline{\varphi}}$, $[\overline{u}]_{\overline{\varphi}}$ and $[\overline{tu}]_{\overline{\varphi}}$ are exactly the Reidemeister classes of $\overline{\varphi}$. The result now follows from the addition formula.

If A has infinite order, it has either repeated eigenvalue ± 1 or its eigenvalues are real and different from ± 1 . In the former case, we have:

PROPOSITION 5.16. Let $E = (\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}$, where $A = \psi_{|\mathbb{Z}^2}$ has infinite order. If A has repeated eigenvalue 1 or -1, then E has the R_{∞} property.

PROOF. Write $\varepsilon \in \{\pm 1\}$ the repeated eigenvalue of A. Changing bases if necessary, we may assume that $A = \begin{pmatrix} \varepsilon & r \\ 0 & \varepsilon \end{pmatrix}$, for some $r \neq 0 \in \mathbb{Z}$. Clearly A and $-A^{-1}$ have different eigenvalues. As a result, they are not conjugate, that is, no $M \in \operatorname{GL}_2(\mathbb{Z})$ satisfies $MA = -A^{-1}M$. By Lemma 5.15, it suffices to show that $MA = A^{-1}M$ implies $R(M) = \infty$ for all $M \in \operatorname{GL}_2(\mathbb{Z})$. To this end, take M in $\operatorname{GL}_2(\mathbb{Z})$ satisfying AMA = M. Writing $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, this means

$$\begin{pmatrix} a + \varepsilon rc & b + r^2c + \varepsilon r(a+d) \\ c & d + \varepsilon rc \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

Recall $r \neq 0$, hence c = 0 and subsequently a + d = 0. So either a or d is 1, implying $R(M) = \infty$.

The situation where A has real eigenvalues is more subtle.

PROPOSITION 5.17. Let $E = (\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}$, where $A = \psi_{|\mathbb{Z}^2}$ has real eigenvalues $\neq \pm 1$. Then the Reidemeister spectrum of E is either $\{\infty\}$ or $\{8, \infty\}$. Moreover, if $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ has the R_{∞} property, then so does E.

PROOF. By Lemma 5.15, only ∞ and R(M) + R(-M) + R(AM) + R(-AM) are candidate Reidemeister numbers, where $M \in \text{GL}_2(\mathbb{Z})$ satisfies $MA = \pm A^{-1}M$.

In [8], Gonçalves and Wong analysed the condition $MA = A^{-1}M$, see Section 4.1. They concluded that $MA = A^{-1}M$ can only hold if det(A) = 1 and $\operatorname{Tr}(M) = 0$. In particular, R(M) is either 2 or ∞ , and $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ has the R_{∞} property precisely when only $R(M) = \infty$ occurs. Moreover, if M satisfies the condition $MA = A^{-1}M$, the matrices -M, AM, -AM satisfy this condition as well. Therefore, the sum R(M) + R(-M) + R(-AM) + R(-AM) is always infinite if $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ has the R_{∞} property, and either 8 or ∞ otherwise.

We now repeat their argument for the condition $MA = -A^{-1}M$. So, take $M \in \operatorname{GL}_2(\mathbb{Z})$ satisfying $MA = -A^{-1}M$. Setting $\delta := \det(A)$, let λ and $\delta \lambda^{-1}$ denote the eigenvalues of A. As $\lambda \neq \pm 1$, these eigenvalues are different. In particular A is diagonalizable over \mathbb{C} . Take P in $\operatorname{GL}_2(\mathbb{C})$ such that

$$PAP^{-1} = \begin{pmatrix} \lambda & 0\\ 0 & \delta\lambda^{-1} \end{pmatrix}$$
 and write $PMP^{-1} = \begin{pmatrix} a & c\\ b & d \end{pmatrix}$.

Denoting $\widetilde{A} := PAP^{-1}$ and $\widetilde{M} := PMP^{-1}$, we still have the relation $\widetilde{M}\widetilde{A} = -\widetilde{A}^{-1}\widetilde{M}$. Equating these matrices explicitly gives

$$\begin{pmatrix} \lambda a & \delta \lambda^{-1}c \\ \lambda b & \delta \lambda^{-1}d \end{pmatrix} = \begin{pmatrix} -\lambda^{-1}a & -\lambda^{-1}c \\ -\delta \lambda b & -\delta \lambda d \end{pmatrix}.$$

As $\lambda \neq \pm i$, this implies a = d = 0. Hence, $\operatorname{Tr}(M) = 0$. As before, this implies $\operatorname{Tr}(AM) = 0$ as well. Note that when $\det(A) = 1$, the relation above forces both b and c to be zero, so that M would be the zero matrix. Thus $\det(A) = -1$. In the odd case of Proposition 5.14 we showed that this implies either $R(M) = \infty$ or $R(AM) = \infty$. In particular, the sum R(M) + R(-M) + R(AM) + R(-AM) is always infinite, and the proof is complete.

Proposition 5.17 prompts the question: If $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ does not have the R_{∞} property, does *E* admit an automorphism having Reidemeister number 8? Examples 5.18 and 5.19 below will show that the answer to this question depends on the situation.

In general, E admits an automorphism having Reidemeister number 8 if there exists $M \in \mathrm{SL}_2(\mathbb{Z})$, satisfying $MA = A^{-1}M$, that extends to an automorphism ϕ of E having finite Reidemeister number. As in fact $G := \mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}$ is characteristic in $E = G \rtimes_{\psi} \mathbb{Z}$ by Lemma 5.15, any such ϕ is of the form

$$\phi(gu^k) = \varphi(g)(g_0u^{-1})^k, \quad g \in G, \ k \in \mathbb{Z},$$

for some g_0 in G and φ in Aut(G), where in turn

$$\varphi(zt^k) = M(z)(z_0t^{-1})^k, \quad z \in \mathbb{Z}^2, \ k \in \mathbb{Z},$$

for some z_0 in \mathbb{Z}^2 . It is easy to check that ϕ is an automorphism if and only if $\mu(g_0) \circ \varphi \circ \psi = \psi^{-1} \circ \varphi$, where $\mu(a) \colon G \to G$ is defined by $g \mapsto a^{-1} g a$. Writing $g_0 = m_0 t^m$ for some $m_0 \in \mathbb{Z}^2$, $m \in \mathbb{Z}$, this implies in particular that for all $z \in \mathbb{Z}^2$,

$$((-1)^m MA)(z) = \mu(g_0)(MA(z)) = A^{-1}M(z).$$

As $MA = A^{-1}M$, the above shows that m is even, and that the condition $\mu(g_0) \circ \varphi \circ \psi = \psi^{-1} \circ \varphi$ is satisfied on \mathbb{Z}^2 . Hence, ϕ is an automorphism if and only if $(\mu(g_0) \circ \varphi \circ \psi)(t) = (\psi^{-1} \circ \varphi)(t)$, or equivalently, $(\mu(m_0) \circ \varphi \circ \psi)(t) = (\psi^{-1} \circ \varphi)(t)$ as m is even. Recalling $\psi(t) = n_0 t$, this means $(M + A^{-1})(n_0) = 2m_0 + (A^{-1} - I)(z_0)$, or, equivalently,

(5.5)
$$(I + AM)(n_0) = 2A(m_0) + (I - A)(z_0).$$

So such ϕ exists exactly when we can find $m_0, z_0 \in \mathbb{Z}^2$ satisfying relation (5.5). Clearly, we can find such m_0, z_0 in the following situations:

- If $n_0 = 0$, that is, E is actually a semidirect product $\mathbb{Z}^2 \rtimes \mathbb{Z}^2$.
- If Tr(A) = 3 or 5, since then I A is invertible.

A less trivial example where we always can find appropriate m_0 , z_0 is the following:

EXAMPLE 5.18. Suppose that $A = \begin{pmatrix} 2 & 3 \\ 3 & 5 \end{pmatrix}$, and set $M := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. It is easy to check that $M \in SL_2(\mathbb{Z})$ satisfies AMA = M, hence $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ does not have the R_{∞} property. Moreover, one easily verifies that $Im(2A) + Im(I - A) = \mathbb{Z}^2$. Hence, suitable $m_0, z_0 \in \mathbb{Z}^2$ exist for any $n_0 \in \mathbb{Z}^2$. By the discussion preceding this example, we conclude that E does not have the R_{∞} property either.

In contrast, the following example shows that the Reidemeister spectrum of E may depend on n_0 .

EXAMPLE 5.19. Suppose that $A = \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$. The matrix M from Example 5.18 shows that $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ does not have the R_{∞} property. In general, recall that any $M \in \mathrm{SL}_2(\mathbb{Z})$ satisfying AMA = M can be written as $M = \begin{pmatrix} m & n \\ p & -m \end{pmatrix}$, where m, n and p satisfy system (4.1). In this particular situation, this system reads:

$$\begin{cases} -m^2 - pn = 1, \\ 4m + 2p + 2n = 0. \end{cases}$$

From the bottom equation -2m = p + n, it is easy to infer that either both p and n are odd, or both p and n are even. Suppose that both p and n are even. Then m is odd by the top equation, so -m = p/2 + n/2 moreover implies that p or n is a quadruple. Consequently, $pn \equiv 0 \mod 8$, thus $-1 \equiv m^2 \mod 8$ is a square modulo 8. This is false, hence p and n must be odd, implying m to be even. We conclude that all entries of I + M are odd. Of course, this holds equally for the matrix I + AM, as AM satisfies A(AM)A = AM as well. As all entries of I - A are even, equation (5.5) allows a solution (m_0, z_0) precisely when the sum of the components of n_0 is even. Equivalently, E has the R_{∞} property precisely when the sum of the components of n_0 is odd.

5.2.3. Conclusion. We summarise our findings in the following table:

eigenvalues of $A := \psi _{\mathbb{Z}^2}$	$\operatorname{Spec}_{\mathrm{R}}((\mathbb{Z}^2 \rtimes_{-I} \mathbb{Z}) \rtimes_{\psi} \mathbb{Z})$
1, 1 or -1, -1	$\{\infty\}$
$\lambda,-\lambda^{-1},\;\lambda\in\mathbb{R}$	$\{\infty\}$
$\lambda, \lambda^{-1}, \ \lambda \in \mathbb{R} \setminus \{\pm 1\}$	$\{\infty\}$ or $\{8,\infty\}$

5.3. The semidirect product $H_n \rtimes_{\psi} \mathbb{Z}$. Let us first fix some notations we will use throughout this subsection. As before, we will write elements of H_n as expressions of the form $x^a y^b z^c$, where z generates the center of H_n and $[y, x] = z^n$. Denote the projection map $H_n \to H_n/Z(H_n)$ by $h \mapsto \overline{h}$. The quotient $H_n/Z(H_n)$ is generated by \overline{x} and \overline{y} , and moreover $\langle \overline{x}, \overline{y} \rangle \cong \mathbb{Z}^2$. Let $A \in \operatorname{GL}_2(\mathbb{Z})$ represent the induced automorphism $\overline{\psi}$ relative to the basis $\{\overline{x}, \overline{y}\}$. Again, we have:

LEMMA 5.20. If A does not have 1 as eigenvalue, the subgroup H_n of $H_n \rtimes_{\psi} \mathbb{Z}$ is characteristic.

PROOF. Write $E = H_n \rtimes_{\psi} \mathbb{Z}$. Since A does not have 1 as eigenvalue, Im(I - A) has finite index in $H_n/Z(H_n)$. Moreover, one easily checks that Im $(I - A) = \overline{[E, E]}$, so $\overline{[E, E]}$ has finite index in $H_n/Z(H_n)$ as well. Hence [E, E] has finite index in H_n and the result follows.

In fact, if A does have eigenvalue 1, we already studied the Reidemeister spectrum of $H_n \rtimes_{\psi} \mathbb{Z}$. Indeed, if A has repeated eigenvalue 1, the group $H_n \rtimes_{\psi} \mathbb{Z}$

is nilpotent; if A has eigenvalues 1 and -1, the group $H_n \rtimes_{\psi} \mathbb{Z}$ is an extension of \mathbb{Z}^2 by \mathbb{Z}^2 .

LEMMA 5.21. If A has eigenvalues 1 and -1, the group $H_n \rtimes_{\psi} \mathbb{Z}$ is an extension of \mathbb{Z}^2 by \mathbb{Z}^2 .

PROOF. Write $E = H_n \rtimes_{\psi} \mathbb{Z}$. The matrix I - A has eigenvalues 0 and 2. Consequently, $\operatorname{Im}(I - A) \cong \mathbb{Z}$, say $\operatorname{Im}(I - A) = \langle \overline{h} \rangle$ for some $h \in H_n$. Take generators $\overline{v}, \overline{w}$ of $H_n/Z(H_n)$ such that $\overline{v}^k = \overline{h}$ for some $k \in \mathbb{Z}$, or equivalently, $\operatorname{Im}(I - A) \subseteq \langle \overline{v} \rangle$. Since $\overline{[E, E]} = \operatorname{Im}(I - A)$, also $\overline{[E, E]} \subseteq \langle \overline{v} \rangle$, that is, $[E, E] \subseteq \langle v, z \rangle$. It easily follows that $\langle v, z \rangle \lhd E$ and that $E/\langle v, z \rangle \cong \mathbb{Z}^2$. Clearly, also $\langle v, z \rangle \cong \mathbb{Z}^2$, and the proof is finished. \Box

So it remains to study $H_n \rtimes_{\psi} \mathbb{Z}$ when A does not have 1 as eigenvalue.

PROPOSITION 5.22. If $A \neq -I$ does not have 1 as eigenvalue, the group $H_n \rtimes_{\psi} \mathbb{Z}$ has the R_{∞} property.

PROOF. The center $Z(H_n)$ is characteristic in $H_n \rtimes_{\psi} \mathbb{Z}$ and $(H_n \rtimes_{\psi} \mathbb{Z})/Z(H_n) \cong \mathbb{Z}^2 \rtimes_A \mathbb{Z}$. Therefore, if $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$ has the R_∞ property, $H_n \rtimes_{\psi} \mathbb{Z}$ has the R_∞ property as well. It thus remains to prove the proposition when A has real eigenvalues and det(A) = 1. To this end, take an automorphism ϕ of $H_n \rtimes_{\psi} \mathbb{Z}$. If ϕ induces the identity on $(H_n \rtimes_{\psi} \mathbb{Z})/H_n \cong \mathbb{Z}$, clearly $R(\phi)$ is infinite. So assume $\phi(t) = h_0 t^{-1}$ for some $h_0 \in H_n$. Let φ denote the restriction of ϕ to H_n . Further, let $\overline{\varphi}$ denote the induced automorphism on the quotient $H_n/Z(H_n)$, say $\overline{\varphi}$ is represented by $M \in \operatorname{GL}_2(\mathbb{Z})$. The addition formula implies that $R(\varphi) = R(\overline{\varphi})R(\varphi|_{Z(H_n)})$.

From the condition $\phi(th) = \phi(t)\phi(h)$ for all $h \in H_n$, it easily follows that $MA = A^{-1}M$. Hence $R(\overline{\varphi}) = \infty$ when $\det(M) = -1$.

At the same time, $\varphi|_{Z(H_n)}$ is multiplication by det(M), hence $R(\varphi|_{Z(H_n)}) = \infty$ when det(M) = 1. We conclude that $R(\varphi)$ is always infinite, so $R(\phi) = \infty$ as well.

In contrast, when A = -I, more is possible:

PROPOSITION 5.23. Let E denote the group $H_n \rtimes_{\psi} \mathbb{Z}$ where $\psi(x) = x^{-1}z^k$ and $\psi(y) = y^{-1}z^l$ for some $k, l \in \mathbb{Z}$.

- (a) If n is odd, E has Reidemeister spectrum $4\mathbb{N} \cup \{\infty\}$.
- (c) If n is even, E has Reidemeister spectrum 4N ∪ {∞} if both k and l are even, and Reidemeister spectrum 8N ∪ {∞} if k or l is odd.

PROOF. We first show that in either case, the Reidemeister spectrum of E is contained in $4\mathbb{N} \cup \{\infty\}$. To this end, let ϕ be an automorphism of E. We may assume that $\phi(t) = h_0 t^{-1}$ for some $h_0 \in H_n$. As before, write φ the restriction to H_n and let $\overline{\varphi}$ denote the induced automorphism on the quotient $H_n/Z(H_n)$.

Further, let $M \in \operatorname{GL}_2(\mathbb{Z})$ denote the matrix representing $\overline{\varphi}$ relative to the basis $\{\overline{x}, \overline{y}\}$. If det(M) = 1, then φ induces the identity on $Z(H_n)$, hence $R(\phi) = \infty$. Furthermore, if det(M) = -1, the addition formula says that $R(\phi) = 4 |\operatorname{Tr}(M)|$ if $\operatorname{Tr}(M)$ is nonzero, and $R(\phi) = \infty$ otherwise. So $R(\varphi) \in 4\mathbb{N} \cup \{\infty\}$.

Next, we determine which M arise as induced automorphisms on $H_n/Z(H_n)$, or more generally, which φ can be extended to an automorphism on E. As elements in $\operatorname{SL}_2(\mathbb{Z})$ always contribute ∞ to $\operatorname{Spec}_{\mathbb{R}}(E)$, we only consider M having $\det(M) = -1$. So, let $\varphi \in \operatorname{Aut}(H_n)$ and write $\varphi(x) = x^a y^c z^m$, $\varphi(y) = x^b y^d z^p$ for some $m, p \in \mathbb{Z}$ and

(5.6)
$$M := \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{GL}_2(\mathbb{Z}), \quad \text{where } \det(M) = -1.$$

Then φ extends to an automorphism $\phi \colon E \to E$ defined by $ht^r \mapsto \varphi(h)(h_0t^{-1})^r$, $h_0 \in H_n$, if and only if all $h \in H_n$ satisfy $\phi(\psi(h)t) = \phi(t)\phi(h)$, that is,

$$(\varphi \circ \psi)(h) h_0 = h_0(\psi^{-1} \circ \varphi)(h).$$

Writing $h_0 = x^e y^f z^g$ (where g will not really matter as z^g is central), a simple calculation shows this to be equivalent to

(5.7)
$$2\binom{m}{p} + (I + M^T)\binom{k}{l} = n\binom{ac}{bd} + \binom{-c}{-d} \binom{e}{b}\binom{e}{f}.$$

If n is odd, any $M \in \operatorname{GL}_2(\mathbb{Z})$ allows $m, p, e, f \in \mathbb{Z}$ such that condition (5.7) above is satisfied. Indeed, since n is odd, 2 is invertible modulo n, so there exist $m, p \in \mathbb{Z}$ such that

$$2\binom{m}{p} + (I + M^T)\binom{k}{l} \in \binom{n\mathbb{Z}}{n\mathbb{Z}}, \quad \text{say} \quad 2\binom{m}{p} + (I + M^T)\binom{k}{l} = \binom{n\,\widetilde{m}}{n\widetilde{p}}$$

for some $\widetilde{m}, \widetilde{p} \in \mathbb{Z}$. Subsequently, setting

$$\begin{pmatrix} e \\ f \end{pmatrix} := \begin{pmatrix} -c & a \\ -d & b \end{pmatrix}^{-1} \begin{pmatrix} \widetilde{m} - ac \\ \widetilde{p} - bd \end{pmatrix}$$

results in a solution (m, p, e, f) to (5.7). Therefore, there exists for every matrix $M_r := \begin{pmatrix} r & 1 \\ 1 & 0 \end{pmatrix}, r \in \mathbb{N}$, an automorphism ϕ_r of E inducing M_r on $H_n/Z(H_n)$. By construction $R(\phi_r) = 4r$, thus E has Reidemeister spectrum $4\mathbb{N} \cup \{\infty\}$.

If n is even, we can similarly find suitable $m, p, e, f \in \mathbb{Z}$ provided

(5.8)
$$\left((I+M^T) \begin{pmatrix} k \\ l \end{pmatrix} \in \begin{pmatrix} 2\mathbb{Z} \\ 2\mathbb{Z} \end{pmatrix}, \right.$$

to guarantee the existence of m and p. Moreover, condition (5.7) readily implies that if n is even, (5.8) is in fact equivalent to the existence of such m, p, e and f.

Hence M lifts to an automorphism of E if and only if it satisfies condition (5.8) above. Therefore, the Reidemeister spectrum of E equals the set

 $\operatorname{Spec}_{\mathrm{R}}(E) = \{4|\operatorname{Tr}(M)| \mid M \text{ satisfies } (5.8), \ \det(M) = -1, \ \operatorname{Tr}(M) \neq 0\} \cup \{\infty\}.$

We distinguish four cases, based on the parity of k and l.

Case 1. If both k and l are even, condition (5.8) is trivially satisfied. As before, the matrices $M_r := \begin{pmatrix} r & 1 \\ 1 & 0 \end{pmatrix}$, $r \in \mathbb{N}$, show that E has Reidemeister spectrum $4\mathbb{N} \cup \{\infty\}$.

Case 2. Next, suppose that k is even, but l is odd. Let $M \in GL_2(\mathbb{Z})$ satisfy condition (5.8), write M as in (5.6). Condition (5.8) implies that both c and 1 + d are even, that is, c is even and d is odd. As M is invertible, a must be odd as well. Hence Tr(M) = a + d is even, showing $Spec_R(E) \subseteq 8\mathbb{N} \cup \{\infty\}$.

Conversely, for $r \in \mathbb{N}$ even, the matrix $M_r := \begin{pmatrix} r+1 & r/2 \\ -2 & -1 \end{pmatrix}$ satisfies condition (5.8), has $\det(M_r) = -1$ and moreover $\operatorname{Tr}(M_r) = r$. Thus in this case, indeed $\operatorname{Spec}_{\mathbf{R}}(E) = 8\mathbb{N} \cup \{\infty\}$.

Case 3. Note that x and y play quite a symmetric role: we only use that $H_n/Z(H_n) = \langle \overline{x}, \overline{y} \rangle$ and that $[y, x] = z^n$, where $Z(H_n) = \langle z \rangle$. Consequently, if k is odd and l is even, we can reduce to Case 2 by swapping x and y and replacing z by z^{-1} .

Case 4. Similarly, if both k and l are odd, replacing x by yx transforms this case into Case 2, as $\psi(yx) = (yx)^{-1}z^{k+l+n}$ and n is even, and [y, yx] = [y, x]. \Box

5.3.1. Conclusion. We summarise our findings in the following table:

ψ		$\operatorname{Spec}_{\mathrm{R}}(H_n \rtimes_{\psi} \mathbb{Z})$
$A := \psi _{\mathbb{Z}^2} \neq -I \text{ and } 1 \notin \Lambda(A)$		$\{\infty\}$
$\psi(x) = x^{-1} z^k, \ \psi(y) = y^{-1} z^l$	(k and l even) or (n odd) (k or l odd) and (n even)	$4\mathbb{N} \cup \{\infty\}\\ 8\mathbb{N} \cup \{\infty\}$

References

- L.S. CHARLAP, Bieberbach Groups and Flat Manifolds, Universitext, Springer-Verlag, New York Inc., 1986.
- [2] K. DEKIMPE, A users' guide to infra-nilmanifolds and almost-Bieberbach groups, Handbook of Group Actions, Vol. III, Adv. Lect. Math. (ALM), 40, Int. Press, Somerwille, MA, 2018, 215–262.
- [3] K. DEKIMPE, B. DE ROCK AND P. PENNINCKX, The R_∞ property for infra-nilmanifolds, Topol. Methods Nonlinear Anal. 34 (2009), no. 2, 353–374.
- [4] K. DEKIMPE AND D. GONÇALVES, The R_∞ property for free groups, free nilpotent groups and free solvable groups, Bull. Lond. Math. Soc. 46 (2014), no. 4, 737–746.

- [5] A. FEL'SHTYN AND R. HILL, Dynamical zeta functions, congruences in Nielsen theory and Reidemeister torsion, Nielsen theory and Reidemeister torsion (Warsaw, 1996), vol. 49, Banach Center Publ., vol. 49, Polish Acad. Sci., Warsaw, 1999, 77–116.
- [6] A. FEL'SHTYN AND T. NASYBULLOV, The R_{∞} and S_{∞} properties for linear algebraic groups, J. Group Theory **19** (2016), no. 5, 901–921.
- [7] A. FEL'SHTYN AND E. TROITSKY, Aspects of the property R_{∞} , J. Group Theory 18 (2015), no. 6, 1021–1034.
- [8] D. GONÇALVES AND P. WONG, Twisted conjugacy classes in exponential growth groups, Bull. London Math. Soc. 35 (2003), no. 2, 261–268.
- D. GONÇALVES AND P. WONG, Twisted conjugacy classes in nilpotent groups, J. Reine Angew. Math. 633 (2009), 11–27.
- [10] D. GONÇALVES AND P. WONG, Nielsen numbers of selfmaps of Sol 3-manifolds, Topology Appl. 159 (2012), no. 18, 3729–3737.
- [11] D.L. GONÇALVES AND P.N.-S. WONG, Twisted conjugacy for virtually cyclic groups and crystallographic groups, Combinatorial and Geometric Group Theory, Trends in Mathematics, Birkhaüser, 2010, 119–147.
- [12] K.Y. HA, J.B. LEE AND P. PENNINCKX, Formulas for the Reidemeister, Lefschetz and Nielsen coincidence number of maps between infra-nilmanifolds, Fixed Point Theory Appl. 2012 (2012), https://doi.org/10.1186/1687-1812-2012-39.
- [13] J. JEZIERSKI AND W. MARZANTOWICZ, Homotopy Methods in Topological Fixed and Periodic Point Theory, Topological Fixed Point Theory and its Applications, vol. 3, Springer, 2006.
- [14] B. JIANG, Nielsen Fixed Point Theory, Contemp. Math., vol. 14, Amer. Math. Soc. 1983.
- [15] T. KIANG, The Theory of Fixed Point Classes, Springer-Verlag, 1989.
- [16] J.B. LEE AND K.B. LEE, Averaging formula for Nielsen numbers of maps on infrasolvmanifolds of type (R), Nagoya Math. J. 196 (2009), 117–134.
- [17] M. NEWMAN, Integral Matrices, Pure and Applied Mathematics, vol. 45, Academic Press, New York, London, 1972.
- [18] A.L. ONISHCHIK AND E.B. VINBERG Lie Groups and Lie Algebras I, Encyclopedia of Mathematics, vol. I, Springer–Verlag, Berlin, Heidelberg, New York, 1993.
- [19] V. ROMAN'KOV, Twisted conjugacy classes in nilpotent groups, J. Pure Appl. Algebra 215 (2011), no. 4, 664–671.
- [20] A. SZCZEPAŃSKI, Geometry of crystallographic groups, Algebra and Discrete Mathematics, vol. 4, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2012.
- [21] K.I. TAHARA, On the finite subgroups of GL(3, Z), Nagoya Math. J. 41 (1971), 169–209.
- [22] J.A. WOLF, Spaces of constant curvature, Publish or Perish, Inc. Berkeley, 1977.

Manuscript received November 22, 2017 accepted January 20, 2018

KAREL DEKIMPE, SAM TERTOOY AND IRIS VAN DEN BUSSCHE KU Leuven Campus Kulak Kortrijk E. Sabbelaan 53 8500 Kortrijk, BELGIUM

E-mail address: karel.dekimpe@kuleuven.be sam.tertooy@kuleuven.be iris.vandenbussche@kuleuven.be

TMNA: Volume 53 – 2019 – N^o 2