

# *Algebra & Number Theory*

Volume 5

2011

No. 8

**The Picard group of a  $K3$  surface and  
its reduction modulo  $p$**

Andreas-Stephan Elsenhans and Jörg Jahnel



**mathematical sciences publishers**

# The Picard group of a $K3$ surface and its reduction modulo $p$

Andreas-Stephan Elsenhans and Jörg Jahnel

We present a method to compute the geometric Picard rank of a  $K3$  surface over  $\mathbb{Q}$ . Contrary to a widely held belief, we show that it is possible to verify Picard rank 1 using reduction at a single prime.

## 1. Introduction

**1.1.** For complex, projective  $K3$  surfaces, the Picard group is a highly interesting invariant. In general, it is isomorphic to  $\mathbb{Z}^n$  for some  $n = 1, \dots, 20$ . A generic  $K3$  surface has Picard rank 1. Nevertheless, the first explicit examples of  $K3$  surfaces over  $\mathbb{Q}$  having geometric Picard rank 1 were constructed by R. van Luijk [2007] as late as 2004. Van Luijk's method is based on reduction modulo  $p$ . It works as follows.

**Approach 1.2** (van Luijk). Let  $S$  be a  $K3$  surface over  $\mathbb{Q}$ .

- (i) At a place  $p$  of good reduction, the Picard group  $\text{Pic}(S_{\overline{\mathbb{Q}}})$  of the surface injects into the Picard group  $\text{Pic}(S_{\overline{\mathbb{F}}_p})$  of its reduction modulo  $p$ .
- (ii) On its part, the group  $\text{Pic}(S_{\overline{\mathbb{F}}_p})$  injects into the second étale cohomology group  $H_{\text{ét}}^2(S_{\overline{\mathbb{F}}_p}, \mathbb{Q}_l(1))$ .
- (iii) Only roots of unity can arise as eigenvalues of the Frobenius  $\text{Frob}$  on the image of  $\text{Pic}(S_{\overline{\mathbb{F}}_p})$  in  $H_{\text{ét}}^2(S_{\overline{\mathbb{F}}_p}, \mathbb{Q}_l(1))$ . The number of eigenvalues of this form, counted with multiplicities, is therefore an upper bound for the Picard rank of  $S_{\overline{\mathbb{F}}_p}$ . One may compute the eigenvalues of  $\text{Frob}$  by counting the points on  $S$ , defined over  $\mathbb{F}_p$  and some finite extensions.

Doing this for one prime, one obtains an upper bound for  $\text{rk Pic}(S_{\overline{\mathbb{F}}_p})$ , which is always even. The Tate conjecture asserts that this bound is actually sharp.

---

Esenhans was supported in part by the Deutsche Forschungsgemeinschaft (DFG) through a funded research project.

*MSC2010:* primary 14C22; secondary 14D15, 14J28, 14Q10.

*Keywords:*  $K3$  surface, Picard group, Picard scheme, deformation, Artin approximation, Van Luijk's method.

Therefore, the best that could happen is to find a prime  $p$  that yields an upper bound of 2 for the rank of  $\text{Pic}(S_{\overline{\mathbb{Q}}})$ .

- (iv) In this case, the assumption that the surface has Picard rank 2 over  $\overline{\mathbb{Q}}$  implies that the discriminants of both Picard groups,  $\text{Pic}(S_{\overline{\mathbb{Q}}})$  and  $\text{Pic}(S_{\overline{\mathbb{F}}_p})$ , belong to the same square class. Note here that reduction modulo  $p$  respects the intersection pairing.
- (v) To obtain a contradiction, one combines information from two primes. It may happen that one has a rank bound of 2 at both places but that different square classes arise for the discriminants. Then, these data are incompatible with Picard rank 2 over  $\overline{\mathbb{Q}}$ . Geometric Picard rank 1 is proven.

**1.3. The improvement.** The idea behind [Approach 1.2](#) is to consider the specialization  $\text{sp} : \text{Pic}(S_{\overline{\mathbb{Q}}}) \hookrightarrow \text{Pic}(S_{\overline{\mathbb{F}}_p})$  as an injection of lattices. Then, the two possibilities  $\text{rk Pic}(S_{\overline{\mathbb{Q}}}) < \text{rk Pic}(S_{\overline{\mathbb{F}}_p})$  and  $\text{rk Pic}(S_{\overline{\mathbb{Q}}}) = \text{rk Pic}(S_{\overline{\mathbb{F}}_p})$  are distinguished. In the latter, the standard fact is used that  $\text{disc Pic}(S_{\overline{\mathbb{Q}}}) / \text{disc Pic}(S_{\overline{\mathbb{F}}_p})$  is a perfect square.

We will show in this article that the assertion for the second case may be refined to  $\text{disc Pic}(S_{\overline{\mathbb{Q}}}) = \text{disc Pic}(S_{\overline{\mathbb{F}}_p})$ . More precisely, we shall prove that, at least for  $p \neq 2$ , the cokernel of  $\text{sp} : \text{Pic}(S_{\overline{\mathbb{Q}}}) \hookrightarrow \text{Pic}(S_{\overline{\mathbb{F}}_p})$  is always torsion-free. This is true actually in a by-far more general situation than just for  $K3$  surfaces.

**Theorem 1.4.** *Let  $R$  be a discrete valuation ring with quotient field  $K$  of characteristic 0 and residue field  $k$  of characteristic  $p > 0$ . Further, let  $\pi : X \rightarrow \text{Spec } R$  be a morphism of schemes that is proper and smooth.*

*Suppose that  $R$  is of ramification degree  $e < p - 1$  and that  $k$  is perfect. Then, the cokernel of the specialization homomorphism  $\text{sp}_{\overline{k}} : \text{Pic}(X_{\overline{k}}) \rightarrow \text{Pic}(X_{\overline{k}})$  is torsion-free.*

**Remarks 1.5.** (a) In the applications, we will have  $R = \mathbb{Z}_{(p)} \subset \mathbb{Q}$ . Then, the assumption simply means  $p \neq 2$ .

- (b) We will show this theorem in [Section 3](#). As an application, one may prove  $\text{rk Pic}(S_{\overline{\mathbb{Q}}}) = 1$  for a  $K3$  surface  $S$  using its reduction at a single prime. This works as follows.

**Approach 1.6.** Let a  $K3$  surface  $S$  over  $\mathbb{Q}$  be given.

- (i) For a prime  $p \neq 2$  of good reduction, perform steps (i), (ii) and (iii) as in [Approach 1.2](#). Thereby, the hope is to prove  $\text{rk Pic}(S_{\overline{\mathbb{F}}_p}) = 2$ . Further, compute the discriminant giving two explicit generators.

Alternatively, to determine the discriminant, one might use the Artin–Tate formula [[Milne 1975](#)]. In this case,  $\text{rk Pic}(S_{\overline{\mathbb{F}}_p}) = 2$  is shown only relative to the Tate conjecture. Observe, however, that a surface with  $\text{rk Pic}(S_{\overline{\mathbb{F}}_p}) = 1$ , due to a failure of the Tate conjecture, would serve our purposes as well.

- (ii) Assume  $\text{rk Pic}(S_{\overline{\mathbb{Q}}}) = 2$ . Then, according to [Theorem 1.4](#), every invertible sheaf on  $S_{\overline{\mathbb{F}}_p}$  must lift to  $S_{\overline{\mathbb{Q}}}$ . Estimate the degree of a hypothetical effective divisor. Finally, use Gröbner bases to verify that such a divisor does not exist.

**Example 1.7.** Consider the  $K3$  surface  $S$  over  $\mathbb{Q}$  given by

$$w^2 = x^5y + x^4y^2 + 2x^3y^3 + x^2y^4 + xy^5 + 4y^6 + 2x^5z + 2x^4z^2 + 4x^3z^3 + 2xz^5 + 4z^6.$$

Then,  $\text{rk Pic}(S_{\overline{\mathbb{Q}}}) = 1$ .

*Proof.* For the reduction of  $S$  at the prime 5, one sees that the branch locus has a tritangent line given by  $z - 2y = 0$ . It meets the branch locus at  $(1 : 0 : 0)$ ,  $(1 : 3 : 1)$ , and  $(0 : 1 : 2)$ .

The numbers of points on  $S$  over  $\mathbb{F}_{5^d}$  are, in this order, 41, 751, 15 626, 392 251, 9 759 376, 244 134 376, 6 103 312 501, 152 589 156 251, 3 814 704 296 876, and 95 367 474 609 376. Thus, the traces of Frob on  $H_{\text{ét}}^2(S_{\overline{\mathbb{F}}_5}, \overline{\mathbb{Q}}_l)$  are 15, 125, 0, 1 625,  $-6 250$ ,  $-6 250$ ,  $-203 125$ , 1 265 625, 7 031 250, and 42 968 750.

Elsenhans and Jahnel [[2008a](#), Algorithm 23] show that the sign in the functional equation is positive. The characteristic polynomial of Frob is therefore completely determined. For its decomposition into prime polynomials, we find (after Tate twist to  $H_{\text{ét}}^2(S_{\overline{\mathbb{F}}_5}, \overline{\mathbb{Q}}_l(1))$ )

$$\begin{aligned} \frac{1}{5}(t-1)^2(5t^{20} - 5t^{19} - 5t^{18} + 10t^{17} - 2t^{16} - 3t^{15} + 4t^{14} - 2t^{13} - 2t^{12} + t^{11} \\ + 3t^{10} + t^9 - 2t^8 - 2t^7 + 4t^6 - 3t^5 - 2t^4 + 10t^3 - 5t^2 - 5t + 5). \end{aligned}$$

This shows  $\text{rk Pic}(S_{\overline{\mathbb{F}}_5}) \leq 2$ .

The irreducible components of the pull-back of the tritangent line are explicit generators for  $\text{Pic}(S_{\overline{\mathbb{F}}_5})$ . Such a component  $l$ , because it is a projective line, has self-intersection number  $l^2 = -2$ . Further,  $lh = 1$  for  $h$  the pull-back of a line. If we had  $\text{rk Pic}(S_{\overline{\mathbb{Q}}}) = 2$ , then the invertible sheaf  $\mathcal{O}(l)$  would lift to  $S_{\overline{\mathbb{Q}}}$ . We would have a divisor  $L$  on  $S_{\overline{\mathbb{Q}}}$  such that  $HL = 1$  and  $L^2 = -2$ . By [[Barth et al. 1984](#), Proposition VIII.3.6.i], such a divisor is automatically effective.

The equation  $HL = 1$  shows that  $L$  is obtained from a line on  $\mathbf{P}^2$ , the pull-back of which splits into two components. This is possible only for a line tritangent to the branch locus. Algorithm 8 of [[Elsenhans and Jahnel 2008a](#)] shows, however, using Gröbner bases, that such a tritangent line does not exist.  $\square$

## 2. The cokernel of the restriction map

**Notation 2.1.** (i) Let  $R$  be a discrete valuation ring of unequal characteristic. We will write  $K := \text{Quot}(R)$  for its quotient field,  $\mathfrak{p}$  for the maximal ideal,  $k := R/\mathfrak{p}$  for the residue field of characteristic  $p$ , and  $v : K \rightarrow \mathbb{Z}$  for the normalized valuation. Let  $e := v(p)$  denote the ramification degree of  $R$ .

- (ii) Let  $X$  be an  $R$ -scheme. Then, we will write  $X_{\mathfrak{p}}$  for the special fiber and  $X_{\eta}$  for the generic fiber of  $X$ . For  $L$  an extension of  $K$ , we will denote by  $X_L$  the base extension of  $X_{\eta}$  to  $L$ . Analogously, for  $l$  an extension of  $k$ , we will write  $X_l$  for the base extension of  $X_{\mathfrak{p}}$  to  $l$ . In the particular case that  $l = \mathbb{F}_q$ , the shortcut  $X_q$  shall be used for  $X_l$ .

**Proposition 2.2.** *Let  $\pi : X \rightarrow \text{Spec } R$  be a morphism of schemes that is proper and flat. Suppose that the special fiber  $X_{\mathfrak{p}}$  is normal.*

*If  $R$  is complete and satisfies the condition  $e < p - 1$ , then the cokernel of the restriction homomorphism  $\text{Pic}(X) \rightarrow \text{Pic}(X_{\mathfrak{p}})$  is torsion-free.*

*Proof.* This result was obtained by M. Raynaud in the course of his investigations on the Picard scheme [Raynaud 1979, Théorème 4.1.2.1]. □

**Remark 2.3.** Assume, also, that the restriction homomorphism  $H^1(X, \mathcal{O}_X) \rightarrow H^1(X_{\mathfrak{p}}, \mathcal{O}_{X_{\mathfrak{p}}})$  is surjective. Then, the assertion of Proposition 2.2 may be established using the following elementary argument, which is also due to M. Raynaud [1979, section 1].

Consider the functors  $T^i$  on the category of all finitely generated  $R$ -modules to finitely generated  $R$ -modules, given by  $T^i(M) := H^i(X, \pi^* \tilde{M})$ . Here,  $\tilde{M}$  denotes the coherent sheaf associated with the  $R$ -module  $M$ . According to [Grothendieck 1963, Proposition (7.7.10), p. 71], the functor  $T^1$  is right exact. Hence, by [ibid., Théorème (7.7.5.II), p. 68],  $T^2$  is left exact. This, in turn, immediately implies that  $H^2(X, \mathcal{O}_X)$  is torsion-free.

Further, the short exact sequence

$$0 \rightarrow \mathcal{U}_1 \rightarrow \mathcal{O}_X^* \rightarrow \mathcal{O}_{X_{\mathfrak{p}}}^* \rightarrow 0$$

shows that  $\text{coker}(\text{Pic}(X) \rightarrow \text{Pic}(X_{\mathfrak{p}}))$  injects into  $H^2(X, \mathcal{U}_1)$ . Finally, as  $e < p - 1$ , the exponential map provides us with an isomorphism

$$\mathcal{O}_X \xrightarrow{\cdot p} p\mathcal{O}_X \xrightarrow{\text{exp}} \mathcal{U}_1.$$

**Remarks 2.4.** (i) The additional assumption of 2.3 is fulfilled in our applications.

- (ii) For prime-to- $p$  torsion, the assertion of Proposition 2.2 is true in a more general situation.

**Proposition 2.5.** *Let  $\pi : X \rightarrow \text{Spec } R$  be a proper morphism of schemes.*

*If  $R$  is Henselian, then the cokernel of the restriction homomorphism*

$$\text{Pic}(X) \rightarrow \text{Pic}(X_{\mathfrak{p}})$$

*has no prime-to- $p$  torsion.*

*Proof.* Let  $l \neq p$  be a prime number. We will show that there is no  $l$ -torsion. For this, we observe at first that, according to a consequence of the theorem on proper base change [Artin et al. 1973, Exp. XII, Corollaire 5.5.iii], the restriction morphism induces bijections

$$H_{\text{ét}}^1(X, \mu_l) \xrightarrow{\cong} H_{\text{ét}}^1(X_p, \mu_l) \quad \text{and} \quad H_{\text{ét}}^2(X, \mu_l) \xrightarrow{\cong} H_{\text{ét}}^2(X_p, \mu_l).$$

Because [Berthelot et al. 1971, Exp. X, diagramme (7.13.10)] the restriction homomorphisms on the Picard groups and étale cohomology commute with the Chern maps, we see that restriction induces a surjection  $\text{Pic}(X)_l \twoheadrightarrow \text{Pic}(X_p)_l$  and an injection  $\text{Pic}(X)/l \hookrightarrow \text{Pic}(X_p)/l$ .

Applied to the two commutative diagrams of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Pic}(X)_l & \longrightarrow & \text{Pic}(X) & \longrightarrow & P_X \longrightarrow 0 \\ & & \downarrow \twoheadrightarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \text{Pic}(X_p)_l & \longrightarrow & \text{Pic}(X_p) & \longrightarrow & P_{X_p} \longrightarrow 0, \\ \\ 0 & \longrightarrow & P_X & \xrightarrow{\cdot l} & \text{Pic}(X) & \longrightarrow & \text{Pic}(X)/l \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \hookrightarrow \\ 0 & \longrightarrow & P_{X_p} & \xrightarrow{\cdot l} & \text{Pic}(X_p) & \longrightarrow & \text{Pic}(X_p)/l \longrightarrow 0, \end{array}$$

the snake lemma now shows that the induced homomorphism

$$\text{coker}(\text{Pic}(X) \rightarrow \text{Pic}(X_p)) \rightarrow \text{coker}(P_X \rightarrow P_{X_p})$$

is a bijection, while

$$\text{coker}(P_X \rightarrow P_{X_p}) \xrightarrow{\cdot l} \text{coker}(\text{Pic}(X) \rightarrow \text{Pic}(X_p))$$

is injective. Consequently,  $\text{coker}(\text{Pic}(X) \rightarrow \text{Pic}(X_p))$  has no  $l$ -torsion. □

### 3. The cokernel of the specialization map

**3.1.** In this section, we continue to use the notation from 2.1. Let  $\pi : X \rightarrow \text{Spec } R$  be a morphism of schemes that is proper and smooth. We have the restriction homomorphisms

$$\text{Pic}(X_\eta) \leftarrow \text{Pic}(X) \rightarrow \text{Pic}(X_p).$$

As  $\pi$  is smooth, the arrow to the left is a bijection [Berthelot et al. 1971, Exp. X, App. 7.8]. Consequently, there is a natural homomorphism  $\text{sp} : \text{Pic}(X_\eta) \rightarrow \text{Pic}(X_p)$ , which is called the *specialization*.

**Lemma 3.2.** *Let  $\pi : X \rightarrow \text{Spec } R$  be a morphism of schemes that is proper and smooth.*

*If  $R$  is complete and satisfies the condition  $e < p - 1$ , then the cokernel of the specialization homomorphism  $\text{sp} : \text{Pic}(X_\eta) \rightarrow \text{Pic}(X_p)$  is torsion-free.*

*Proof.* The assertion follows directly from [Proposition 2.2](#). □

**3.3.** Let  $K'/K$  be an extension field equipped with a discrete valuation extending that on  $K$ . Denote by  $R'$  the discrete valuation ring and by  $k'$  the residue field. The morphism  $X \times_{\text{Spec } R} \text{Spec } R' \rightarrow \text{Spec } R'$ , obtained by base change, induces a specialization homomorphism  $\text{sp}_{K'} : \text{Pic}(X_{K'}) \rightarrow \text{Pic}(X_{k'})$ .

There are the following two applications.

- (i) Suppose  $R$  to be complete. Then, for every finite extension  $K'/K$ , there is a unique [\[Serre 1968, Chap. II, §2, Proposition 3\]](#) discrete valuation extending the valuation on  $K$ . The direct limit of the homomorphisms  $\text{sp}_{K'} : \text{Pic}(X_{K'}) \rightarrow \text{Pic}(X_{k'})$  is a natural homomorphism  $\text{sp}_{\bar{K}} : \text{Pic}(X_{\bar{K}}) \rightarrow \text{Pic}(X_{\bar{k}})$ , again called the specialization.
- (ii) For general  $R$ , fix an embedding  $\bar{K} \hookrightarrow \widehat{\bar{K}}$  of the algebraic closure of  $K$  into that of its completion. By functoriality, this embedding induces a homomorphism  $\text{Pic}(X_{\bar{K}}) \rightarrow \text{Pic}(X_{\widehat{\bar{K}}})$ . Composing with  $\text{sp}_{\widehat{\bar{K}}}$ , constructed in (i), one has a specialization homomorphism  $\text{sp}_{\bar{K}} : \text{Pic}(X_{\bar{K}}) \rightarrow \text{Pic}(X_{\bar{k}})$ .

**Proposition 3.4.** *Let  $\pi : X \rightarrow \text{Spec } R$  be a morphism of schemes that is proper and smooth.*

*Suppose  $R$  is complete and satisfies the condition  $e < p - 1$ , and let  $k$  be perfect. Then, the cokernel of the specialization homomorphism  $\text{sp}_{\bar{K}} : \text{Pic}(X_{\bar{K}}) \rightarrow \text{Pic}(X_{\bar{k}})$  is torsion-free.*

*Proof.* By [\[Serre 1968, Chap. III, §5, Corollaire 1 du Théorème 3\]](#),  $K$  has a unique maximal unramified extension  $K^{\text{nr}}$ , which is actually the filtered direct limit of all finite unramified extensions  $K'/K$ .

An unramified extension does not change the ramification degree. Hence, by [Lemma 3.2](#), the homomorphisms  $\text{sp}_{K'} : \text{Pic}(X_{K'}) \rightarrow \text{Pic}(X_{k'})$  have torsion-free cokernels. As the filtered direct limit is an exact functor, the same is true for  $\text{sp}_{K^{\text{nr}}} : \text{Pic}(X_{K^{\text{nr}}}) \rightarrow \text{Pic}(X_{\bar{k}})$ .

We claim that the specialization homomorphism  $\text{sp}_{\bar{K}}$  has the same image in  $\text{Pic}(X_{\bar{k}})$  as  $\text{sp}_{K^{\text{nr}}}$ . For this, let  $\mathcal{L} \in \text{Pic}(X_{\bar{K}})$ . The inertia group  $I := \text{Gal}(\bar{K}/K^{\text{nr}})$  sends  $\mathcal{L}$  to a finite orbit  $\{\mathcal{L}_1, \dots, \mathcal{L}_m\}$ . The specializations of  $\mathcal{L}_1, \dots, \mathcal{L}_m$  in  $\text{Pic}(X_{\bar{k}})$  are all the same. Therefore,

$$m \cdot \text{sp}_{\bar{K}}(\mathcal{L}) = \text{sp}_{\bar{K}}(\mathcal{L}^{\otimes m}) = \text{sp}_{\bar{K}}(\mathcal{L}_1 \otimes \dots \otimes \mathcal{L}_m) = \text{sp}_{K^{\text{nr}}}(\mathcal{L}_1 \otimes \dots \otimes \mathcal{L}_m),$$

since  $\mathcal{L}_1 \otimes \cdots \otimes \mathcal{L}_m$  is  $I$ -invariant. Hence,  $m \cdot \text{sp}_{\bar{K}}(\mathcal{L}) \in \text{im } \text{sp}_{K^{\text{nr}}}$ . As  $\text{sp}_{K^{\text{nr}}}$  has a torsion-free cokernel, we see that  $\text{sp}_{\bar{K}}(\mathcal{L}) \in \text{im } \text{sp}_{K^{\text{nr}}}$ , too. □

**Remark 3.5.** The argument above uses that  $\text{Pic}(X_L) = \text{Pic}(X_K)^{\text{Gal}(L/K)}$ . This equality is certainly not correct, in general. It is true as soon as  $Y(K) \neq \emptyset$  for every connected component  $Y$  of  $X$ .

As  $\pi$  is smooth, we indeed have  $Y(K^{\text{nr}}) \neq \emptyset$ . To see this, let  $s : \text{Spec } l \rightarrow Y_k$  be a point defined over a finite extension. By [Grothendieck 1967, Proposition (17.5.3), p. 68],  $s$  may be lifted to a morphism  $\text{Spf } S \rightarrow Y$  for  $S$  the corresponding unramified extension of  $R$ . Then [Grothendieck 1961, Théorème (5.4.1), p. 156] yields the desired point.

**Theorem 3.6.** *Let  $R$  be a discrete valuation ring with quotient field  $K$  of characteristic 0 and residue field  $k$  of characteristic  $p > 0$ . Further, let  $\pi : X \rightarrow \text{Spec } R$  be a morphism of schemes that is proper and smooth.*

*Suppose that  $R$  is of ramification degree  $e < p - 1$  and that  $k$  is perfect. Then, the cokernel of the specialization homomorphism  $\text{sp}_{\bar{K}} : \text{Pic}(X_{\bar{K}}) \rightarrow \text{Pic}(X_{\bar{k}})$  is torsion-free.*

**Corollary 3.7.** *Let  $p \neq 2$  be a prime number and  $X$  be a scheme proper and flat over  $\mathbb{Z}$ . Suppose that the special fiber  $X_p$  is nonsingular.*

*Then, the cokernel of the specialization homomorphism*

$$\text{sp}_{\bar{\mathbb{Q}}} : \text{Pic}(X_{\bar{\mathbb{Q}}}) \rightarrow \text{Pic}(X_{\bar{\mathbb{F}}_p})$$

*is torsion-free.*

**Remark 3.8.** The technical condition on the ramification degree cannot be omitted. In fact, D. Maulik and B. Poonen [2010, Example 3.12] constructed counterexamples to the assertion of Theorem 3.6 in the situation that  $e \geq p - 1$ .

**Remarks 3.9** (elementary reductions). (i) Let  $R'$  be a discrete valuation ring, finite and flat over  $R$ . Then, the assertion for  $\text{pr}_2 : X \times_{\text{Spec } R} \text{Spec } R' \rightarrow \text{Spec } R'$ , obtained by base-change, implies that for  $\pi$ .

(ii) In particular, we may suppose that  $\pi : X \rightarrow \text{Spec } R$  has a section.

(iii) We may suppose that the fibers of  $\pi$  are geometrically connected.

Indeed, as  $\pi : X \rightarrow \text{Spec } R$  is proper and smooth, one has  $\pi_* \mathbb{O}_X = \tilde{S}$  for  $S$  a finite étale  $R$ -algebra [Grothendieck 1963, Remarque (7.8.10.i), p. 75]. Hence, there exists a discrete valuation ring  $R'$ , étale over  $R$ , such that  $S \otimes_R R'$  is a direct product of finitely many copies of  $R'$ . This means that the connected components of  $X \times_{\text{Spec } R} \text{Spec } R'$  have geometrically connected fibers. Knowing the assertion for each component separately, the proof will be complete.



**Proposition 3.10.** *Let  $R$  be a discrete valuation ring of characteristic 0 and let  $\pi : X \rightarrow \text{Spec } R$  be a proper and smooth morphism of schemes. Suppose that  $\pi$  has a section and that the fibers of  $\pi$  are geometrically connected.*

*Then, the specialization homomorphisms*

$$\text{sp}_{\bar{K}} : \text{Pic}(X_{\bar{K}}) \rightarrow \text{Pic}(X_{\bar{k}}) \quad \text{and} \quad \text{sp}_{\widehat{K}} : \text{Pic}(X_{\widehat{K}}) \rightarrow \text{Pic}(X_{\bar{k}})$$

*have the same image.*

*Proof.* As  $\text{sp}_{\bar{K}}$  factors via  $\text{sp}_{\widehat{K}}$ , we clearly have  $\text{im } \text{sp}_{\bar{K}} \subseteq \text{im } \text{sp}_{\widehat{K}}$ . We will show the reverse inclusion in several steps. Let an invertible sheaf  $\mathcal{L} \in \text{Pic}(X_{\widehat{K}})$  be given. We have to construct an invertible sheaf  $\mathcal{L}' \in \text{Pic}(X_{\bar{K}})$  having the same specialization as  $\mathcal{L}$ .

*First step* (the Picard scheme). Our assumptions on  $\pi$  imply that it is cohomologically flat in dimension zero [Grothendieck 1963, Proposition (7.8.6), p. 74]. Hence, by [Artin 1969b, Theorem 7.3], the Picard functor  $\text{Pic}_{X/R}$  is representable by an algebraic space  $P := \mathbf{Pic}_{X/R}$  that is locally of finite type over  $R$ . According to [Grothendieck 1962, Exp. 236, Théorème 2.1.i],  $P$  is separated. This is enough to ensure that  $P$  is actually a scheme [Raynaud 1970, Théorème (3.3.1)]. Further, every closed subset  $Z \subseteq P$ , being of finite type, is proper over  $R$ .

*Second step* (the representing morphism). The invertible sheaf  $\mathcal{L} \in \text{Pic}(X_{\widehat{K}})$  is defined over a finite extension  $L$  of  $\widehat{K}$ . Hence, it defines a morphism  $i : \text{Spec } L \rightarrow P$ . As  $\widehat{K}$  is complete, there is a unique prolongation to  $L$  of the discrete valuation on  $\widehat{K}$ . That is, we have a discrete valuation ring  $S \supseteq \widehat{R}$ . There is a unique continuation  $j : \text{Spec } S \rightarrow P$  of  $i$ .

*Third step* (Artin approximation). By Lemma 3.12, we have  $S = \widehat{S}$  for a discrete valuation ring  $\underline{S}$ , finite over  $R$ . Write  $\underline{L}$  for the quotient field of  $\underline{S}$ . This is a finite extension of  $K$ .

We now recall that discrete valuation rings of characteristic zero are excellent [Grothendieck 1965, Scholie (7.8.3.iii), p. 214]. In particular, Artin's approximation results [1969a] are applicable. According to [1969a, Corollary (2.5)], there are an étale extension  $S'$  of  $\underline{S}$  and a morphism  $j' : \text{Spec } S' \rightarrow P$  of schemes that coincides, up to extensions of the base field, with  $j$  on the special fiber.

Corresponding to  $j'$ , there is some  $\xi \in \text{Pic}_{X/R}(\text{Spec } S')$ .

*Fourth step* (an invertible sheaf). As the fibers of  $X$  are geometrically connected, we have  $\pi_* \mathcal{O}_X = \mathcal{O}_{\text{Spec } R}$ . Further, since  $\pi$  has a section, one has [Grothendieck 1962, Exp. 232, Proposition 2.1]

$$\text{Pic}_{X/R}(T) = \text{Pic}(X \times_{\text{Spec } R} T) / \text{Pic}(T)$$

for every  $R$ -scheme  $T$ . In particular,

$$\begin{aligned} \text{Pic}_{X/R}(\text{Spec } S') &= \text{Pic}(X \times_{\text{Spec } R} \text{Spec } S') / \text{Pic}(\text{Spec } S') \\ &= \text{Pic}(X \times_{\text{Spec } R} \text{Spec } S'). \end{aligned}$$

Hence,  $\xi$  defines an invertible sheaf on  $X \times_{\text{Spec } R} \text{Spec } S'$ . Let  $\mathcal{L}' \in \text{Pic}(X_{\underline{L}})$  be its restriction to the generic fiber. Then, by construction,  $\mathcal{L}'$  has the same specialization as  $\mathcal{L}$ . The assertion follows.  $\square$

**Remark 3.11.** Suppose that  $H^1(X, \mathcal{O}_X) = 0$ . Then, [Proposition 3.10](#) is significantly more elementary. In fact, the Picard scheme  $P_K$  is of dimension zero [[Grothendieck 1962](#), Exp. 236, Proposition 2.10.iii] in this case. Hence, every point on  $P_K$  is defined over  $\bar{K}$ . No approximation argument is necessary.

Actually, the assumption  $H^1(X, \mathcal{O}_X) = 0$  is fulfilled in the examples, discussed in [1.7](#) and below in [Section 4](#).

**Lemma 3.12.** *Let  $R$  be a discrete valuation ring with quotient field  $K$  of characteristic zero and  $L/\widehat{K}$  a finite field extension of its completion.*

*Then, there exists a subfield  $\underline{L} \subset L$ , finite over  $K$ , such that  $\widehat{\underline{L}} = L$ .*

*Proof.* Choose a primitive element  $x$  of  $L$  over  $\widehat{K}$  and let  $f \in \widehat{K}[X]$  be its minimal polynomial. Then, the assertion is an immediate consequence of [[Serre 1968](#), Chapitre II, §2, Exercice 2].  $\square$

**3.13. Proof of [Theorem 3.6](#).** Consider the completion  $\widehat{R}$  of  $R$  and denote by  $\widehat{K}$  the corresponding quotient field. The ramification degree of  $\widehat{R}$  is the same as that of  $R$ . Therefore, [Proposition 3.4](#) shows that the specialization homomorphism  $\text{sp}_{\widehat{K}} : \text{Pic}(X_{\widehat{K}}) \rightarrow \text{Pic}(X_{\bar{k}})$  has a torsion-free cokernel. Further, by [Proposition 3.10](#),  $\text{sp}_{\widehat{K}}$  has the same image in  $\text{Pic}(X_{\bar{k}})$  as  $\text{sp}_{\bar{K}}$ . This implies the assertion.  $\square$

#### 4. The obstruction to first order deformations

The obstructions to lifting invertible sheaves were essential for the elementary proof of [Proposition 2.2](#), as discussed in [2.3](#). In some cases, they can be made explicit.

**Proposition 4.1.** *Let  $S$  be a K3 surface of degree 2 over  $\mathbb{Q}$ , given explicitly by*

$$w^2 = f_6(x, y, z)$$

*for  $f_6 \in \mathbb{Z}[x, y, z]$  of degree 6. Suppose, for a prime  $p \neq 2$  of good reduction, there is an  $\mathbb{F}_p$ -rational line “ $\ell = 0$ ”, tritangent to the ramification locus of  $S_p$ . Write  $l$  for an irreducible component of the pull-back of the tritangent.*

*One has  $f_6 \equiv f_3^2 + \ell f_5 \pmod{p}$  for homogeneous forms  $f_3, f_5 \in \mathbb{Z}[x, y, z]$ . Put*

$$G(x, y, z) := (f_6 - f_3^2 - \ell f_5) / p.$$

Then,  $\mathcal{O}(l)$  lifts to  $S_{p^2}$  if and only if  $G$  vanishes in  $\mathbb{F}_p[x, y, z]/(\ell, f_3, f_5)$ .

*Proof.* Suppose that  $\mathcal{O}(l)$  has a lift  $\mathcal{L} \in \text{Pic}(X_{p^2})$ . Then,  $\mathcal{L}/p\mathcal{L} \cong \mathcal{O}(l)$ . Since multiplication by  $p$  induces an isomorphism  $\mathcal{L}/p\mathcal{L} \cong p\mathcal{L}$ , we automatically have a short exact sequence

$$0 \rightarrow \mathcal{O}(l) \rightarrow \mathcal{L} \rightarrow \mathcal{O}(l) \rightarrow 0.$$

As  $H^1(X_p, \mathcal{O}(l)) = 0$ , the restriction map  $H^0(X_{p^2}, \mathcal{L}) \rightarrow H^0(X_p, \mathcal{O}(l))$  is a surjection. That is, the divisor  $l$  on  $X_p$  necessarily lifts to an effective Cartier divisor on  $X_{p^2}$ .

This is possible only when the line defined by  $\ell$  may be lifted to  $P_{p^2}^2$  in such a way that it is still a tritangent. On the other hand, if  $\ell$  may be lifted to  $P_{p^2}^2$  such that it is still a tritangent, then clearly  $\mathcal{O}(l)$  lifts to  $X_{p^2}$ .

Explicitly, the condition means that  $f_6$  is a square modulo  $p^2$  and some lift of  $\ell$ . Writing

$$f_6 \equiv (f_3 + pf'_3)^2 + (\ell + p\ell')(f_5 + pf'_5) \pmod{p^2},$$

one immediately sees that this is equivalent to the assertion that  $G$  vanishes in  $\mathbb{F}_p[x, y, z]/(\ell, f_3, f_5)$ . □

**Remark 4.2.** There is another proof that consists of the determination of the cohomological obstruction to lifting  $\mathcal{O}(l)$ , that is, of the image of  $\mathcal{O}(l)$  under the connecting homomorphism  $d : \text{Pic}(X_p) \rightarrow H^2(X_p, \mathcal{O}_{X_p})$  that is induced by the short exact sequence

$$0 \rightarrow \mathcal{O}_{X_p} \rightarrow \mathcal{O}_{X_{p^2}}^* \rightarrow \mathcal{O}_{X_p}^* \rightarrow 0.$$

The obstruction may easily be computed in Čech cohomology for a suitable affine open covering of  $X_{p^2}$ . Via the corresponding isomorphism  $H^2(X_p, \mathcal{O}_{X_p}) \cong \mathbb{F}_p$ , our result is indeed  $((-G) \pmod{(p, \ell, f_3, f_5)})$ . The necessary calculations are, however, rather lengthy and shall not be reproduced here.

**4.3.** In the examples below, we will use the obstruction in its explicit form, as given in [Proposition 4.1](#). The methods for point counting, which we apply, are explained in some detail in [[Elsenhans and Jahnel 2008a](#); [2008b](#); [2010](#)].

**Example 4.4.** Let  $S$  be a  $K3$  surface over  $\mathbb{Q}$  given by  $w^2 = f_6(x, y, z)$ . Suppose

$$\begin{aligned} f_6(x, y, z) \equiv & x^6 + 2x^5z + 2x^4y^2 + 2x^4z^2 + 2x^3y^3 + 2x^3z^3 \\ & + 2x^2y^4 + 2x^2y^3z + x^2z^4 + xy^3z^2 + 2xz^5 + y^6 \pmod{3}. \end{aligned}$$

Assume further that the coefficient of  $y^2z^4$  is not divisible by 9.

Then,  $\text{rk Pic}(S_{\mathbb{Q}}) = 1$ .

*Proof.* A direct calculation shows that, modulo 3, the right hand side is  $f_3^2 + xf_5$  for  $f_3 = 2x^3 + 2x^2z + xz^2 + 2y^3$  and  $f_5 = 2x^3y^2 + x^2z^3 + 2xy^4 + 2z^5$ . Thus, the branch locus of  $S_3$  has a tritangent line given by  $x = 0$ .

The numbers of points over  $\mathbb{F}_{3^d}$  are, in this order, 19, 127, 676, 6 751, 58 564, 532 414, 4 791 232, 43 038 703, 387 383 311, and 3 486 675 052. For the decomposition of the characteristic polynomial of the Frobenius on  $H_{\text{ét}}^2(S_{\mathbb{F}_3}, \mathbb{Q}_l(1))$ , we find

$$\frac{1}{3}(t-1)^2(3t^{20} - 3t^{19} - 3t^{18} + 8t^{17} - 3t^{16} - 4t^{15} + 6t^{14} - 4t^{13} + 2t^{12} + 4t^{11} - 7t^{10} + 4t^9 + 2t^8 - 4t^7 + 6t^6 - 4t^5 - 3t^4 + 8t^3 - 3t^2 - 3t + 3).$$

This shows  $\text{rk Pic}(S_{\mathbb{F}_3}) \leq 2$ .

Let  $l$  be an irreducible component of the pull-back of the tritangent line. We have to show that the obstruction to lifting  $\mathbb{O}(l)$  is nonzero. For this, we observe that  $x$ ,  $f_3$ , and  $f_5$  do not generate the monomial  $y^2z^4$ . However,  $G$  contains this monomial by its very definition.  $\square$

**Example 4.5.** Consider the  $K3$  surface  $S$  over  $\mathbb{Q}$ , given by  $w^2 = f_6(x, y, z)$  for

$$\begin{aligned} f_6(x, y, z) = & 4x^6 + 2x^5y + 12x^5z + 2x^4y^2 + 4x^4yz + 12x^4z^2 + 24x^3y^3 - 57x^3y^2z \\ & - 9x^3yz^2 + 6x^3z^3 + 8x^2y^4 - 5x^2y^3z - 72x^2y^2z^2 + 7x^2yz^3 \\ & + 4x^2z^4 + 20xy^4z - 52xy^3z^2 - 57xy^2z^3 + 7xyz^4 + 4y^5z \\ & - 7y^4z^2 - 18y^3z^3 + 7y^2z^4 + 12yz^5 + 2z^6. \end{aligned}$$

Then,  $\text{rk Pic}(S_{\mathbb{Q}}) = 3$ .

*Proof.* We have

$$\begin{aligned} f_6 = & (2x^3 + 2x^2z + 2y^2z + yz^2 + z^3)^2 \\ & + (2x^2 + 2xz + yz + z^2)(x^3y + 2x^3z + x^2y^2 + x^2yz + 2x^2z^2 + 12xy^3 \\ & - 34xy^2z - 9xyz^2 - 2xz^3 + 4y^4 - 15y^3z - 7y^2z^2 + 9yz^3 + z^4) \end{aligned}$$

and

$$\begin{aligned} f_6 = & 4(x^3 + 2x^2y + 2x^2z + xy^2 + xyz + xz^2 + y^2z + yz^2 + z^3)^2 \\ & - (x^2 + xz + yz + z^2)(14x^3y + 4x^3z + 22x^2y^2 + 22x^2yz + 8x^2z^2 - 8xy^3 \\ & + 61xy^2z + 9xyz^2 + 6xz^3 - 4y^4 + 15y^3z + 11y^2z^2 - 6yz^3 + 2z^4). \end{aligned}$$

Hence, there are two conics  $C_1$  and  $C_2$ , each of which is six times tangent to the ramification locus of  $S$ . The irreducible components of their pull-backs yield the

intersection matrix

$$\begin{pmatrix} -2 & 6 & 1 & 3 \\ 6 & -2 & 3 & 1 \\ 1 & 3 & -2 & 6 \\ 3 & 1 & 6 & -2 \end{pmatrix},$$

which is of rank 3. Hence,  $\text{rk Pic}(S_{\overline{\mathbb{Q}}}) \geq 3$ .

On the other hand,  $S$  has good reduction at the prime  $p = 3$ . Point counting over extensions of  $\mathbb{F}_3$  shows that the characteristic polynomial of the Frobenius operating on  $H_{\text{ét}}^2(S_{\overline{\mathbb{F}}_3}, \mathbb{Q}_l(1))$  is

$$\frac{1}{3}(t - 1)^4(3t^{18} + 3t^{17} + 2t^{16} + 2t^{15} + 4t^{14} + 5t^{13} + 4t^{12} + 3t^{11} + 6t^{10} + 8t^9 + 6t^8 + 3t^7 + 4t^6 + 5t^5 + 4t^4 + 2t^3 + 2t^2 + 3t + 3).$$

Consequently, we have  $\text{rk Pic}(S_{\overline{\mathbb{F}}_3}) \leq 4$ .

In particular, the assumption  $\text{rk Pic}(S_{\overline{\mathbb{Q}}}) > 3$  implies  $\text{rk Pic}(S_{\overline{\mathbb{Q}}}) = \text{rk Pic}(S_{\overline{\mathbb{F}}_3})$ . **Theorem 3.6** guarantees that the specialization map  $\text{sp}_{\overline{\mathbb{Q}}} : \text{Pic}(S_{\overline{\mathbb{Q}}}) \rightarrow \text{Pic}(S_{\overline{\mathbb{F}}_3})$  must be bijective. Giving one invertible sheaf  $\mathcal{L} \in \text{Pic}(S_{\overline{\mathbb{F}}_3})$  with a nontrivial obstruction will be enough to yield a contradiction.

For this, observe that the ramification locus of  $S_3$  has a tritangent line given by  $x + y + z = 0$ . Indeed,

$$f_6(x, y, z) \equiv (x^3 + x^2y + xy^2 + y^3)^2 + (x + y + z)(2x^3y^2 + x^3yz + 2x^2yz^2 + 2xy^4 + xy^3z + xy^2z^2 + 2xyz^3 + xz^4 + 2y^5 + 2y^4z + yz^4 + 2z^5) \pmod{3}.$$

Modulo the ideal  $(3, x + y + z)$ , we have

$$f_3 \equiv x^3 + x^2y + xy^2 + y^3, \\ f_5 \equiv -(x^5 + x^3y^2 + x^2y^3 + xy^4 + y^5), \quad G \equiv x^6 + 2x^5y + x^4y^2 + 2xy^5 + y^6.$$

Trying to generate  $G$  by  $3, x + y + z, f_3$ , and  $f_5$  now leads to a system of seven linear equations in six unknowns that is easily seen to be unsolvable.  $\square$

**Remarks 4.6.** (i) It is not at all hard to generate more examples similar to 1.7 and 4.4. Choosing the coefficients in  $\mathbb{F}_p$  at random, one usually finds Picard rank 2 over  $\overline{\mathbb{F}}_p$  after a few trials. One may work with small primes, only, say  $p \leq 7$ .

Clearly, for our arguments, it is of importance to have explicit generators for  $\text{Pic}(S_{\overline{\mathbb{F}}_p})$ . In practice, it turns out that a second generator may often be found. We have no formal reason for this. However, [Kovács 1994] might give an indication.

In Example 4.4, we applied a linear transform in order to make the obstruction depend only on a single coefficient. In general, one would have a linear form in the coefficients.

(ii) [Example 4.5](#) is a bit more particular. Both conics, which are six times tangent to the ramification sextic, simultaneously lift to  $\mathbb{Q}$ . This is not at all the generic behavior.

(iii) It seems to be substantially more difficult to construct examples for which  $\mathrm{rk}\mathrm{Pic}(X) \leq \mathrm{rk}\mathrm{Pic}(X_p) - 2$  may be shown. To understand the problem, recall the obstruction homomorphism  $\delta : \mathrm{Pic}(X_p) \rightarrow H^2(X, \mathbb{C}_X)$ , introduced in [Remark 2.3](#). In [Proposition 4.1](#), we calculated  $\delta(\mathcal{O}(l))$  at a precision of one  $p$ -adic digit.

In order to verify  $\mathrm{rk}\mathrm{Pic}(X) \leq \mathrm{rk}\mathrm{Pic}(X_p) - 2$ , one would have to ensure that  $\mathrm{rk}_{\mathbb{Z}}(\mathrm{im}\delta) \geq 2$ . This, however, is impossible as long as only  $p$ -adic approximations of finitely many values  $\delta(\mathcal{L})$  are known.

There are methods known to show

$$\mathrm{rk}\mathrm{Pic}(X) \leq \mathrm{rk}\mathrm{Pic}(X_{p_1}) - 2 \quad \text{and} \quad \mathrm{rk}\mathrm{Pic}(X) \leq \mathrm{rk}\mathrm{Pic}(X_{p_2}) - 2$$

when one works with two primes [\[Elsenhans and Jahnel 2011\]](#).

## References

- [Artin 1969a] M. Artin, “Algebraic approximation of structures over complete local rings”, *Inst. Hautes Études Sci. Publ. Math.* 36 (1969), 23–58. [MR 42 #3087](#) [Zbl 0181.48802](#)
- [Artin 1969b] M. Artin, “Algebraization of formal moduli, I”, pp. 21–71 in *Global Analysis: Papers in Honor of K. Kodaira*, Univ. Tokyo Press, Tokyo, 1969. [MR 41 #5369](#) [Zbl 0205.50402](#)
- [Artin et al. 1973] M. Artin, A. Grothendieck, and J. L. Verdier (editors), *Théorie des topos et cohomologie étale des schémas, Tome 3*, Lecture Notes in Mathematics **305**, Springer, Berlin, 1973. [MR 50 #7132](#) [Zbl an:0245.00002](#)
- [Barth et al. 1984] W. Barth, C. Peters, and A. Van de Ven, *Compact complex surfaces*, *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)* **4**, Springer, Berlin, 1984. [MR 86c:32026](#) [Zbl 0718.14023](#)
- [Berthelot et al. 1971] P. Berthelot, A. Grothendieck, and L. Illusie (editors), *Théorie des intersections et théorème de Riemann–Roch (SGA 6)*, Lecture Notes in Mathematics **225**, Springer, Berlin, 1971. [MR 50 #7133](#) [Zbl 0218.14001](#)
- [Elsenhans and Jahnel 2008a] A.-S. Elsenhans and J. Jahnel, “ $K3$  surfaces of Picard rank one and degree two”, pp. 212–225 in *Algorithmic number theory*, edited by A. J. van der Poorten and A. Stein, Lecture Notes in Comput. Sci. **5011**, Springer, Berlin, 2008. [MR 2010h:11102](#) [Zbl 1205.11073](#)
- [Elsenhans and Jahnel 2008b] A.-S. Elsenhans and J. Jahnel, “ $K3$  surfaces of Picard rank one which are double covers of the projective plane”, pp. 63–77 in *Higher-dimensional geometry over finite fields*, edited by D. Kaledin and Y. Tschinkel, NATO Sci. Peace Secur. Ser. D Inf. Commun. Secur. **16**, IOS, Amsterdam, 2008. [MR 2009j:14047](#) [Zbl 1182.14036](#)
- [Elsenhans and Jahnel 2010] A.-S. Elsenhans and J. Jahnel, “On Weil polynomials of  $K3$  surfaces”, pp. 126–141 in *Algorithmic number theory*, edited by G. Hanrot et al., Lecture Notes in Comput. Sci. **6197**, Springer, Berlin, 2010. [MR 2011m:11130](#) [Zbl 05793676](#)
- [Elsenhans and Jahnel 2011] A.-S. Elsenhans and J. Jahnel, “On the computation of the Picard group for  $K3$  surfaces”, *Math. Proc. Cambridge Philos. Soc.* **151**:2 (2011), 263–270. [MR 2823134](#) [Zbl 1223.14044](#)

- [Grothendieck 1961] A. Grothendieck, *Étude cohomologique des faisceaux cohérents, I (EGA III)*, Inst. Hautes Études Sci. Publ. Math. **11**, 1961. [MR 29 #1209](#) [Zbl 0118.36206](#)
- [Grothendieck 1962] A. Grothendieck, *Fondements de la géométrie algébrique: Extraits du Séminaire Bourbaki, 1957–1962*, Secrétariat mathématique, Paris, 1962. [MR 26 #3566](#) [Zbl 0239.14002](#)
- [Grothendieck 1963] A. Grothendieck, *Étude cohomologique des faisceaux cohérents, II (EGA III)*, Inst. Hautes Études Sci. Publ. Math. **17**, 1963. [MR 29 #1210](#) [Zbl 0122.16102](#)
- [Grothendieck 1965] A. Grothendieck, *Étude locale des schémas et des morphismes de schémas, II (EGA IV)*, Publications Mathématiques **24**, Institut des Hautes Études Scientifiques, Paris, 1965. [MR 33 #7330](#) [Zbl 0135.39701](#)
- [Grothendieck 1967] A. Grothendieck, *Étude locale des schémas et des morphismes de schémas, IV (EGA IV)*, Publications Mathématiques **32**, Institut des Hautes Études Scientifiques, Paris, 1967. [MR 39 #220](#) [Zbl 0153.22301](#)
- [Kovács 1994] S. J. Kovács, “The cone of curves of a  $K3$  surface”, *Math. Ann.* **300**:4 (1994), 681–691. [MR 96a:14044](#) [Zbl 0813.14026](#)
- [van Luijk 2007] R. van Luijk, “ $K3$  surfaces with Picard number one and infinitely many rational points”, *Algebra Number Theory* **1**:1 (2007), 1–15. [MR 2008d:14058](#) [Zbl 1123.14022](#)
- [Maulik and Poonen 2010] D. Maulik and B. Poonen, “Néron–Severi groups under specialization”, preprint, version 3, 2010. [arXiv 0907.4781v3](#)
- [Milne 1975] J. S. Milne, “On a conjecture of Artin and Tate”, *Ann. of Math. (2)* **102**:3 (1975), 517–533. [MR 54 #2659](#) [Zbl 0343.14005](#)
- [Raynaud 1970] M. Raynaud, “Spécialisation du foncteur de Picard”, *Inst. Hautes Études Sci. Publ. Math.* **38** (1970), 27–76. [MR 44 #227](#) [Zbl 0207.51602](#)
- [Raynaud 1979] M. Raynaud, “‘ $p$ -torsion’ du schéma de Picard”, pp. 87–148 in *Journées de Géométrie Algébrique de Rennes* (Rennes, 1978), vol. 2, Astérisque **64**, Soc. Math. France, Paris, 1979. [MR 81f:14026](#) [Zbl 0434.14024](#)
- [Serre 1968] J.-P. Serre, *Corps locaux*, 2nd ed., Publications de l’Université de Nancago **8**, Hermann, Paris, 1968. [MR 50 #7096](#)

Communicated by János Kollár

Received 2010-03-31

Revised 2011-03-01

Accepted 2011-04-01

[stephan.elsenhans@uni-bayreuth.de](mailto:stephan.elsenhans@uni-bayreuth.de)

Mathematisches Institut, Universität Bayreuth,  
Universitätsstraße 30, D-95440 Bayreuth, Germany  
<http://www.staff.uni-bayreuth.de/~btm216>

[jahnel@mathematik.uni-siegen.de](mailto:jahnel@mathematik.uni-siegen.de)

Fachbereich 6 Mathematik, Universität Siegen,  
Walter-Flex-Straße 3, D-57068 Siegen, Germany  
<http://www.uni-math.gwdg.de/jahnel>

# Algebra & Number Theory

[msp.berkeley.edu/ant](http://msp.berkeley.edu/ant)

## EDITORS

### MANAGING EDITOR

Bjorn Poonen  
Massachusetts Institute of Technology  
Cambridge, USA

### EDITORIAL BOARD CHAIR

David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Georgia Benkart	University of Wisconsin, Madison, USA	Shigefumi Mori	RIMS, Kyoto University, Japan
Dave Benson	University of Aberdeen, Scotland	Raman Parimala	Emory University, USA
Richard E. Borcherds	University of California, Berkeley, USA	Jonathan Pila	University of Oxford, UK
John H. Coates	University of Cambridge, UK	Victor Reiner	University of Minnesota, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Karl Rubin	University of California, Irvine, USA
Brian D. Conrad	University of Michigan, USA	Peter Sarnak	Princeton University, USA
Hélène Esnault	Universität Duisburg-Essen, Germany	Joseph H. Silverman	Brown University, USA
Hubert Flenner	Ruhr-Universität, Germany	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Ronald Solomon	Ohio State University, USA
Andrew Granville	Université de Montréal, Canada	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Joseph Gubeladze	San Francisco State University, USA	J. Toby Stafford	University of Michigan, USA
Ehud Hrushovski	Hebrew University, Israel	Bernd Sturmfels	University of California, Berkeley, USA
Craig Huneke	University of Kansas, USA	Richard Taylor	Harvard University, USA
Mikhail Kapranov	Yale University, USA	Ravi Vakil	Stanford University, USA
Yujiro Kawamata	University of Tokyo, Japan	Michel van den Bergh	Hasselt University, Belgium
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Yuri Manin	Northwestern University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Barry Mazur	Harvard University, USA	Andrei Zelevinsky	Northeastern University, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne	Efim Zelmanov	University of California, San Diego, USA
Susan Montgomery	University of Southern California, USA		

## PRODUCTION

[contact@msp.org](mailto:contact@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [www.jant.org](http://www.jant.org) for submission instructions.


The subscription price for 2011 is US \$150/year for the electronic version, and \$210/year (+\$35 shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840, USA.

Algebra & Number Theory (ISSN 1937-0652) at Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

ANT peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
<http://msp.org/>

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2011 by Mathematical Sciences Publishers



# Algebra & Number Theory

Volume 5    No. 8    2011

---

The behavior of Hecke $L$ -functions of real quadratic fields at $s = 0$ BYUNGHEUP JUN and JUNGYUN LEE	1001
The Picard group of a $K3$ surface and its reduction modulo $p$ ANDREAS-STEPHAN ELSENHANS and JÖRG JAHNEL	1027
Linear determinantal equations for all projective schemes JESSICA SIDMAN and GREGORY G. SMITH	1041
Involutions, weights and $p$ -local structure GEOFFREY R. ROBINSON	1063
Parametrizing quartic algebras over an arbitrary base MELANIE MATCHETT WOOD	1069
Coleman maps and the $p$ -adic regulator ANTONIO LEI, DAVID LOEFFLER and SARAH LIVIA ZERBES	1095
Conjecture de Shafarevitch effective pour les revêtements cycliques ROBIN DE JONG and GAËL RÉMOND	1133