

## ON BALLICO-HEFEZ CURVES AND ASSOCIATED SUPERSINGULAR SURFACES

THANH HOAI HOANG AND ICHIRO SHIMADA

### Abstract

Let  $p$  be a prime integer, and  $q$  a power of  $p$ . The Ballico-Hefez curve is a non-reflexive nodal rational plane curve of degree  $q + 1$  in characteristic  $p$ . We investigate its automorphism group and defining equation. We also prove that the surface obtained as the cyclic cover of the projective plane branched along the Ballico-Hefez curve is unirational, and hence is supersingular. As an application, we obtain a new projective model of the supersingular  $K3$  surface with Artin invariant 1 in characteristic 3 and 5.

### 1. Introduction

We work over an algebraically closed field  $k$  of positive characteristic  $p > 0$ . Let  $q = p^v$  be a power of  $p$ .

In positive characteristics, algebraic varieties often possess interesting properties that are not observed in characteristic zero. One of those properties is the failure of reflexivity. In [4], Ballico and Hefez classified irreducible plane curves  $X$  of degree  $q + 1$  such that the natural morphism from the conormal variety  $C(X)$  of  $X$  to the dual curve  $X^\vee$  has inseparable degree  $q$ . The Ballico-Hefez curve in the title of this note is one of the curves that appear in their classification. It is defined in Fukasawa, Homma and Kim [8] as follows.

**DEFINITION 1.1.** The *Ballico-Hefez curve* is the image of the morphism  $\phi : \mathbf{P}^1 \rightarrow \mathbf{P}^2$  defined by

$$[s : t] \mapsto [s^{q+1} : t^{q+1} : st^q + s^q t].$$

**THEOREM 1.2** (Ballico and Hefez [4], Fukasawa, Homma and Kim [8]).  
(1) *Let  $B$  be the Ballico-Hefez curve. Then  $B$  is a curve of degree  $q + 1$  with*

---

2000 *Mathematics Subject Classification.* primary 14H45, secondary 14J25, 14J28.

*Key words and phrases.* plane curve, positive characteristic, supersingularity,  $K3$  surface.

Partially supported by JSPS Grant-in-Aid for Challenging Exploratory Research No. 23654012 and JSPS Grants-in-Aid for Scientific Research (C) No. 25400042.

Received May 21, 2013; revised February 4, 2014.

$(q^2 - q)/2$  ordinary nodes, the dual curve  $B^\vee$  is of degree 2, and the natural morphism  $C(B) \rightarrow B^\vee$  has inseparable degree  $q$ .

(2) Let  $X \subset \mathbf{P}^2$  be an irreducible singular curve of degree  $q + 1$  such that the dual curve  $X^\vee$  is of degree  $> 1$  and the natural morphism  $C(X) \rightarrow X^\vee$  has inseparable degree  $q$ . Then  $X$  is projectively isomorphic to the Ballico-Hefez curve.

Recently, geometry and arithmetic of the Ballico-Hefez curve have been investigated by Fukasawa, Homma and Kim [8] and Fukasawa [7] from various points of view, including coding theory and Galois points. As is pointed out in [8], the Ballico-Hefez curve has many properties in common with the Hermitian curve; that is, the Fermat curve of degree  $q + 1$ , which also appears in the classification of Ballico and Hefez [4]. In fact, we can easily see that the image of the line

$$x_0 + x_1 + x_2 = 0$$

in  $\mathbf{P}^2$  by the morphism  $\mathbf{P}^2 \rightarrow \mathbf{P}^2$  given by

$$[x_0 : x_1 : x_2] \mapsto [x_0^{q+1} : x_1^{q+1} : x_2^{q+1}]$$

is projectively isomorphic to the Ballico-Hefez curve. Hence, up to linear transformation of coordinates, the Ballico-Hefez curve is defined by an equation

$$x_0^{1/(q+1)} + x_1^{1/(q+1)} + x_2^{1/(q+1)} = 0$$

in the style of ‘‘Coxeter curves’’ (see Griffith [9]).

In this note, we prove the the following:

**PROPOSITION 1.3.** *Let  $B$  be the Ballico-Hefez curve. Then the group*

$$\text{Aut}(B) := \{g \in \text{PGL}_3(k) \mid g(B) = B\}$$

*of projective automorphisms of  $B \subset \mathbf{P}^2$  is isomorphic to  $\text{PGL}_2(\mathbf{F}_q)$ .*

**PROPOSITION 1.4.** *The Ballico-Hefez curve is defined by the following equations:*

• *When  $p = 2$ ,*

$$x_0^q x_1 + x_0 x_1^q + x_2^{q+1} + \sum_{i=0}^{v-1} x_0^{2^i} x_1^{2^i} x_2^{q+1-2^{i+1}} = 0, \quad \text{where } q = 2^v.$$

• *When  $p$  is odd,*

$$2(x_0^q x_1 + x_0 x_1^q) - x_2^{q+1} - (x_2^2 - 4x_1 x_0)^{(q+1)/2} = 0.$$

*Remark 1.5.* In fact, the defining equation for  $p = 2$  has been obtained by Fukasawa in an apparently different form (see Remark 3 of [6]).

Another property of algebraic varieties peculiar to positive characteristics is the failure of Lüroth's theorem for surfaces; a non-rational surface can be unirational in positive characteristics. A famous example of this phenomenon is the Fermat surface of degree  $q + 1$ . Shioda [18] and Shioda-Katsura [19] showed that the Fermat surface  $F$  of degree  $q + 1$  is unirational (see also [16] for another proof). This surface  $F$  is obtained as the cyclic cover of  $\mathbf{P}^2$  with degree  $q + 1$  branched along the Fermat curve of degree  $q + 1$ , and hence, for any divisor  $d$  of  $q + 1$ , the cyclic cover of  $\mathbf{P}^2$  with degree  $d$  branched along the Fermat curve of degree  $q + 1$  is also unirational.

We prove an analogue of this result for the Ballico-Hefez curve. Let  $d$  be a divisor of  $q + 1$  larger than 1. Note that  $d$  is prime to  $p$ .

**PROPOSITION 1.4.** *Let  $\gamma : S_d \rightarrow \mathbf{P}^2$  be the cyclic covering of  $\mathbf{P}^2$  with degree  $d$  branched along the Ballico-Hefez curve. Then there exists a dominant rational map  $\mathbf{P}^2 \cdots \rightarrow S_d$  of degree  $2q$  with inseparable degree  $q$ .*

Note that  $S_d$  is not rational except for the case  $(d, q + 1) = (3, 3)$  or  $(2, 4)$ .

A smooth surface  $X$  is said to be *supersingular* (in the sense of Shioda) if the second  $l$ -adic cohomology group  $H^2(X)$  of  $X$  is generated by the classes of curves. Shioda [18] proved that every smooth unirational surface is supersingular. Hence we obtain the following:

**COROLLARY 1.7.** *Let  $\rho : \tilde{S}_d \rightarrow S_d$  be the minimal resolution of  $S_d$ . Then the surface  $\tilde{S}_d$  is supersingular.*

We present a finite set of curves on  $\tilde{S}_d$  whose classes span  $H^2(\tilde{S}_d)$ . For a point  $P$  of  $\mathbf{P}^1$ , let  $l_P \subset \mathbf{P}^2$  denote the line tangent at  $\phi(P) \in B$  to the branch of  $B$  corresponding to  $P$ . It was shown in [8] that, if  $P$  is an  $\mathbf{F}_{q^2}$ -rational point of  $\mathbf{P}^1$ , then  $l_P$  and  $B$  intersect only at  $\phi(P)$ , and hence the strict transform of  $l_P$  by the composite  $\tilde{S}_d \rightarrow S_d \rightarrow \mathbf{P}^2$  is a union of  $d$  rational curves  $l_P^{(0)}, \dots, l_P^{(d-1)}$ .

**PROPOSITION 1.8.** *The cohomology group  $H^2(\tilde{S}_d)$  is generated by the classes of the following rational curves on  $\tilde{S}_d$ ; the irreducible components of the exceptional divisor of the resolution  $\rho : \tilde{S}_d \rightarrow S_d$  and the rational curves  $l_P^{(i)}$ , where  $P$  runs through the set  $\mathbf{P}^1(\mathbf{F}_{q^2})$  of  $\mathbf{F}_{q^2}$ -rational points of  $\mathbf{P}^1$  and  $i = 0, \dots, d - 1$ .*

Note that, when  $(d, q + 1) = (4, 4)$  and  $(2, 6)$ , the surface  $\tilde{S}_d$  is a  $K3$  surface. In these cases, we can prove that the classes of rational curves given in Proposition 1.8 generate the Néron-Severi lattice  $\text{NS}(\tilde{S}_d)$  of  $\tilde{S}_d$ , and that the discriminant of  $\text{NS}(\tilde{S}_d)$  is  $-p^2$ . Using this fact and the result of Ogas [13, 14] and Rudakov-Shafarevich [15] on the uniqueness of a supersingular  $K3$  surface with Artin invariant 1, we prove the following:

**PROPOSITION 1.9.** (1) *If  $p = q = 3$ , then  $\tilde{S}_4$  is isomorphic to the Fermat quartic surface*

$$w^4 + x^4 + y^4 + z^4 = 0.$$

(2) *If  $p = q = 5$ , then  $\tilde{S}_2$  is isomorphic to the Fermat sextic double plane*

$$w^2 = x^6 + y^6 + z^6.$$

Recently, many studies on these supersingular  $K3$  surfaces with Artin invariant 1 in characteristics 3 and 5 have been carried out. See [10, 12] for characteristic 3 case, and [11, 17] for characteristic 5 case.

Thanks are due to Masaaki Homma and Satoru Fukasawa for their comments. We also thank the referee for his/her suggestion on the first version of this paper.

## 2. Basic properties of the Ballico-Hefez curve

We recall some properties of the Ballico-Hefez curve  $B$ . See Fukasawa, Homma and Kim [8] for the proofs.

It is easy to see that the morphism  $\phi : \mathbf{P}^1 \rightarrow \mathbf{P}^2$  is birational onto its image  $B$ , and that the degree of the plane curve  $B$  is  $q + 1$ . The singular locus  $\text{Sing}(B)$  of  $B$  consists of  $(q^2 - q)/2$  ordinary nodes, and we have

$$\phi^{-1}(\text{Sing}(B)) = \mathbf{P}^1(\mathbf{F}_{q^2}) \setminus \mathbf{P}^1(\mathbf{F}_q).$$

In particular, the singular locus  $\text{Sing}(S_d)$  of  $S_d$  consists of  $(q^2 - q)/2$  ordinary rational double points of type  $A_{d-1}$ . Therefore, by Artin [1, 2], the surface  $S_d$  is not rational if  $(d, q + 1) \neq (3, 3), (2, 4)$ .

Let  $t$  be the affine coordinate of  $\mathbf{P}^1$  obtained from  $[s : t]$  by putting  $s = 1$ , and let  $(x, y)$  be the affine coordinates of  $\mathbf{P}^2$  such that  $[x_0 : x_1 : x_2] = [1 : x : y]$ . Then the morphism  $\phi : \mathbf{P}^1 \rightarrow \mathbf{P}^2$  is given by

$$t \mapsto (t^{q+1}, t^q + t).$$

For a point  $P = [1 : t]$  of  $\mathbf{P}^1$ , the line  $l_P$  is defined by

$$x - t^q y + t^{2q} = 0.$$

Suppose that  $P \notin \mathbf{P}^1(\mathbf{F}_{q^2})$ . Then  $l_P$  intersects  $B$  at  $\phi(P) = (t^{q+1}, t^q + t)$  with multiplicity  $q$  and at the point  $(t^{q^2+q}, t^{q^2} + t^q) \neq \phi(P)$  with multiplicity 1. In particular, we have  $l_P \cap \text{Sing}(B) = \emptyset$ .

Suppose that  $P \in \mathbf{P}^1(\mathbf{F}_{q^2}) \setminus \mathbf{P}^1(\mathbf{F}_q)$ . Then  $l_P$  intersects  $B$  at the node  $\phi(P)$  of  $B$  with multiplicity  $q + 1$ . More precisely,  $l_P$  intersects the branch of  $B$  corresponding to  $P$  with multiplicity  $q$ , and the other branch transversely.

Suppose that  $P \in \mathbf{P}^1(\mathbf{F}_q)$ . Then  $\phi(P)$  is a smooth point of  $B$ , and  $l_P$  intersects  $B$  at  $\phi(P)$  with multiplicity  $q + 1$ . In particular, we have  $l_P \cap \text{Sing}(B) = \emptyset$ .

Combining these facts, we see that  $\phi(\mathbf{P}^1(\mathbf{F}_{q^2}))$  coincides with the set of smooth inflection points of  $B$ . (See [8] for the definition of inflection points.)

### 3. Proof of Proposition 1.3

We denote by  $\phi_B : \mathbf{P}^1 \rightarrow B$  the birational morphism  $t \mapsto (t^{q+1}, t^q + t)$  from  $\mathbf{P}^1$  to  $B$ . We identify  $\text{Aut}(\mathbf{P}^1)$  with  $\text{PGL}_2(k)$  by letting  $\text{PGL}_2(k)$  act on  $\mathbf{P}^1$  by

$$[s : t] \mapsto [as + bt : cs + dt] \quad \text{for} \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \text{PGL}_2(k).$$

Then  $\text{PGL}_2(\mathbf{F}_q)$  is the subgroup of  $\text{PGL}_2(k)$  consisting of elements that leave the set  $\mathbf{P}^1(\mathbf{F}_q)$  invariant. Since  $\phi_B$  is birational, the projective automorphism group  $\text{Aut}(B)$  of  $B$  acts on  $\mathbf{P}^1$  via  $\phi_B$ . The subset  $\phi_B(\mathbf{P}^1(\mathbf{F}_q))$  of  $B$  is projectively characterized as the set of smooth inflection points of  $B$ , and we have  $\mathbf{P}^1(\mathbf{F}_q) = \phi_B^{-1}(\phi_B(\mathbf{P}^1(\mathbf{F}_q)))$ . Hence  $\text{Aut}(B)$  is contained in the subgroup  $\text{PGL}_2(\mathbf{F}_q)$  of  $\text{PGL}_2(k)$ . Thus, in order to prove Proposition 1.3, it is enough to show that every element

$$g := \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \text{with} \quad a, b, c, d \in \mathbf{F}_q$$

of  $\text{PGL}_2(\mathbf{F}_q)$  is coming from the action of an element of  $\text{Aut}(B)$ . We put

$$\tilde{g} := \begin{bmatrix} a^2 & b^2 & ab \\ c^2 & d^2 & cd \\ 2ac & 2bd & ad + bc \end{bmatrix},$$

and let the matrix  $\tilde{g}$  act on  $\mathbf{P}^2$  by the left multiplication on the column vector  ${}^t[x_0 : x_1 : x_2]$ . Then we have

$$\phi \circ g = \tilde{g} \circ \phi,$$

because we have  $\lambda^q = \lambda$  for  $\lambda = a, b, c, d \in \mathbf{F}_q$ . Therefore  $g \mapsto \tilde{g}$  gives an isomorphism from  $\text{PGL}_2(\mathbf{F}_q)$  to  $\text{Aut}(B)$ .

### 4. Proof of Proposition 1.4

We put

$$F(x, y) := \begin{cases} x + x^q + y^{q+1} + \sum_{i=0}^{v-1} x^{2^i} y^{q+1-2^{i+1}} & \text{if } p = 2 \text{ and } q = 2^v, \\ 2x + 2x^q - y^{q+1} - (y^2 - 4x)^{(q+1)/2} & \text{if } p \text{ is odd,} \end{cases}$$

that is,  $F$  is obtained from the homogeneous polynomial in Proposition 1.4 by putting  $x_0 = 1$ ,  $x_1 = x$ ,  $x_2 = y$ . Since the polynomial  $F$  is of degree  $q+1$  and the plane curve  $B$  is also of degree  $q+1$ , it is enough to show that  $F(t^{q+1}, t^q + t) = 0$ .

Suppose that  $p = 2$  and  $q = 2^v$ . We put

$$S(x, y) := \sum_{i=0}^{v-1} \left( \frac{x}{y^2} \right)^{2^i}.$$

Then  $S(x, y)$  is a root of the Artin-Schreier equation

$$s^2 + s = \left(\frac{x}{y^2}\right)^q + \frac{x}{y^2}.$$

Hence  $S_1 := S(t^{q+1}, t^q + t)$  is a root of the equation  $s^2 + s = b$ , where

$$b := \left[ \frac{t^{q+1}}{(t^q + t)^2} \right]^q + \frac{t^{q+1}}{(t^q + t)^2} = \frac{t^{2q^2+q+1} + t^{q^2+3q} + t^{q^2+q+2} + t^{3q+1}}{(t^q + t)^{2q+2}}.$$

We put

$$S'(x, y) := \frac{x + x^q + y^{q+1}}{y^{q+1}}.$$

We can verify that  $S_2 := S'(t^{q+1}, t^q + t)$  is also a root of the equation  $s^2 + s = b$ . Hence we have either  $S_1 = S_2$  or  $S_1 = S_2 + 1$ . We can easily see that both of the rational functions  $S_1$  and  $S_2$  on  $\mathbf{P}^1$  have zero at  $t = \infty$ . Hence  $S_1 = S_2$  holds, from which we obtain  $F(t^{q+1}, t^q + t) = 0$ .

Suppose that  $p$  is odd. We put

$$S(x, y) := 2x + 2x^q - y^{q+1}, \quad S_1 := S(t^{q+1}, t^q + t), \quad \text{and}$$

$$S'(x, y) := (y^2 - 4x)^{(q+1)/2}, \quad S_2 := S'(t^{q+1}, t^q + t).$$

Then it is easy to verify that both of  $S_1^2$  and  $S_2^2$  are equal to

$$t^{2q^2+2q} - 2t^{2q^2+q+1} + t^{2q^2+2} - 2t^{q^2+3q} + 4t^{q^2+2q+1} - 2t^{q^2+q+2} + t^{4q} - 2t^{3q+1} + t^{2q+2}.$$

Therefore either  $S_1 = S_2$  or  $S_1 = -S_2$  holds. Comparing the coefficients of the top-degree terms of the polynomials  $S_1$  and  $S_2$  of  $t$ , we see that  $S_1 = S_2$ , whence  $F(t^{q+1}, t^q + t) = 0$  follows.

## 5. Proof of Propositions 1.6 and 1.8

We consider the universal family

$$L := \{(P, Q) \in \mathbf{P}^1 \times \mathbf{P}^2 \mid Q \in l_P\}$$

of the lines  $l_P$ , which is defined by

$$x - t^q y + t^{2q} = 0$$

in  $\mathbf{P}^1 \times \mathbf{P}^2$ , and let

$$\pi_1 : L \rightarrow \mathbf{P}^1, \quad \pi_2 : L \rightarrow \mathbf{P}^2$$

be the projections. We see that  $\pi_1 : L \rightarrow \mathbf{P}^1$  has two sections

$$\sigma_1 : t \mapsto (t, x, y) = (t, t^{q+1}, t^q + t),$$

$$\sigma_q : t \mapsto (t, x, y) = (t, t^{q^2+q}, t^{q^2} + t^q).$$

For  $P \in \mathbf{P}^1$ , we have  $\pi_2(\sigma_1(P)) = \phi(P)$  and  $l_P \cap B = \{\pi_2(\sigma_1(P)), \pi_2(\sigma_q(P))\}$ . Let  $\Sigma_1 \subset L$  and  $\Sigma_q \subset L$  denote the images of  $\sigma_1$  and  $\sigma_q$ , respectively. Then  $\Sigma_1$  and  $\Sigma_q$  are smooth curves, and they intersect transversely. Moreover, their intersection points are contained in  $\pi_1^{-1}(\mathbf{P}^1(\mathbf{F}_{q^2}))$ .

We denote by  $\bar{M}$  the fiber product of  $\gamma: S_d \rightarrow \mathbf{P}^2$  and  $\pi_2: L \rightarrow \mathbf{P}^2$  over  $\mathbf{P}^2$ . The pull-back  $\pi_2^*B$  of  $B$  by  $\pi_2$  is equal to the divisor  $q\Sigma_1 + \Sigma_q$ . Hence  $\bar{M}$  is defined by

$$(5.1) \quad \begin{cases} z^d = (y - t^q - t)^q (y - t^{q^2} - t^q), \\ x - t^q y + t^{2q} = 0. \end{cases}$$

We denote by  $M \rightarrow \bar{M}$  the normalization, and by

$$\alpha: M \rightarrow L, \quad \eta: M \rightarrow S_d$$

the natural projections. Since  $d$  is prime to  $q$ , the cyclic covering  $\alpha: M \rightarrow L$  of degree  $d$  branches exactly along the curve  $\Sigma_1 \cup \Sigma_q$ . Moreover, the singular locus  $\text{Sing}(M)$  of  $M$  is located over  $\Sigma_1 \cap \Sigma_q$ , and hence is contained in  $\alpha^{-1}(\pi_1^{-1}(\mathbf{P}^1(\mathbf{F}_{q^2})))$ .

Since  $\eta$  is dominant and  $\rho: \tilde{S}_d \rightarrow S_d$  is birational,  $\eta$  induces a rational map

$$\eta': M \cdots \rightarrow \tilde{S}_d.$$

Let  $A$  denote the affine open curve  $\mathbf{P}^1 \setminus \mathbf{P}^1(\mathbf{F}_{q^2})$ . We put

$$L_A := \pi_1^{-1}(A), \quad M_A := \alpha^{-1}(L_A).$$

Note that  $M_A$  is smooth. Let  $\pi_{1,A}: L_A \rightarrow A$  and  $\alpha_A: M_A \rightarrow L_A$  be the restrictions of  $\pi_1$  and  $\alpha$ , respectively. If  $P \in A$ , then  $l_P$  is disjoint from  $\text{Sing}(B)$ , and hence  $\eta(\alpha^{-1}(\pi_1^{-1}(P))) = \gamma^{-1}(l_P)$  is disjoint from  $\text{Sing}(S_d)$ . Therefore the restriction of  $\eta'$  to  $M_A$  is a morphism. It follows that we have a proper birational morphism

$$\beta: \tilde{M} \rightarrow M$$

from a smooth surface  $\tilde{M}$  to  $M$  such that  $\beta$  induces an isomorphism from  $\beta^{-1}(M_A)$  to  $M_A$  and that the rational map  $\eta'$  extends to a morphism  $\tilde{\eta}: \tilde{M} \rightarrow \tilde{S}_d$ . Summing up, we obtain the following commutative diagram:

$$(5.2) \quad \begin{array}{ccccccc} M_A & \hookrightarrow & \tilde{M} & \xrightarrow{\tilde{\eta}} & \tilde{S}_d & & \\ & & \square & \downarrow \beta & \downarrow \rho & & \\ M_A & \hookrightarrow & M & \xrightarrow{\eta} & S_d & & \\ \alpha_A \downarrow & & \square & \downarrow \alpha & \downarrow \gamma & & \\ L_A & \hookrightarrow & L & \xrightarrow{\pi_2} & \mathbf{P}^2 & & \\ \pi_{1,A} \downarrow & & \square & \downarrow \pi_1 & & & \\ A & \hookrightarrow & \mathbf{P}^1 & & & & \end{array}$$

Since the defining equation  $x - t^q y + t^{2q} = 0$  of  $L$  in  $\mathbf{P}^1 \times \mathbf{P}^2$  is a polynomial in  $k[x, y][t^q]$ , and its discriminant as a quadratic equation of  $t^q$  is  $y^2 - 4x \neq 0$ , the projection  $\pi_2$  is a finite morphism of degree  $2q$  and its inseparable degree is  $q$ . Hence  $\eta$  is also a finite morphism of degree  $2q$  and its inseparable degree is  $q$ . Therefore, in order to prove Proposition 1.6, it is enough to show that  $M$  is rational. We denote by  $k(M) = k(\bar{M})$  the function field of  $M$ . Since  $x = t^q y - t^{2q}$  on  $\bar{M}$ , the field  $k(M)$  is generated over  $k$  by  $y, z$  and  $t$ . Let  $c$  denote the integer  $(q+1)/d$ , and put

$$\tilde{z} := \frac{z}{(y - t^q - t)^c} \in k(M).$$

Then, from the defining equation (5.1) of  $\bar{M}$ , we have

$$\tilde{z}^d = \frac{y - t^{q^2} - t^q}{y - t^q - t}.$$

Therefore we have

$$y = \frac{\tilde{z}^d(t^q + t) - (t^{q^2} + t^q)}{\tilde{z}^d - 1},$$

and hence  $k(M)$  is equal to the purely transcendental extension  $k(\tilde{z}, t)$  of  $k$ . Thus Proposition 1.6 is proved.

We put

$$\Xi := \tilde{M} \setminus M_A = \beta^{-1}(\alpha^{-1}(\pi_1^{-1}(\mathbf{P}^1(\mathbf{F}_{q^2}))))).$$

Since the cyclic covering  $\alpha : M \rightarrow L$  branches along the curve  $\Sigma_1 = \sigma_1(\mathbf{P}^1)$ , the section  $\sigma_1 : \mathbf{P}^1 \rightarrow L$  of  $\pi_1$  lifts to a section  $\tilde{\sigma}_1 : \mathbf{P}^1 \rightarrow M$  of  $\pi_1 \circ \alpha$ . Let  $\tilde{\Sigma}_1$  denote the strict transform of the image of  $\tilde{\sigma}_1$  by  $\beta : \tilde{M} \rightarrow M$ .

LEMMA 5.1. *The Picard group  $\text{Pic}(\tilde{M})$  of  $\tilde{M}$  is generated by the classes of  $\tilde{\Sigma}_1$  and the irreducible components of  $\Xi$ .*

*Proof.* Since  $\Sigma_1 \cap \Sigma_q \cap L_A = \emptyset$ , the morphism

$$\pi_{1,A} \circ \alpha_A : M_A \rightarrow A$$

is a smooth  $\mathbf{P}^1$ -bundle. Let  $D$  be an irreducible curve on  $\tilde{M}$ , and let  $e$  be the degree of

$$\pi_1 \circ \alpha \circ \beta|_D : D \rightarrow \mathbf{P}^1.$$

Then the divisor  $D - e\tilde{\Sigma}_1$  on  $\tilde{M}$  is of degree 0 on the general fiber of the smooth  $\mathbf{P}^1$ -bundle  $\pi_{1,A} \circ \alpha_A$ . Therefore  $(D - e\tilde{\Sigma}_1)|_{M_A}$  is linearly equivalent in  $M_A$  to a multiple of a fiber of  $\pi_{1,A} \circ \alpha_A$ . Hence  $D$  is linearly equivalent to a linear combination of  $\tilde{\Sigma}_1$  and irreducible curves in the boundary  $\Xi = \tilde{M} \setminus M_A$ .  $\square$

The rational curves on  $\tilde{\mathcal{S}}_d$  listed in Proposition 1.8 are exactly equal to the irreducible components of

$$\rho^{-1} \left( \gamma^{-1} \left( \bigcup_{P \in \mathbf{P}^1(\mathbf{F}_{q^2})} l_P \right) \right).$$

Let  $V \subset H^2(\tilde{\mathcal{S}}_d)$  denote the linear subspace spanned by the classes of these rational curves. We will show that  $V = H^2(\tilde{\mathcal{S}}_d)$ .

Let  $h \in H^2(\tilde{\mathcal{S}}_d)$  denote the class of the pull-back of a line of  $\mathbf{P}^2$  by the morphism  $\gamma \circ \rho : \tilde{\mathcal{S}}_d \rightarrow \mathbf{P}^2$ . Suppose that  $P \in \mathbf{P}^1(\mathbf{F}_q)$ . Then  $l_P$  is disjoint from  $\text{Sing}(B)$ . Therefore we have

$$h = [(\gamma \circ \rho)^*(l_P)] = [l_P^{(0)}] + \cdots + [l_P^{(d-1)}] \in V.$$

Let  $\tilde{B}$  denote the strict transform of  $B$  by  $\gamma \circ \rho$ . Then  $\tilde{B}$  is written as  $d \cdot R$ , where  $R$  is a reduced curve on  $\tilde{\mathcal{S}}_d$  whose support is equal to  $\tilde{\eta}(\tilde{\Sigma}_1)$ . On the other hand, the class of the total transform  $(\gamma \circ \rho)^*B$  of  $B$  by  $\gamma \circ \rho$  is equal to  $(q+1)h$ . Since the difference of the divisors  $d \cdot R$  and  $(\gamma \circ \rho)^*B$  is a linear combination of exceptional curves of  $\rho$ , we have

$$(5.3) \quad \tilde{\eta}_*([\tilde{\Sigma}_1]) \in V.$$

By the commutativity of the diagram (5.2), we have

$$\tilde{\eta}(\Xi) \subset \rho^{-1} \left( \gamma^{-1} \left( \bigcup_{P \in \mathbf{P}^1(\mathbf{F}_{q^2})} l_P \right) \right).$$

Hence, for any irreducible component  $\Gamma$  of  $\Xi$ , we have

$$(5.4) \quad \tilde{\eta}_*([\Gamma]) \in V.$$

Let  $C$  be an arbitrary irreducible curve on  $\tilde{\mathcal{S}}_d$ . Then we have

$$\tilde{\eta}_* \tilde{\eta}^*([C]) = 2q[C].$$

By Lemma 5.1, there exist integers  $a, b_1, \dots, b_m$  and irreducible components  $\Gamma_1, \dots, \Gamma_m$  of  $\Xi$  such that the divisor  $\eta^*C$  of  $\tilde{M}$  is linearly equivalent to

$$a\tilde{\Sigma}_1 + b_1\Gamma_1 + \cdots + b_m\Gamma_m.$$

By (5.3) and (5.4), we obtain

$$[C] = \frac{1}{2q} \tilde{\eta}_* \tilde{\eta}^*([C]) \in V.$$

Therefore  $V \subset H^2(\tilde{\mathcal{S}}_d)$  is equal to the linear subspace spanned by the classes of all curves. Combining this fact with Corollary 1.7, we obtain  $V = H^2(\tilde{\mathcal{S}}_d)$ .

## 6. Supersingular $K3$ surfaces

In this section, we prove Proposition 1.9. First, we recall some facts on supersingular  $K3$  surfaces. Let  $Y$  be a supersingular  $K3$  surface in characteristic

$p$ , and let  $\text{NS}(Y)$  denote its Néron-Severi lattice, which is an even hyperbolic lattice of rank 22. Artin [3] showed that the discriminant of  $\text{NS}(Y)$  is written as  $-p^{2\sigma}$ , where  $\sigma$  is a positive integer  $\leq 10$ . This integer  $\sigma$  is called the *Artin invariant* of  $Y$ . Ogus [13, 14] and Rudakov-Shafarevich [15] proved that, for each  $p$ , a supersingular  $K3$  surface with Artin invariant 1 is unique up to isomorphism. Let  $X_p$  denote the supersingular  $K3$  surface with Artin invariant 1 in characteristic  $p$ . It is known that  $X_3$  is isomorphic to the Fermat quartic surface, and that  $X_5$  is isomorphic to the Fermat sextic double plane. (See, for example, [12] and [17], respectively.) Therefore, in order to prove Proposition 1.9, it is enough to prove the following:

**PROPOSITION 6.1.** *Suppose that  $(d, q + 1) = (4, 4)$  or  $(2, 6)$ . Then, among the curves on  $\tilde{S}_d$  listed in Proposition 1.8, there exist 22 curves whose classes together with the intersection pairing form a lattice of rank 22 with discriminant  $-p^2$ .*

*Proof.* Suppose that  $p = q = 3$  and  $d = 4$ . We put  $\alpha := \sqrt{-1} \in \mathbf{F}_9$ , so that  $\mathbf{F}_9 := \mathbf{F}_3(\alpha)$ . Consider the projective space  $\mathbf{P}^3$  with homogeneous coordinates  $[w : x_0 : x_1 : x_2]$ . By Proposition 1.4, the surface  $S_4$  is defined in  $\mathbf{P}^3$  by an equation

$$w^4 = 2(x_0^3 x_1 + x_0 x_1^3) - x_2^4 - (x_2^2 - x_1 x_0)^2.$$

Hence the singular locus  $\text{Sing}(S_4)$  of  $S_4$  consists of the three points

$$\begin{aligned} Q_0 &:= [0 : 1 : 1 : 0] \quad (\text{located over } \phi([1 : \alpha]) = \phi([1 : -\alpha]) \in B), \\ Q_1 &:= [0 : 1 : 2 : 1] \quad (\text{located over } \phi([1 : 1 + \alpha]) = \phi([1 : 1 - \alpha]) \in B), \\ Q_2 &:= [0 : 1 : 2 : 2] \quad (\text{located over } \phi([1 : 2 + \alpha]) = \phi([1 : 2 - \alpha]) \in B), \end{aligned}$$

and they are rational double points of type  $A_3$ . The minimal resolution  $\rho : \tilde{S}_4 \rightarrow S_4$  is obtained by blowing up twice over each singular point  $Q_a$  ( $a \in \mathbf{F}_3$ ). The rational curves  $l_p^{(i)}$  on  $\tilde{S}_4$  given in Proposition 1.8 are the strict transforms of the following 40 lines  $\bar{L}_\tau^{(v)}$  in  $\mathbf{P}^3$  contained in  $S_4$ , where  $v = 0, \dots, 3$ :

$$\begin{aligned} \bar{L}_0^{(v)} &:= \{x_1 = w - \alpha^v x_2 = 0\}, \\ \bar{L}_1^{(v)} &:= \{x_0 + x_1 - x_2 = w - \alpha^v (x_2 + x_0) = 0\}, \\ \bar{L}_2^{(v)} &:= \{x_0 + x_1 + x_2 = w - \alpha^v (x_2 - x_0) = 0\}, \\ \bar{L}_\infty^{(v)} &:= \{x_0 = w - \alpha^v x_2 = 0\}, \\ \bar{L}_{\pm\alpha}^{(v)} &:= \{-x_0 + x_1 \pm \alpha x_2 = w - \alpha^v x_2 = 0\}, \\ \bar{L}_{1\pm\alpha}^{(v)} &:= \{\pm\alpha x_0 + x_1 + (-1 \pm \alpha)x_2 = w - \alpha^v (x_2 + x_0) = 0\}, \\ \bar{L}_{2\pm\alpha}^{(v)} &:= \{\mp\alpha x_0 + x_1 + (1 \pm \alpha)x_2 = w - \alpha^v (x_2 - x_0) = 0\}. \end{aligned}$$

We denote by  $L_\tau^{(v)}$  the strict transform of  $\bar{L}_\tau^{(v)}$  by  $\rho$ . Note that the image of  $\bar{L}_\tau^{(v)}$  by the covering morphism  $S_4 \rightarrow \mathbf{P}^2$  is the line  $l_{\phi([1:\tau])}$ . Note also that, if  $\tau \in \mathbf{F}_3 \cup \{\infty\}$ , then  $\bar{L}_\tau^{(v)}$  is disjoint from  $\text{Sing}(S_4)$ , whereas if  $\tau = a + b\alpha \in \mathbf{F}_9 \setminus \mathbf{F}_3$  with  $a \in \mathbf{F}_3$  and  $b \in \mathbf{F}_3 \setminus \{0\} = \{\pm 1\}$ , then  $\bar{L}_\tau^{(v)} \cap \text{Sing}(S_4)$  consists of a single point  $Q_a$ . Looking at the minimal resolution  $\rho$  over  $Q_a$  explicitly, we see that the three exceptional  $(-2)$ -curves in  $\tilde{S}_4$  over  $Q_a$  can be labeled as  $E_{a-\alpha}$ ,  $E_a$ ,  $E_{a+\alpha}$  in such a way that the following hold:

- $\langle E_{a-\alpha}, E_a \rangle = \langle E_a, E_{a+\alpha} \rangle = 1$ ,  $\langle E_{a-\alpha}, E_{a+\alpha} \rangle = 0$ .
- Suppose that  $b \in \{\pm 1\}$ . Then  $L_{a+b\alpha}^{(v)}$  intersects  $E_{a+b\alpha}$ , and is disjoint from the other two irreducible components  $E_a$  and  $E_{a-b\alpha}$ .
- The four intersection points of  $L_{a+b\alpha}^{(v)}$  ( $v = 0, \dots, 3$ ) and  $E_{a+b\alpha}$  are distinct.

Using these, we can calculate the intersection numbers among the  $9 + 40$  curves  $E_\tau$  and  $L_{\tau'}^{(v)}$  ( $\tau \in \mathbf{F}_9$ ,  $\tau' \in \mathbf{F}_9 \cup \{\infty\}$ ,  $v = 0, \dots, 3$ ). From among them, we choose the following 22 curves:

$$\begin{aligned} & E_{-\alpha}, E_0, E_\alpha, E_{1-\alpha}, E_1, E_{1+\alpha}, E_{2-\alpha}, E_2, E_{2+\alpha}, \\ & L_0^{(0)}, L_0^{(1)}, L_0^{(2)}, L_0^{(3)}, L_1^{(0)}, L_1^{(1)}, L_2^{(0)}, L_2^{(1)}, L_\infty^{(1)}, \\ & L_{-\alpha}^{(0)}, L_{-\alpha}^{(1)}, L_{1-\alpha}^{(2)}, L_{2-\alpha}^{(0)}. \end{aligned}$$

Their intersection numbers are calculated as in Table 6.1. We can easily check that this matrix is of determinant  $-9$ . Therefore the Artin invariant of  $\tilde{S}_4$  is 1.

-2	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	
1	-2	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
0	1	-2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
0	0	0	-2	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
0	0	0	1	-2	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	1	-2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	-2	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
0	0	0	0	0	0	1	-2	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	1	-2	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	1	-2	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	1	0	0	0	0
0	0	0	0	0	0	0	0	0	1	1	-2	0	1	0	1	0	0	0	0	0	1	1
0	0	0	0	0	0	0	0	0	0	0	1	0	-2	1	0	0	0	0	1	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	1	1	-2	0	0	1	0	0	0	0	1
0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	-2	1	0	0	0	0	1
0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	-2	1	1	0	1	0
0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	1	-2	0	1	0	0	0
1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	-2	0	0	0	0
1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	-2	1	1	0
0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	1	-2	0	0
0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	1	0	0	0	1	0	-2	0

 Table 6.1. Gram matrix of  $\text{NS}(\tilde{S}_4)$  for  $q = 3$

The proof for the case  $p = q = 5$  and  $d = 2$  is similar. We put  $\alpha := \sqrt{2}$  so that  $\mathbf{F}_{25} = \mathbf{F}_5(\alpha)$ . In the weighted projective space  $\mathbf{P}(3, 1, 1, 1)$  with homogeneous coordinates  $[w : x_0 : x_1 : x_2]$ , the surface  $S_2$  for  $p = q = 5$  is defined by

$$w^2 = 2(x_0^5 x_1 + x_0 x_1^5) - x_2^6 - (x_2^2 + x_0 x_1)^3.$$

The singular locus  $\text{Sing}(S_2)$  consists of ten ordinary nodes

$$Q_{\{a+b\alpha, a-b\alpha\}} \quad (a \in \mathbf{F}_5, b \in \{1, 2\})$$

located over the nodes  $\phi([1 : a + b\alpha]) = \phi([1 : a - b\alpha])$  of the branch curve  $B$ . Let  $E_{\{a+b\alpha, a-b\alpha\}}$  denote the exceptional  $(-2)$ -curve in  $\tilde{S}_2$  over  $Q_{\{a+b\alpha, a-b\alpha\}}$  by the minimal resolution. As the 22 curves, we choose the following eight exceptional  $(-2)$ -curves

$$\begin{aligned} &E_{\{-\alpha, \alpha\}}, \quad E_{\{-2\alpha, 2\alpha\}}, \quad E_{\{1-\alpha, 1+\alpha\}}, \quad E_{\{1-2\alpha, 1+2\alpha\}}, \\ &E_{\{2-\alpha, 2+\alpha\}}, \quad E_{\{3-2\alpha, 3+2\alpha\}}, \quad E_{\{4-\alpha, 4+\alpha\}}, \quad E_{\{4-2\alpha, 4+2\alpha\}}, \end{aligned}$$

and the strict transforms of the following 14 curves on  $S_2$ :

$$\begin{aligned} &\{x_1 = w - 2\alpha x_2^3 = 0\}, \\ &\{x_1 = w + 2\alpha x_2^3 = 0\}, \\ &\{x_0 + x_1 + 4x_2 = w + 2\alpha(3x_0 + x_2)^3 = 0\}, \\ &\{3x_0 + x_1 + 3\alpha x_2 = w - 2\alpha x_2^3 = 0\}, \\ &\{2x_0 + x_1 + 4\alpha x_2 = w + 2\alpha x_2^3 = 0\}, \\ &\{3x_0 + x_1 + 2\alpha x_2 + 3x_0 = w - 2\alpha x_2^3 = 0\}, \\ &\{(3 + 3\alpha)x_0 + x_1 + (4 + \alpha)x_2 = w + 2\alpha(3x_0 + x_2)^3 = 0\}, \\ &\{(4 + \alpha)x_0 + x_1 + (4 + 2\alpha)x_2 = w + 2\alpha(3x_0 + x_2)^3 = 0\}, \\ &\{(2 + 3\alpha)x_0 + x_1 + (3 + 3\alpha)x_2 = w - 2\alpha(x_0 + x_2)^3 = 0\}, \\ &\{(1 + \alpha)x_0 + x_1 + (3 + \alpha)x_2 = w - 2\alpha(x_0 + x_2)^3 = 0\}, \\ &\{(1 + \alpha)x_0 + x_1 + (2 + 4\alpha)x_2 = w - 2\alpha(x_2 + 4x_0)^3 = 0\}, \\ &\{(2 + 3\alpha)x_0 + x_1 + (2 + 2\alpha)x_2 = w + 2\alpha(x_2 + 4x_0)^3 = 0\}, \\ &\{(3 + 3\alpha)x_0 + x_1 + (1 + 4\alpha)x_2 = w - 2\alpha(x_2 + 2x_0)^3 = 0\}, \\ &\{(4 + 4\alpha)x_0 + x_1 + (1 + 2\alpha)x_2 = w - 2\alpha(x_2 + 2x_0)^3 = 0\}. \end{aligned}$$

Their intersection matrix is given in Table 6.2. It is of determinant  $-25$ . Therefore the Artin invariant of  $\tilde{S}_2$  is 1.  $\square$



- [10] T. KATSURA AND S. KONDŌ, Rational curves on the supersingular  $K3$  surface with Artin invariant 1 in characteristic 3, *J. Algebra* **352** (2012), 299–321.
- [11] T. KATSURA, S. KONDŌ AND I. SHIMADA, On the supersingular  $K3$  surface in characteristic 5 with Artin invariant 1, *Michigan Math. J.* **63** (2014), 803–844.
- [12] S. KONDŌ AND I. SHIMADA, The automorphism group of a supersingular  $K3$  surface with Artin invariant 1 in characteristic 3, *Int. Math. Res. Not. IMRN* **2014** (2014), 1885–1924.
- [13] A. OGUS, Supersingular  $K3$  crystals, *Journées de Géométrie Algébrique de Rennes (Rennes, 1978)*, Vol. II, Astérisque **64**, Soc. Math. France, Paris, 1979, 3–86.
- [14] A. OGUS, A crystalline Torelli theorem for supersingular  $K3$  surfaces, *Arithmetic and geometry*, Vol. II, *Progr. Math.* **36**, Birkhäuser Boston, Boston, MA, 1983, 361–394.
- [15] A. N. RUDAKOV AND I. R. SHAFAREVICH, Surfaces of type  $K3$  over fields of finite characteristic, *Current problems in mathematics* **18**, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Informatsii, Moscow, 1981, 115–207. Reprinted in I. R. Shafarevich, *Collected mathematical papers*, Springer-Verlag, Berlin, 1989, 657–714.
- [16] I. SHIMADA, Unirationality of certain complete intersections in positive characteristics, *Tohoku Math. J.* **44** (1992), 379–393.
- [17] I. SHIMADA, Projective models of the supersingular  $K3$  surface with Artin invariant 1 in characteristic 5, *J. Algebra* **403** (2014), 273–299.
- [18] T. SHIODA, An example of unirational surfaces in characteristic  $p$ , *Math. Ann.* **211** (1974), 233–236.
- [19] T. SHIODA AND T. KATSURA, On Fermat varieties, *Tohoku Math. J.* **31** (1979), 97–115.

Thanh Hoai Hoang  
DEPARTMENT OF MATHEMATICS  
GRADUATE SCHOOL OF SCIENCE  
HIROSHIMA UNIVERSITY  
1-3-1 KAGAMIYAMA  
HIGASHI-HIROSHIMA, 739-8526  
JAPAN  
E-mail: hoangthanh2127@yahoo.com

Ichiro Shimada  
DEPARTMENT OF MATHEMATICS  
GRADUATE SCHOOL OF SCIENCE  
HIROSHIMA UNIVERSITY  
1-3-1 KAGAMIYAMA  
HIGASHI-HIROSHIMA, 739-8526  
JAPAN  
E-mail: shimada@math.sci.hiroshima-u.ac.jp