

On the equivalence of several definitions of compact infra-solvmanifolds

By Shintarô KUROKI^{*,**} and Li YU^{***}

(Communicated by Kenji FUKAYA, M.J.A., Oct. 15, 2013)

Abstract: We show the equivalence of several definitions of compact infra-solvmanifolds that appear in various math literatures.

Key words: Infra-solvmanifold; solvable Lie group; homogeneous space; discrete group action; holonomy.

Infra-solvmanifolds are a special class of aspherical manifolds studied by mathematicians for a long time. By the literal meaning, an infra-solvmanifold is a smooth manifold finitely covered by a solvmanifold (which is the quotient of a connected solvable Lie group by a closed subgroup). Infra-solvmanifolds include all flat Riemannian manifolds and infra-nilmanifolds.

It is known that compact infra-solvmanifolds are *smoothly rigid*, i.e. any homotopy equivalence between two compact infra-solvmanifolds is homotopic to a diffeomorphism (see [3] and [8]). Geometrically, a compact infra-solvmanifold M can be described (see [7, Proposition 3.1]) as a manifold which admits a sequence of Riemannian metric $\{g_n\}$ with uniformly bounded sectional curvature so that (M, g_n) collapses in the *Gromov-Hausdorff* sense to a *flat orbifold*.

In the history of study on this topic, several definitions of compact infra-solvmanifolds were proposed in various math literatures. This is mainly because people look at these manifolds from very different angles (topologically, algebraically or geometrically). The following are five such definitions.

Definition 1. Let G be a connected, simply connected solvable Lie group, K be a maximal compact subgroup of the group $\text{Aut}(G)$ of auto-

morphisms of G , and Γ be a discrete subgroup of $G \rtimes K$. If $[\Gamma : G \cap \Gamma] < \infty$ and the action of Γ on G is cocompact and free, the orbit space $\Gamma \backslash G$ is called an *infra-solvmanifold modeled on G* . See [7, Definition 1.1].

Definition 2. A *compact infra-solvmanifold* is a manifold of the form $\Gamma \backslash G$, where G is a connected, simply connected solvable Lie group, and Γ is a torsion-free discrete subgroup of $\text{Aff}(G) = G \rtimes \text{Aut}(G)$ which acts on G cocompactly and the closure of $\text{hol}(\Gamma)$ in $\text{Aut}(G)$ is compact where $\text{hol} : \text{Aff}(G) \rightarrow \text{Aut}(G)$ is the holonomy projection.

Definition 3. A *compact infra-solvmanifold* is a double coset space $\Gamma \backslash G / K$ where G is a virtually connected and virtually solvable Lie group, K is a maximal compact subgroup of G and Γ is a torsion-free, cocompact, discrete subgroup of G . See [2, Definition 2.10].

A *virtually connected* Lie group is a Lie group with finitely many connected components.

Definition 4. A *compact infra-solvmanifold* M is an orbit space of the form $M = \Delta \backslash S$ where S is a connected, simply connected solvable Lie group acted upon cocompactly by a torsion-free closed subgroup $\Delta \subset \text{Aff}(S)$ satisfying

- the identity component Δ_0 of Δ is contained in the nil-radical of S ,
- the closure of $\text{hol}(\Delta)$ in $\text{Aut}(S)$ is compact.

Definition 5. A *compact infra-solvmanifold* is a manifold of the form $\Delta \backslash G$, where G is a connected, simply connected solvable Lie group, and Δ is a torsion-free subgroup of $\text{Aff}(G)$ which acts cocompactly on G and the closure of $\text{hol}(\Delta)$ in $\text{Aut}(G)$ is compact.

Note that the subgroup Δ in Definition 4 and Definition 5 is not necessarily discrete. The Defi-

2010 Mathematics Subject Classification. Primary 22E25, 22E40, 22F30; Secondary 53C30.

^{*}) Osaka City University Advanced Mathematical Institute (OCAMI), 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan.

^{**}) Department of Mathematics, University of Toronto, Room 6290, 40 St. George Street, Toronto, Ontario, M5S 2E4, Canada.

^{***}) Department of Mathematics and IMS, Nanjing University, Nanjing, 210093, P. R. China.


inition 4 is from [3, Definition 1.1] and Definition 5 is from [1, Definition 1.1].

The main purpose of this paper is to explain why the above five definitions of compact infra-solvmanifolds are equivalent. The authors are fully aware that the reason is probably known to many people. However, we do not find any formal proof of this equivalence and feel this phenomenon a little confusing. So we want to write a proof here for the convenience of future reference. However, we do not intend to give a completely new treatment of this subject. Our proof will directly quote some results on infra-solvmanifolds from [1], [3] and [8] and use many subtle facts in Lie theory. This paper can be treated as an elementary exposition of compact infra-solvmanifolds and some related concepts.

Now let us start to prove the following main result of the paper.

Proposition 1. *The five definitions Definition 1–Definition 5 of compact infra-solvmanifolds are all equivalent.*

Proof. We divide our proof into several parts. The framework of the proof is shown below.



Definition 1 \iff **Definition 2**. Let M be a compact infra-solvmanifold in the sense of Definition 1. First of all, any $g \in \Gamma$ can be decomposed as $g = k_g u_g$ where $k_g \in K \subset \text{Aut}(G)$ and $u_g \in G$. The holonomy projection $hol : G \rtimes \text{Aut}(G) \rightarrow \text{Aut}(G)$ sends g to k_g . Since hol is a group homomorphism, its image $hol(\Gamma)$ is a subgroup of K . By assumption, $|hol(\Gamma)| = [\Gamma : G \cap \Gamma]$ is finite, so $hol(\Gamma)$ is compact. In addition, since G is a connected simply-connected solvable Lie group, G is diffeomorphic to an Euclidean space. Then by Smith fixed point theorem ([6, Theorem I]), Γ acting on G freely implies that Γ is torsion-free. So M satisfies Definition 2.

Conversely, if M is a compact infra-solvmanifold in Definition 2, then [8, Theorem 3 (a) \implies (f)] tells us that there exists a connected, simply-connected solvable Lie group G' so that the Γ (which defines M) can be thought of as discrete subgroup of $G' \rtimes F$ where F is a finite subgroup of

$\text{Aut}(G')$. Moreover, there exists an equivariant diffeomorphism from G to G' with respect to the action of Γ (by [8, Theorem 1 and Theorem 2]). Hence

$$M = \Gamma \backslash G \underset{\text{diff.}}{\cong} \Gamma \backslash G'.$$

Then $[\Gamma : \Gamma \cap G'] = |hol_{G'}(\Gamma)| \leq |F|$ is finite.

It remains to show that the action of Γ on G' is free. Since F is finite, we can choose an F -invariant Riemannian metric on G' (in fact we only need F to be compact). If the action of an element $g \in \Gamma$ on G' has a fixed point, say h_0 . Let L_{h_0} be the left translation of G' by h_0 . Then $L_{h_0}^{-1} g L_{h_0}$ fixes the identity element, which implies $L_{h_0}^{-1} g L_{h_0} \in F$. So there exists a compact neighborhood U of h_0 so that $g \cdot U = U$ and g acts isometrically on U with respect to the Riemannian metric just as the element $L_{h_0}^{-1} g L_{h_0} \in F$ acts around e . So

$$A = Dg : T_{h_0} U \rightarrow T_{h_0} U$$

is an orthogonal transformation. Then A is conjugate in $O(n)$ to the block diagonal matrix of the form

$$\begin{pmatrix} B(\theta_1) & & 0 \\ & \ddots & \\ 0 & & B(\theta_m) \end{pmatrix},$$

where $B(0) = 1$, $B(\pi) = -1$, and

$$B(\theta_j) = \begin{pmatrix} \cos \theta_j & -\sin \theta_j \\ \sin \theta_j & \cos \theta_j \end{pmatrix}, \quad 1 \leq j \leq m.$$

Since Γ is torsion-free, g is an infinite order element. This implies that at least one θ_j is irrational. Then for any $v \neq 0 \in T_{h_0} G'$, $|\{A^n v\}_{n \in \mathbb{Z}}| = \infty$. So there exists $h \in U$ so that $|\{g^n \cdot h\}_{n \in \mathbb{Z}}| = \infty$. Then the set $\{g^n \cdot h\}_{n \in \mathbb{Z}} \subset U$ has at least one accumulation point. This contradicts the fact that the orbit space $\Gamma \backslash G'$ is Hausdorff (since $\Gamma \backslash G' = M$ is a manifold).

Combing the above arguments, M is an infra-solvmanifold modeled on G' in the sense of Definition 1.

Definition 1 \implies **Definition 3**. Let M be an infra-solvmanifold in the sense of Definition 1. Let $hol(\Gamma)$ be the image of the the holonomy projection $hol : \Gamma \rightarrow \text{Aut}(G)$. Then $hol(\Gamma)$ is a finite subgroup of $\text{Aut}(G)$.

Define $\tilde{G} = G \rtimes hol(\Gamma)$ which is virtually solvable. It is easy to see that $hol(\Gamma)$ is a maximal

compact subgroup of \tilde{G} and, Γ is a cocompact, discrete subgroup of \tilde{G} . Then $M \cong \Gamma \backslash \tilde{G} / \text{hol}(\Gamma)$. So M satisfies Definition 3.

Definition 3 \implies Definition 2. Let $M = \Gamma \backslash G / K$ be a compact infra-solvmanifold in the sense of Definition 3. Since K is a maximal compact subgroup of G , G / K is contractible. Note that K is not necessarily a normal subgroup of G . Hence G / K may not directly inherit a group structure from G .

Let G_0 be the connected component of G containing the identity element. Then by [4, Theorem 14.1.3 (ii)], $K_0 = K \cap G_0$ is connected and K_0 is a maximal compact subgroup of G_0 . Moreover, K intersects each connected component of G and

$$K / K_0 \cong G / G_0.$$

By the classical Lie theory, the Lie algebra of a compact Lie group is a direct product of an abelian Lie algebra and some simple Lie algebras. Then since the Lie algebra $\text{Lie}(G)$ of G is solvable and K is compact, the Lie algebra $\text{Lie}(K) \subset \text{Lie}(G)$ must be abelian. This implies that K_0 is a torus and hence a maximal torus in G_0 .

In addition, since G is virtually solvable, G_0 is actually solvable. This is because the radical R of G is a normal subgroup of G_0 and $\dim(R) = \dim(G_0)$ (since G is virtually solvable). So G_0 / R is discrete. Then since G_0 is connected, G_0 must be equal to R .

Let $Z(G)$ be the center of G and define

$$C = Z(G) \cap K.$$

Then C is clearly a normal subgroup of G . Let

$$G' = G / C, \quad K' = K / C.$$

Let $\rho : G \rightarrow G'$ be the quotient map. Since $\Gamma \cap K = \{1\}$, $\Gamma \cong \rho(\Gamma) \subset G'$, we can think of Γ as a subgroup of G' . So we have

$$(1) \quad M = \Gamma \backslash G / K \cong \Gamma \backslash G' / K'.$$

Let $G'_0 = \rho(G_0)$ be the identity component of G' . Then G'_0 is a finite index solvable normal subgroup of G' .

$$(2) \quad \begin{aligned} \text{Lie}(G'_0) &= \text{Lie}(G') = \text{Lie}(G) / \text{Lie}(C) \\ &= \text{Lie}(G_0) / \text{Lie}(C). \end{aligned}$$

Besides, let $K'_0 = K' \cap G'_0$ which is a maximal torus of G'_0 . We have

$$(3) \quad \begin{aligned} \text{Lie}(K'_0) &= \text{Lie}(K') = \text{Lie}(K) / \text{Lie}(C) \\ &= \text{Lie}(K_0) / \text{Lie}(C). \end{aligned}$$

Claim-1. G'_0 is linear and so G' is linear.

A group is called *linear* if it admits a faithful finite-dimensional representation. By [4, Theorem 16.2.9 (b)], a connected solvable Lie group S is linear if and only if $\mathfrak{t} \cap [\mathfrak{s}, \mathfrak{s}] = \{0\}$ where \mathfrak{s} and \mathfrak{t} are Lie algebras of S and its maximal torus T_S , respectively. And for a general connected solvable group S , the Lie subalgebra $\mathfrak{t} \cap [\mathfrak{s}, \mathfrak{s}]$ is always central in \mathfrak{s} . So for our G_0 and its maximal torus K_0 , the Lie subalgebra $\text{Lie}(K_0) \cap [\text{Lie}(G_0), \text{Lie}(G_0)]$ is central in $\text{Lie}(G_0) = \text{Lie}(G)$. So $\text{Lie}(K_0) \cap [\text{Lie}(G_0), \text{Lie}(G_0)] \subset \text{Lie}(Z(G))$ and then

$$(4) \quad \begin{aligned} \text{Lie}(K_0) \cap [\text{Lie}(G_0), \text{Lie}(G_0)] \\ \subset \text{Lie}(K) \cap \text{Lie}(Z(G)) = \text{Lie}(C). \end{aligned}$$

Then for the Lie group G'_0 and its maximal torus K'_0 , we have

$$(5) \quad \begin{aligned} \text{Lie}(K'_0) \cap [\text{Lie}(G'_0), \text{Lie}(G'_0)] \\ = \frac{\text{Lie}(K_0)}{\text{Lie}(C)} \cap \left[\frac{\text{Lie}(G_0)}{\text{Lie}(C)}, \frac{\text{Lie}(G_0)}{\text{Lie}(C)} \right] = 0. \end{aligned}$$

So by [4, Theorem 16.2.9 (b)], G'_0 is linear. Moreover, suppose V is a faithful finite-dimensional representation of G'_0 . Then $\mathbf{R}[G'] \otimes_{\mathbf{R}[G'_0]} V$ is a faithful finite-dimensional representation of G' where $\mathbf{R}[G']$ and $\mathbf{R}[G'_0]$ are the group rings of G' and G'_0 over \mathbf{R} , respectively. So the Claim-1 is proved.

From the Claim-1 and (1), we can just assume that our group G is linear at the beginning. Under this assumption, G_0 is a connected linear solvable group. So there exists a simply connected solvable normal Lie subgroup S of G_0 so that $G_0 = S \rtimes K_0$ and $[G_0, G_0] \subset S$ (see [4, Lemma 16.2.3]). So

$$\text{Lie}(G_0) = \text{Lie}(K_0) \oplus \text{Lie}(S).$$

More specifically, we can take $S = p^{-1}(V)$ where $p : G_0 \rightarrow G_0 / [G_0, G_0]$ is the quotient map and V is a vector subgroup of the abelian group $G_0 / [G_0, G_0]$ so that $G_0 / [G_0, G_0] \cong p(K_0) \times V$ (see the proof of [4, Theorem 16.2.3]). Note that the vector subgroup V is not unique, nor is S .

Claim-2. We can choose S to be normal in G and so $G \cong S \times K$.

Indeed since K is compact, we can choose a metric on $\text{Lie}(G_0)$ which is invariant under the adjoint action of K . Then we can choose V so that $\text{Lie}(S)$ is orthogonal to $\text{Lie}(K_0)$ in $\text{Lie}(G_0)$. Then because K_0 is normal in K , the adjoint action of K on $\text{Lie}(G_0)$ preserves $\text{Lie}(K_0)$, so it also preserves

the orthogonal complement $\text{Lie}(S)$ of $\text{Lie}(K_0)$. This implies that S is preserved under the adjoint action of K .

Let $G_0, h_1G_0, \dots, h_mG_0$ be all the connected components of G . Since K intersects each connected component of G , we can assume that $h_i \in K$ for all $1 \leq i \leq m$. Then any element $g \in G$ can be written as $g = g_0h_i$ for some $g_0 \in G_0$ and $h_i \in K$. So $gSg^{-1} = g_0h_iSh_i^{-1}g_0^{-1} \subset g_0Sg_0^{-1} \subset S$. The Claim-2 is proved.

From the semidirect product $G = S \rtimes K$, we can define an injective group homomorphism $\alpha : G \rightarrow \text{Aff}(S) = S \rtimes \text{Aut}(S)$ as follows. For any $g \in G$, we can write $g = s_gk_g$ for a unique $s_g \in S$ and $k_g \in K$ since $S \cap K = S \cap K_0 = \{1\}$. Then $\alpha(g) : S \rightarrow S$ is the composition of the adjoint action of k_g on S and the left translation on S by s_g , i.e.

$$\alpha(g) = L_{s_g} \circ \text{Ad}_{k_g}.$$

Claim-3. $\alpha(\Gamma) \backslash S$ is diffeomorphic to the double coset space $\Gamma \backslash G / K$.

Notice that each left coset in G/K contains a unique element of S , so we have

$$G/K = \{sK; s \in S\}.$$

For any $\gamma \in \Gamma$, let $\gamma = s_\gamma k_\gamma$ where $s_\gamma \in S$ and $k_\gamma \in K$. Then we have

$$\begin{aligned} \gamma sK &= s_\gamma k_\gamma sK = s_\gamma k_\gamma s k_\gamma^{-1} K \\ &= \alpha(\gamma)(s)K, \forall s \in S. \end{aligned}$$

So the natural action of Γ on the left coset space G/K can be identified with the action of $\alpha(\Gamma) \subset \text{Aff}(S)$ on S . The Claim-3 is proved.

Let $\text{Ad} : K \rightarrow \text{Aut}(S)$ denote the adjoint action of K on S . Since K is compact and Ad is continuous, $\text{Ad}(K) \subset \text{Aut}(S)$ is also compact. Notice that $\alpha(\Gamma)$ is a subgroup of $S \rtimes \text{Ad}(K) \subset S \rtimes \text{Aut}(S)$. So the closure $\overline{\text{hol}(\alpha(\Gamma))}$ of the holonomy group $\text{hol}(\alpha(\Gamma))$ in $\text{Aut}(S)$ is contained in $\text{Ad}(K)$. So $\overline{\text{hol}(\alpha(\Gamma))}$ is compact. This implies that $\Gamma \backslash G / K \cong \alpha(\Gamma) \backslash S$ is an infra-solvmanifold in the sense of Definition 2.

Remark 1. A simply-connected solvable Lie group is always linear, but for non-simply-connected solvable Lie groups, this is not always so. A counterexample is the quotient group of the Heisenberg group by an infinite cyclic group (see [5, p. 169 Example 5.67]).

It is obviously true that Definition 2 \implies Definition 4 \implies Definition 5. So it remains to show Definition 5 \implies Definition 2.

Definition 5 \implies Definition 2. This follows from a nontrivial result in [1]. Indeed, the fundamental group of the manifold $\Delta \backslash G$ in Definition 5 is $\Gamma = \Delta / \Delta_0$, which is a torsion-free virtually polycyclic group (see [2]). It is shown in [1] that such a group Γ determines a virtually solvable real linear algebraic group H_Γ which contains Γ as a discrete and Zariski-dense subgroup. H_Γ is called the *real algebraic hull* of Γ in [1]. In addition, we have $H_\Gamma = U \rtimes T$ where T is a maximal reductive subgroup of H_Γ and U is the unipotent radical of H_Γ . The splitting of H_Γ gives an injective group homomorphism $\alpha : H_\Gamma \rightarrow \text{Aff}(U)$ and a corresponding affine action of $\Gamma < H_\Gamma$ on U . It is shown in [1] that $M_\Gamma = \alpha(\Gamma) \backslash U$ is a compact infra-solvmanifold whose fundamental group is Γ . The M_Γ is called the *standard Γ -manifold*. Now since $\alpha(\Gamma)$ is a discrete subgroup of $\text{Aff}(U)$, M_Γ is a compact infra-solvmanifold in the sense of Definition 2. Moreover $\Delta \backslash G$ is diffeomorphic to M_Γ by [1, Theorem 1.4].

So we finish the proof of Proposition 1. \square

Remark 2. If we remove the ‘‘cocompact’’ condition in Definition 1–Definition 5, we may get noncompact infra-solvmanifolds, which are vector bundles over some compact infra-solvmanifolds (see [8, Theorem 6]).

Acknowledgements. The first author is supported in part by the JSPS Strategic Young Researcher Overseas Visits Program for Accelerating Brain Circulation ‘‘Deepening and Evolution of Mathematics and Physics, Building of International Network Hub based on OCAMI’’. The second author is supported by Natural Science Foundation of China (Grant no. 11001120) and the PAPD (priority academic program development) of Jiangsu higher education institutions. In addition, both authors want to thank Profs. M. Masuda, J. B. Lee, W. Tuschmann, J. Hillman and F. Farrell for some helpful information and comments. Besides, the first author wants to thank Prof. Y. Karshon for providing him excellent circumstances to do research.

References

- [1] O. Baues, Infra-solvmanifolds and rigidity of subgroups in solvable linear algebraic groups, *Topology* **43** (2004), no. 4, 903–924.
- [2] F. T. Farrell and L. E. Jones, *Classical aspherical manifolds*, CBMS Regional Conference Series in Mathematics, 75, Published for the Conference

- Board of the Mathematical Sciences, Washington, DC, 1990.
- [3] F. T. Farrell and L. E. Jones, Compact infrasolvmanifolds are smoothly rigid, in *Geometry from the Pacific Rim (Singapore, 1994)*, 85–97, de Gruyter, Berlin, 1997.
- [4] J. Hilgert and K.-H. Neeb, *Structure and geometry of Lie groups*, Springer Monographs in Mathematics, Springer, New York, 2012.
- [5] K. H. Hofmann and S. A. Morris, *The structure of compact groups*, second revised and augmented edition, de Gruyter Studies in Mathematics, 25, de Gruyter, Berlin, 2006.
- [6] P. A. Smith, Fixed-point theorems for periodic transformations, *Amer. J. Math.* **63** (1941), 1–8.
- [7] W. Tuschmann, Collapsing, solvmanifolds and inhomogeneous spaces, *Differential Geom. Appl.* **7** (1997), no. 3, 251–264.
- [8] B. Wilking, Rigidity of group actions on solvable Lie groups, *Math. Ann.* **317** (2000), no. 2, 195–237.