

An Existence Result for Impulsive Stochastic Functional Differential Equations with Multiple Delays

Anguraj A* and Banupriya K

P.S.G. College of Arts and Science, Coimbatore, Tamil Nadu, India

Abstract

In this paper we consider Impulsive stochastic neutral functional differential equations with multiple delays. By using Schaefer's fixed point theorem, we prove the existence of solutions for stochastic differential equations with impulses.

Keywords: Impulsive differential equations; Stochastic differential equations; Multiple delay; Fixed point theorem

AMS Mathematical Subject Classification: 34A37, 34K50, 34K45, 60H10.

Introduction

The theory of impulsive differential equations is an important area of scientific activity. Many evolution processes are characterized by the fact that at certain moments of time they experience a change of state abruptly. These short term perturbations act instantaneously, that is in the form of impulses. For example, that many biological phenomena involving thresholds, optimal control models in economics and frequency modulated systems, do exhibit impulsive effects. So the impulsive differential equations appear as a natural description of observed evolution phenomena of several real world problems. Existence of solutions of impulsive differential equations has been studied by many authors. If the impulses are random the solution becomes a stochastic process. Existence of solutions of differential equations with random impulses have been studied by many authors [1-3].

Furthermore, besides impulsive effects, stochastic effects likewise exist in real systems. There is a wide range of interesting process in robotics, economics and biology that can be described as differential equations with non-deterministic dynamics such phenomena are described by stochastic differential equations. The solution of stochastic differential equation is a stochastic process. However the solution of differential equation with random impulses is different from the solution of stochastic differential equations. Existence, Uniqueness and qualitative analysis of solutions of stochastic differential equations have discussed by several authors [4,5].

Since both impulsive and stochastic effects exist it is very difficult to investigate the existence of solution of impulsive stochastic differential equations. In [6] Anguraj and Vinodkumar discussed the existence, uniqueness and stability of impulsive stochastic semi linear neutral functional differential with infinite delays. Lakrib [7] discussed about the existence results for impulsive neutral functional differential equations with multiple delays. Based on the existing literature, stochastic impulsive differential equations involved mainly on controllability and stability. To the best of our knowledge, there is no work reported on impulsive stochastic differential equations with multiple delays. The purpose of this paper is to discuss about the existence results of impulsive stochastic neutral functional differential equations with multiple delays. Our approach is based on Schaefer's fixed point theorem.

In this paper we study the existence results for stochastic impulsive differential equations with multiple delays

$$d[x(t) - f(t, x_t)] = [g(t, x_t) + \sum_{i=1}^p x(t - \tau_i)]dt + a(t, x_t)dB_t \quad (1.1)$$

$$a.e., t \in J = [0, 1], t \neq t_k, k = 1, 2, \dots, m.$$

$$\Delta x(t_k) = I_k(x(t_k^-)), k = 1, 2, \dots, m$$

$$x_0 = \varphi$$

where $f : J \times R^n \rightarrow R^n$, $g : J \times R^n \rightarrow R^n$ and $a : J \times R^n \rightarrow R^n$ are Borel Measurable functions, $J_0 = [-r, 0]$, $r = \max \{ \tau_i : i = 1, 2, \dots, p \}$ and $\varphi : [-r, 0] \rightarrow R^n$. Further more the fixed moments of time t_k satisfy $0 = t_0 < t_1 < \dots < t_m < t_{m+1} = 1$, where $x(t_k^+)$ and $x(t_k^-)$ represent the right and left limits of $x(t)$ at $t = t_k$, respectively. And $\Delta x(t = t_k) = x(t_k^+) - x(t_k^-)$, $k = 1, 2, \dots, m$ represent the jump in the state at time t_k with I_k determining the size of the jump.

Preliminaries

Let $(\bar{\Omega}, F\{F_t\}_{t \geq 0}, P)$ be a complete probability space with a filtration $\{F_t\}_{t \geq 0}$ satisfying the conditions that it is right continuous and F_0 contains all P- null sets and $w(t) = (w_1(t), \dots, w_m(t))^T$ is an m-dimensional Brownian motion defined on $(\bar{\Omega}, F\{F_t\}_{t \geq 0}, P)$. Let $C = C[[-r, 0], R^n]$ denote the family of all continuous R^n -valued function φ on $[-r, 0]$ with the norm $\|\varphi\| = \sup_{-r \leq \theta \leq 0} |\varphi(\theta)|$ where $|\cdot|$ is Euclidean norm of R^n . Denote by $C_{F_0}^b[[-r, 0], R^n]$ the family of all bounded F_0 - measurable, $C[[-r, 0], R^n]$ -valued random variables φ , satisfying $\|\varphi\|_{L^2}^2 = \sup_{-r \leq \theta \leq 0} E|\varphi(\theta)|^2 < \infty$, where E denotes the expectation of stochastic process [8-10]. The initial condition $\varphi \in C_{F_0}^b[[-r, 0], R^n]$.

Let $PC(J, R^n)$ the space of piecewise continuous functions $x : J \rightarrow R^n$ such that x is continuous everywhere except for $t = t_k$ at which $x(t_k^-)$ and $x(t_k^+)$ exist and $x(t_k^-) = x(t_k)$, $k = 1, \dots, m$. If we set $\Omega = \{x : J_1 \rightarrow R^n, x \in R^n \cap PC(J, R^n)\}$ where $J_1 = [-r, 1]$ then Ω is a Banach space normed by $\|x\| = \sup \{ |x(t)| : t \in J_1 \}$, $x \in \Omega$.

*Corresponding author: Anguraj A, P.S.G. College of Arts and Science, Coimbatore-641 014, Tamil Nadu, India, Tel: 0422 430 3300; E-mail: angurajpsg@yahoo.com

Received April 11, 2015; Accepted June 09, 2015; Published June 16, 2015

Citation: Anguraj A, Banupriya K (2015) An Existence Result for Impulsive Stochastic Functional Differential Equations with Multiple Delays. J Phys Math 6: 139. doi:10.4172/2090-0902.1000139

Copyright: © 2015 Anguraj A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Obviously, for any $x \in \Omega$ and $t \in J$, the function x_t defined by $x_t(\theta) = x(t + \theta)$, for $\theta \in J_0$, belongs to R_n .

By $L^1(J, R^n)$ we denote the Banach space of measurable functions $x: J \rightarrow R^n$ which are Lebesgue integrable, normed by

$$\|x\| = \int_0^1 x(t) dt$$

Definition 2.1: A stochastic process $\{x(t) \in C_{F_t}^b[[-r, 0], R^n], t \in [-r, 1]\}$, is called a solution of equations (1.1) if

- (i) $x(t)$ is F_t adapted;
- (ii) $x(t)$ satisfies the integral equation

$$x(t) = \begin{cases} \phi(t) & \text{for } t \in J_0, \\ \phi(0) - f(0, \phi(0)) + f(t, x_t) + \int_0^t g(s, x_s) ds + \sum_{i=1}^p \int_{-\tau_i}^0 \phi(s) ds \\ + \sum_{i=1}^p \int_0^{t-\tau_i} x(s) ds + \int_0^t a(s, x_s) dB(s) + \sum_{0 < t_k < t} I_k(x(t_k^-)) & \text{for } t \in J \end{cases} \quad (2.1)$$

Definition 2.2: (Schaefer fixed point theorem). Let X be a normed linear space and let $\Gamma: X \rightarrow X$ be a completely continuous map, that is, it is a continuous mapping which is compact on each bounded subset of X . If the set $\zeta = \{x \in X: \lambda x = \Gamma x \text{ for some } \lambda > 1\}$ is bounded, then Γ has a fixed point [11].

Hypotheses

H_1 : The function $f, g: J \times R^n \rightarrow R^n$ is such that

$$\|f(t, x_t)\|^2 \vee \|g(t, x_t)\|^2 \leq c_1 \|x_t\|^2 + c_2$$

for each $t \in [0, 1]$ and $c_1, c_2 > 0$ are constants.

H_2 : The function $a: J \times R^n \rightarrow R^n$ is Caratheodory, that is,

(i) $t \mapsto a(t, x)$ is measurable for each $x \in R$,

(ii) $x \mapsto a(t, x)$ is continuous for a.e. $t \in J$.

H_3 : There exists a function $q \in L^1(J, R^n)$ with $q(t) > 0$ for a.e. $t \in J$ and a continuous non decreasing function $\psi: [0, \infty) \rightarrow [0, \infty)$ such that

$$|a(t, x)| \leq q(t)\psi(\|x\|)$$

for a.e. $t \in J$ and each $x \in R^n$ with

$$\int_C \frac{ds}{s + \psi(s)} = \infty \quad (3.1)$$

Where

$$C = \frac{1}{1 - 2c_1} \left[\|\phi\| \left(1 + c_1 + \sum_{i=1}^p \tau_i \right) + 3c_2 \right]$$

H_4 : The function $I_k: R \rightarrow R$ and there exist positive constant $c_k, k=1, 2, \dots, m$, such that $\|I_k(x)\| \leq c_k, x \in R$

Theorem 2.1: Suppose that the conditions $(H_1) - (H_4)$ are satisfied then there exists a solution of the problem (1.1) on J_1 [12-14].

Proof: Transform the problem (1.1) - (1.3) into a fixed point problem. Consider the operator $\Gamma: \Omega \rightarrow \Omega$ defined by

$$\Gamma(x(t)) = \begin{cases} \phi(t) \text{ for } t \in J_0, \\ \phi(0) - f(0, \phi(0)) + f(t, x_t) + \int_0^t g(s, x_s) ds + \sum_{i=1}^p \int_{-\tau_i}^0 \phi(s) ds \\ + \sum_{i=1}^p \int_0^{t-\tau_i} x(s) ds + \int_0^t a(s, x_s) dB(s) + \sum_{0 < t_k < t} I_k(x(t_k^-)) \text{ for } t \in J \end{cases}$$

Step 1: Γ has bounded values for bounded sets in Ω .

Let B be a bounded set in Ω . Then there exists a real number $\rho > 0$ such that $E\|x\|^2 \leq \rho$, for all $x \in B$.

Let $x \in B$ and $t \in J$, we have

$$\begin{aligned} \Gamma x(t) &= \phi(0) - f(0, \phi(0)) + f(t, x_t) + \int_0^t g(s, x_s) ds + \sum_{i=1}^p \int_{-\tau_i}^0 \phi(s) ds \\ &+ \sum_{i=1}^p \int_0^{t-\tau_i} x(s) ds + \int_0^t a(s, x_s) dB(s) + \sum_{0 < t_k < t} I_k(x(t_k^-)) \\ E\|\Gamma x(t)\|^2 &\leq E\|\phi\|^2 + c_1 E\|\phi\|^2 + c_2 + c_1 E\|x_t\|^2 + c_2 + E \int_0^t \|g(s, x_s)\|^2 ds + \sum_{i=1}^p \tau_i E\|\phi\|^2 \\ &+ p(E\|x_t\|^2) + E \int_0^t \|a(s, x_s)\|^2 ds + \sum_{0 < t_k < t} E I_k(x(t_k^-)) \\ &\leq E\|\phi\|^2 (1 + c_1 + \sum_{i=1}^p \tau_i) + 2c_2 + (c_1 + p)E\|x_t\|^2 + \int_0^t (c_1 E\|x_s\|^2 + c_2) ds \\ &+ \int_0^t q(s) \psi(E\|x_s\|^2) ds + m \sum_{0 < t_k < t} c_k^2 \\ &\leq \|\phi\|^2 (1 + c_1 + \sum_{i=1}^p \tau_i) + 2c_2 + (c_1 + p)\rho + (c_1 \rho + c_2) + \psi(\rho) \int_0^t q(s) ds \\ &+ m \sum_{0 < t_k < t} c_k^2 \\ &\leq E\|\phi\|^2 (1 + c_1 + \sum_{i=1}^p \tau_i) + 3c_2 + 2c_1 \rho + p\rho + \psi(\rho) \|q\| + m \sum_{0 < t_k < t} c_k^2 \\ &\leq \eta \text{ (say)} \end{aligned}$$

If $t \in J_0$, then $E\|\Gamma x\|^2 \leq \|\phi\|^2$ and hence $E\|\Gamma x\|^2 \leq \eta$ for all $x \in B$, that is Γ is bounded on bounded subsets of Ω .

Step 2: Γ maps bounded sets into equicontinuous sets.

Let B be as in Step 1 and $x \in B$. Let t and $h \neq 0$ be such that $t, t + h \in J \setminus \{t_1, t_2, \dots, t_m\}$

Now

$$\begin{aligned} E\|\Gamma x(t+h) - \Gamma x(t)\|^2 &\leq E\|f(t+h, x_{t+h}) - f(t, x_t)\|^2 + E \int_t^{t+h} \|g(s, x_s)\|^2 ds \\ &+ p h E\|x\|^2 + E \int_t^{t+h} \|a(s, x_s)\|^2 ds + \sum_{t < t_k < t+h} E\|I_k(x(t_k^-))\|^2 \\ &\leq E\|f(t+h, x_{t+h}) - f(t, x_t)\|^2 + E \int_t^{t+h} \|g(s, x_s)\|^2 ds \\ &+ p h \rho + E \int_t^{t+h} \|a(s, x_s)\|^2 ds + \sum_{t < t_k < t+h} E\|I_k(x(t_k^-))\|^2 \end{aligned}$$

as $h \rightarrow 0$, the right hand side of the above inequality tends to zero. This implies the equicontinuity on $J \setminus \{t_1, t_2, \dots, t_m\}$.

It remains to examine at $t=t_i, i=1, 2, \dots, m$. Let $h \neq 0$ be such that $\{t_k : k \neq i\} \cap [t_i - |h|, t_i + |h|] = \emptyset$ Thus we have

$$\|\Gamma x(t_i + h) - \Gamma x(t_i)\| \leq E\|f(t+h, x_{t+h}) - f(t, x_t)\|^2$$

$$+E \int_{t_i}^{t_i+h} (c_1 \|x_s\|^2 + c_2) ds + p \rho h + \psi(\rho) \int_{t_i}^{t_i+h} q(s) ds$$

The right hand side of the above inequality tends to zero as $h \rightarrow 0$. The equicontinuity on J_0 follows from the uniform continuity of ϕ on this interval.

Step 3: Now we show that Γ is continuous

Let $\{x_n\} \subset \Omega$ be a sequence such that $x_n \rightarrow x$. We will show that $\Gamma x_n \rightarrow \Gamma x$.

For $t \in J$

$$\begin{aligned} E \|\Gamma x_n - \Gamma x\|^2 &= E \left\| \int_0^t (f(t, x_n) - f(t, x)) + \int_0^t (g(s, x_n) - g(s, x)) ds + p \int_0^t (x_n(s) - x(s)) ds \right. \\ &+ \left. \int_0^t (a(s, x_n) - a(s, x)) dB(s) + \sum_{k=1}^m (I_k(x_n(t_k^-)) - I_k(x(t_k^-))) \right\|^2 \\ &\leq 3E \left\| \int_0^t (f(t, x_n) - f(t, x)) \right\|^2 + 3E \int_0^t \left\| (g(s, x_n) - g(s, x)) \right\|^2 ds \\ &+ 3pE \int_0^t \|x_n(s) - x(s)\|^2 ds + 3E \int_0^t \left\| (a(s, x_n) - a(s, x)) \right\|^2 ds \\ &+ 3m \sum_{k=1}^m E \left\| (I_k(x_n(t_k^-)) - I_k(x(t_k^-))) \right\|^2 \quad (3.2) \end{aligned}$$

Using H_3 it can be easily shown that the function $t \mapsto g(t, x_n) - g(t, x)$ is Lebesgue integrable. By the continuity of f and $I_k, k=1, 2, \dots, m$ and the dominated convergence theorem, the right hand side of inequality (3.2) tends to zero as $n \rightarrow \infty$, which completes the proof that Γ is continuous [15-18].

As a sequence of steps 1 to 3, together with the Arzela-Ascoli theorem, we conclude that Γ is completely continuous.

To complete the proof of the theorem, it suffices to prove the following step.

Step 4:

There exists a priori bound of the set

$$\zeta = \{x \in \Omega : \lambda x = \Gamma x \text{ for some } \lambda > 1\}$$

Let $x \in \zeta$ and $\lambda > 1$ be such that $\lambda x = \Gamma x$. Then $x|_{[-r, t_1]}$ satisfies for each $t \in [0, t_1]$,

$$\begin{aligned} x(t) &= \lambda^{-1} [\phi(0) - f(0, \phi(0)) + f(t, x_t) + \int_0^t g(s, x_s) ds + \int_0^t a(s, x_s) dB(s) \\ &+ \sum_{i=1}^p \int_{-\tau_i}^0 \phi(s) ds + \sum_{i=1}^p \int_0^{t-\tau_i} x(s) ds] \end{aligned}$$

It is easy to verify that

$$\begin{aligned} E \|x(t)\|^2 &\leq E \|\phi\|^2 \left(1 + c_1 + \sum_{i=1}^p \tau_i \right) + 2c_1 \|x_t\|^2 \\ &+ 3c_2 + \int_0^t (q(s) \psi(E \|x_s\|^2) + pE \|x_s\|^2) ds \quad (3.3) \end{aligned}$$

Consider the function $v_1(t) = \sup \{E \|x(s)\|^2 : s \in [-r, t]\}$, for $t \in [0, t_1]$. We have $\|x_t\|^2 \leq v_1(t)$ for all $t \in [0, t_1]$ and there is a point $t^* \in [-r, t]$ such that $v_1(t) = \|x_{t^*}\|^2$. If $t^* < 0$, we have $v_1(t) \leq \|\phi\|^2$ for all $t \in [0, t_1]$. Now, if $t^* \geq 0$ from (3.3) it follows that,

for $t \in [0, t_1]$,

$$v_1(t) \leq E \|\phi\|^2 \left(1 + c_1 + \sum_{i=1}^p \tau_i \right) + 2c_1 v_1(t) + 3c_2 + \int_0^t (q(s) \psi(v_1(s)) + p v_1(s)) ds$$

And hence

$$v_1(t) \leq C_1^1 + C_1^2 \int_0^t Q(s) [\psi(v_1(s)) + v_1(s)] ds$$

Where

$$C_1^1 = C_1^2 \left[\|\phi\|^2 \left(1 + c_1 + \sum_{i=1}^p \tau_i \right) + 3c_2 \right]$$

$$C_1^2 = \frac{1}{1 - 2c_1} \text{ And } Q(t) = \max \{q(t), p\}, \text{ for } t \in [0, t_1]$$

Set,

$$w_1(t) = C_1^1 + C_1^2 \int_0^t Q(s) [\psi(v_1(s)) + v_1(s)] ds \text{ for } t \in [0, t_1]$$

Then we have $v_1(t) \leq w_1(t)$ for all $t \in [0, t_1]$

A direct differentiation of w_1 yields

$$\begin{cases} w_1'(t) \leq Q(t) [\psi(w_1(t)) + w_1(t)], \text{ a.e. } t \in [0, t_1] \\ w_1(0) = C_1^1 \end{cases}$$

By integration, this gives

$$\int_0^t \frac{w_1(s)}{\psi(w_1(s)) + w_1(s)} ds \leq \int_0^t Q(s) ds \leq \|Q\| \quad t \in [0, t_1] \quad (3.4)$$

By a change of variables, inequality (3.4) becomes,

$$\int_{C_1^1}^{w_1(t)} \frac{ds}{\psi(s) + s} \leq \|Q\|, \quad t \in [0, t_1]$$

By (3.1) and the mean value theorem, there is a constant $M_1 = M_1(t) > 0$ such that $w_1(t) \leq M_1$ for all $t \in [0, t_1]$

That is $v_1(t) \leq M_1$ for all $t \in [0, t_1]$

At last we choose M_1 such that $\|\phi\| \leq M_1$ to get

$$\sup \{E \|x(t)\|^2 : t \in [-r, t_1]\} = v_1(t) \leq M_1$$

Now, consider $X|_{[-r, t_2]}$, satisfies for each $t \in [0, t_2]$

$$\begin{aligned} x(t) &= \lambda^{-1} [\phi(0) - f(0, \phi(0)) + f(t, x_t) + \int_0^t g(s, x_s) ds + \int_0^t a(s, x_s) dB(s) \\ &+ \sum_{i=1}^p \int_{-\tau_i}^0 \phi(s) ds + \sum_{i=1}^p \int_0^{t-\tau_i} x(s) ds + I_1(x(t_1))] \end{aligned}$$

$$\begin{aligned} E \|x(t)\|^2 &\leq E \|\phi\|^2 \left(1 + c_1 + \sum_{i=1}^p \tau_i \right) + 2c_1 \|x_t\|^2 + 3c_2 \\ &+ \int_0^t (q(s) \psi(E \|x_s\|^2) + pE \|x_s\|^2) ds + \sup E \|I_1(u)\|^2 : \|u\| \leq M_1 \end{aligned}$$

Denote $v_2(t) = \sup \{E \|x(s)\|^2 : s \in [-r, t]\}$, for $t \in [0, t_2]$. Then for each $t \in [0, t_2]$, we have $\|x(t)\|$, such that $\|x_t\|^2 \leq v_2(t)$.

Let $t^* \in [-r, t]$ be such that $v_2(t) = \|x_{t^*}\|^2$.

If $t^* < 0$, we have $v_2(t) \leq \|\phi\|^2$ for all $t \in [0, t_2]$.

Now if $t^* \geq 0$, then by (3.5) we have for $t \in [0, t_2]$

$$\begin{aligned} v_2(t) &\leq E \|\phi\|^2 \left(1 + c_1 + \sum_{i=1}^p \tau_i \right) + 2c_1 v_2(t) + 3c_2 + \int_0^t (q(s) \psi(v_2(s)) + p v_2(s)) \\ &+ \sup \{E \|I_1(u)\|^2 : \|u\| \leq M_1\} \end{aligned}$$

$$v_2(t) \leq C_2^1 + C_2^2 \int_0^t [Q(s) \psi(v_2(s)) + p v_2(s)]$$

Where

$$C_2^1 C_2^2 E \|\phi\|^2 \left(1 + c_1 + \sum_{i=1}^p \tau_i \right) + 3c_2 + \sup E \|I_1(u)\|^2 : \|u\| \leq M_1$$

$$C_2^2 = \frac{1}{1 - 2c_1}$$

And $Q(t) = \text{Max} \{q(t), p\}$, for $t \in [0, t_2]$.

If we set,

$$w_2(t) = C_2^1 + C_2^2 \int_0^t Q(s) [\psi(v_2(s)) + v_2(s)] ds \text{ for } t \in [0, t_2]$$

Then we get $v_2(t) \leq w_2(t)$ for all $t \in [0, t_2]$ and

$$\begin{cases} w_2(t) \leq Q(t) [\psi(w_2(t)) + w_2(t)], \text{ a.e. } t \in [0, t_2] \\ w_2(0) = C_2^1 \end{cases}$$

These yields,

$$\int_0^t \frac{w_2(s)}{\psi(w_2(s)) + pw_2(s)} ds \leq \int_0^t Q(s) ds \leq \|Q\|, \quad t \in [0, t_2]$$

$$\int_{C_2^1}^{w_2(t)} \frac{ds}{\psi(s) + s} \leq \|Q\|, \quad t \in [0, t_2]$$

Again by (3.1) and the mean value theorem, there is a constant $M_2 = M_2(t_1, t_2) > 0$ such that $w_2(t) \leq M_2$ for all $t \in [0, t_2]$, and then $v_2(t) \leq M_2$ for all $t \in [0, t_2]$. Finally, if we

Choose M_2 such that $\varphi \leq M_2$, we get,

$$\sup \{E \|x(t)\|^2 : t \in [-r, t_2]\} = v_2(t_2) \leq M_2$$

Continue this process for $x|_{[-r, t_3]}, \dots, x|_{J_1}$, we obtain that there exists a constant $M = M(t_1, \dots, t_m) > 0$ such that $x \leq M$.

This finish to show that the ζ is bounded in Ω .

As a result the conclusion of theorem holds and consequently the problem (1.1) has a solution x on J_1 . This completes the proof [19].

References

1. Anguraj A, Shujin Wu, Vinodkumar A (2011) The existence and exponential stability of semilinear functional differential equations with random impulses under non-uniqueness. *Nonlinear Analysis* 74: 331-342.
2. Wu S (2007) The Euler scheme for random impulsive differential equations. *Applied Mathematics and Computation* 191: 164-175.
3. Wu SJ, Han D (2005) exponential stability of functional differential systems with impulsive effect on random moments. *Computer and Mathematics with Applications* 50: 321-328.

4. Ren Y, Xia N (2009) A note on the existence and uniqueness of the solution to neutral stochastic functional differential equations with infinite delay. *applied mathematics and computation* 214: 457-461.
5. Sakthivel R, Luo J (2009) asymptotic stability of nonlinear impulsive stochastic differential equations. *Statistics and Probability Letters* 9: 1219-1223.
6. Anguraj A, Vinodkumar A (2009) Existence, Uniqueness and stability Results of Impulsive Stochastic Semi linear Neutral Functional Differential Equations with Infinite delays. *Electronic Journal of Qualitative Theory of Differential Equations* 67: 1-13.
7. Lakrib M (2008) An existence result for impulsive neutral functional differential equations with multiple delay. *Electronic Journal of Differential equations* 36: 1-7.
8. Oksendal B (2002) *Stochastic Differential Equations An Introduction with Applications: (5thedn)*, springer, Berlin, Germany.
9. Mao X (1997) *stochastic differential equations and their Applications*. Horwood Publishing, Chichester, UK.
10. Balachandran K, Kiruthika S (2010) Existence of solutions of Abstract fractional Impulsive Semi linear evolution Equations, *Electronic. Journal of Qualitative Theory of Differential Equations* 4: 1-12.
11. Sakthivel R, Luo J (2009) asymptotic stability of impulsive stochastic partial differential equations with infinite delays. *J Math Anal Appl* 356: 1-6.
12. Pan L (2012) Existence of mild solution for impulsive stochastic differential equations with non-local conditions. *Differential Equations and Applications* 4: 485-494
13. Guendouzi T, Mehdi K (2013) existence of mild solutions for impulsive fractional stochastic equations with infinite delay. *Malaya Journal of Matematik* 4: 30-43.
14. Anguraj A, Karthikeyan K (2009) Existence of solutions for impulsive neutral functional differential equations with nonlocal conditions. *Nonlinear Anal* 70: 2717-2721.
15. Yang Z, Xu D, Xiang L (2006) Exponential p-stability of impulsive stochastic differential equations with delays. *Physics Letters* 359: 129-137.
16. Benchora M, Henderson J, Ntouyas SK (2006) *Impulsive Differential equations and Inclusions*. Hindawi Publishing Corporation, New York, USA.
17. Bao H, Cao J (2009) Existence and Uniqueness of solutions to neutral stochastic functional differential equations with infinite delay. *Applied mathematics and Computation* 215: 1732-1743.
18. Sakthivel R, Luo J (2009) asymptotic stability of nonlinear impulsive stochastic differential equations. *Statistics and Probability Letters* 9: 1219-1223.
19. Hernandez EM, Rabello M, Hernan R, Henriquez HR (2007) existence of solutions for impulsive partial neutral functional differential equations. *J Math Anal Appl* 331: 1135-1158.

Citation: Anguraj A, Banupriya K (2015) An Existence Result for Impulsive Stochastic Functional Differential Equations with Multiple Delays. J Phys Math 6: 139. doi:10.4172/2090-0902.1000139

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 400 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://omicsonline.com/open-access/physical-mathematics.php>