

## HARD LEFSCHETZ THEOREM FOR SASAKIAN MANIFOLDS

BENIAMINO CAPPELLETTI-MONTANO, ANTONIO DE NICOLA  
& IVAN YUDIN

### Abstract

We prove that on a compact Sasakian manifold  $(M, \eta, g)$  of dimension  $2n + 1$ , for any  $0 \leq p \leq n$  the wedge product with  $\eta \wedge (d\eta)^p$  defines an isomorphism between the spaces of harmonic forms  $\Omega_{\Delta}^{n-p}(M)$  and  $\Omega_{\Delta}^{n+p+1}(M)$ . Therefore it induces an isomorphism between the de Rham cohomology spaces  $H^{n-p}(M)$  and  $H^{n+p+1}(M)$ . Such isomorphism is proven to be independent of the choice of a compatible Sasakian metric on a given contact manifold. As a consequence, an obstruction for a contact manifold to admit Sasakian structures is found.

### 1. Introduction

Sasakian manifolds, introduced by Sasaki [23] in 1960, can be described as an odd-dimensional counterpart of Kähler manifolds. Starting from the 90s, a renewed interest in Sasakian geometry was stimulated by new findings in theoretical physics (see e.g. [11, 16, 20, 21]), especially after the Maldacena conjecture [19] on the duality between conformal field theory and supergravity on anti-de-Sitter space-time. As a consequence, many important geometric and topological properties of Sasakian manifolds were discovered, see e.g. [3, 4, 10, 15].

It is well known that Sasakian geometry is naturally related to Kähler geometry from two sides: on the one hand Sasakian manifolds can be defined as those manifolds whose metric cone is Kähler, on the other hand the 1-dimensional foliation defined by the Reeb vector field is transversely Kähler.

A remarkable property of compact Kähler manifolds is given by the celebrated Hard Lefschetz Theorem, stating that the cup product with the suitable powers of the symplectic form gives isomorphisms between the de Rham cohomology groups of complementary degrees. This result was first stated by Lefschetz in [17] but the first complete proof was given by Hodge in [13]. One of the applications of the Hard Lefschetz

---

Received 06/11/2013.

Theorem is that it gives an obstruction for the existence of a Kähler metric on a compact symplectic manifold.

Later on, an odd dimensional version of the Hard Lefschetz Theorem was proven for compact coKähler manifolds in [6]. Thus one may ask whether a similar property also holds in the context of Sasakian geometry. So far, the only result that was obtained in this direction is the transversal Hard Lefschetz Theorem, proved by El Kacimi-Alaoui in [7], which holds for the basic cohomology with respect to any homologically orientable transversely Kähler foliation. In this paper, our aim is to prove that a version of the Hard Lefschetz Theorem holds for the de Rham cohomology of any compact Sasakian manifold.

Our approach needs to be different from the one adopted in Kähler geometry. Indeed, although Sasakian and Kähler manifolds share many properties, in this case the picture for Sasakian manifolds shows profound peculiarities. Let  $(M, \omega, g)$  be a compact Kähler manifold of dimension  $2n$  and  $\Omega_{\Delta}^p(M)$  the space of harmonic  $p$ -forms on  $M$ . We recall that the Hard Lefschetz Theorem states that for any  $0 \leq p \leq n$ , the maps

$$\begin{aligned}\omega^p \wedge - : \Omega_{\Delta}^{n-p}(M) &\rightarrow \Omega_{\Delta}^{n+p}(M) \\ \alpha &\mapsto \omega^p \wedge \alpha\end{aligned}$$

are isomorphisms. Now, let  $(M, \eta, g)$  be a compact Sasakian manifold of dimension  $2n + 1$ . As a natural generalization of the above Lefschetz isomorphism, one can consider, for each  $0 \leq p \leq n$ , the maps

$$\begin{aligned}\eta \wedge (d\eta)^p \wedge - : \Omega_{\Delta}^{n-p}(M) &\rightarrow \Omega_{\Delta}^{n+p+1}(M) \\ (1.1) \quad \alpha &\mapsto \eta \wedge (d\eta)^p \wedge \alpha.\end{aligned}$$

However, at this step a serious problem already arises. Namely, differently from the Kähler case, it is not true that the wedge multiplication by either  $d\eta$  or  $\eta \wedge d\eta$  maps harmonic forms into harmonic forms, so in principle the definition of the above maps may happen to be ill posed.

Nevertheless, we discover some spectral properties of the Laplace operator on differential forms which allow to overcome this obstacle. More precisely, given  $\alpha \in \Omega_{\Delta}^{n-p}(M)$ , we show that the forms  $\eta \wedge (d\eta)^k \wedge \alpha$  and  $(d\eta)^{k-1} \wedge \alpha$  are eigenforms of the Laplacian with positive integer eigenvalues for all  $0 \leq k \leq p-1$ . These eigenforms and their eigenvalues are visualized in Figure 1 at page 58 in the case  $n = 5$ . In the figure, the points on the horizontal axis represent the spaces of harmonic forms on  $M$ . All other points with coordinates  $(p, \nu)$  denote suitable subspaces of

$$\{\beta \in \Omega^p(M) \mid \Delta\beta = 4\nu\beta\}.$$

The segments represent the isomorphisms (3.1) and (3.7) between the corresponding vector spaces. As shown in the figure, the  $\Delta$ -eigenvalues

of  $\eta \wedge (d\eta)^k \wedge \alpha$  initially increase with  $k$  up to degree  $n$  and then decrease until they reach zero for  $k = p$ . Thus each mapping in (1.1) is actually well defined, in the sense that its target space is  $\Omega_\Delta^{n+p+1}(M)$ . Moreover, in Theorem 3.6 we prove that such map is in fact an isomorphism by explicitly giving its inverse map. This isomorphism obviously induces via Hodge theory an isomorphism between the corresponding de Rham cohomology groups. Namely, for each  $0 \leq p \leq n$  we obtain the isomorphism

$$\begin{aligned} Lef_{n-p}: H^{n-p}(M) &\rightarrow H^{n+p+1}(M) \\ [\beta] &\mapsto [\eta \wedge (d\eta)^p \wedge \Pi_\Delta \beta], \end{aligned}$$

where  $\Pi_\Delta \beta$  denotes the orthogonal projection of  $\beta$  on the space of harmonic forms. Note that, contrary to the symplectic case, we are forced to use the metric structure in the definition of  $Lef_p$ . Thus, *a priori*, one could expect that different Sasakian metrics on  $(M, \eta)$  could lead to different Lefschetz isomorphisms. To the utter surprise of the authors, this is not the case. In Theorem 4.5 we prove that the Lefschetz isomorphism is independent of the metric. This provides an obstruction for a contact manifold to admit Sasakian structures. In the last section, we introduce the notion of Lefschetz contact manifold and we prove that its odd Betti numbers up to the middle dimension are even.

**Acknowledgement.** Research partially supported by CMUC and FCT (Portugal), through European program COMPETE/FEDER, grants PEst-C/MAT/UI0324/2011, PTDC/MAT/099880/2008 and MTM2009-13383 (A.D.N.), SFRH/BPD/31788/2006 (I.Y.), Prin 2010/11 – Varietà reali e complesse: geometria, topologia e analisi armonica – Italy (B.C.M.)

## 2. Preliminaries

In this section we recall the definition of Sasakian manifolds and list some of their properties. For further details we refer the reader to [1, 2].

Let  $M$  be a smooth manifold of dimension  $2n + 1$ . A 1-form  $\eta$  on  $M$  is called a *contact form* if  $\eta \wedge (d\eta)^n$  nowhere vanishes. In this case the pair  $(M, \eta)$  is called a (strict) *contact manifold*. We write  $\Phi$  for  $\frac{1}{2}d\eta$  and we denote by  $\xi$  the *Reeb vector field* on  $(M, \eta)$ , that is the unique vector field on  $M$  such that  $i_\xi \eta = 1$  and  $i_\xi d\eta = 0$ .

Let  $(M, \eta)$  be a contact manifold and  $g$  a Riemannian metric on  $M$ . We define the endomorphism  $\phi: TM \rightarrow TM$  by  $\Phi(X, Y) = g(X, \phi Y)$ .

Then  $(M, \eta, g)$  is called a *Sasakian manifold* if the following conditions hold:

- (i)  $\phi^2 = -I + \eta \otimes \xi$ , where  $I$  is the identity operator;
- (ii)  $g(\phi X, \phi Y) = g(X, Y) - \eta(X)\eta(Y)$  for any vector fields  $X$  and  $Y$  on  $M$ ;

(iii) the normality condition is satisfied, namely

$$[\phi, \phi]_{FN} + 2d\eta \otimes \xi = 0,$$

where  $[-, -]_{FN}$  is the Frölicher-Nijenhuis bracket (cf. [14]).

It is well known that in any Sasakian manifold one has

$$(2.1) \quad \phi\xi = 0, \quad \eta \circ \phi = 0.$$

Now we will introduce notation for some linear operators on the de Rham algebra  $\Omega^*(M)$ . For a  $p$ -form  $\alpha$  on  $M$ , we denote by  $\epsilon_\alpha$  the operator given by

$$\epsilon_\alpha \beta = \alpha \wedge \beta,$$

where  $\beta \in \Omega^*(M)$ . If  $(M, \eta)$  is a compact contact manifold, then  $\epsilon_\eta$  is adjoint to  $i_\xi$  with respect to the usual global scalar product on  $\Omega^*(M)$ , that is  $\epsilon_\eta = i_\xi^*$ . Since  $i_\xi \eta = 1$ , for any  $\omega \in \Omega^*(M)$ , we get

$$(2.2) \quad \{i_\xi, \epsilon_\eta\} \omega = \omega,$$

where the curly brackets are used to denote the anti-commutator of two operators.

Let  $(M, \eta, g)$  be a compact Sasakian manifold. We define the operators  $L$  and  $\Lambda$  on  $\Omega^*(M)$  by

$$L = \epsilon_\Phi, \quad \Lambda = L^*.$$

Then, since  $d$  is a graded derivation on  $\Omega^*(M)$  of degree 1 and  $\eta$  is a 1-form, we get

$$(2.3) \quad \{d, \epsilon_\eta\} = \epsilon_{d\eta} = 2\epsilon_\Phi = 2L.$$

Therefore

$$(2.4) \quad \{i_\xi, \delta\} = \{d, \epsilon_\eta\}^* = 2L^* = 2\Lambda.$$

Hereafter, we will use some elements of Frölicher-Nijenhuis calculus, developed in [8] (see also [14, Section 8]). In this framework, for every vector valued  $(k+1)$ -form  $\psi$ , the graded derivation  $i_\psi$  of degree  $k$  on  $\Omega^*(M)$  is defined. It acts on  $\omega \in \Omega^p(M)$  by

$$(i_\psi \omega)(X_1, \dots, X_{p+k}) = \sum_{\sigma} (-1)^{\sigma} \omega(\psi(X_{\sigma(1)}, \dots, X_{\sigma(k+1)}), \dots, X_{\sigma(p+k)}),$$

where the summation is taken over all  $(k+1, p-1)$ -shuffles. In the case  $k+1=0$ , that is when  $\psi=X$  is a vector field, we reobtain the usual interior product  $i_X$ . If  $\psi$  is an endomorphism of  $TM$ , that is when  $k=0$ , the formula above can be rewritten as

$$i_\psi \omega(X_1, \dots, X_p) = \sum_{s=1}^p \omega(X_1, \dots, \psi X_s, \dots, X_p).$$

For any vector valued  $(k+1)$ -form  $\psi$  the operator  $\mathcal{L}_\psi$  is defined to be  $i_\psi d - (-1)^k d i_\psi$ . Note that every  $\mathcal{L}_\psi$  is a graded derivation of degree

$k + 1$  on  $\Omega^*(M)$ . If  $\psi = X$  is a vector field, then  $\mathcal{L}_X$  is the usual Lie derivative.

When  $\psi$  is the identity operator  $I: TM \rightarrow TM$ , the operator  $i_\psi$  will be denoted by  $\deg$ , motivated by the fact that for any  $p$ -form  $\omega$  we have

$$\deg \omega = i_I \omega = p\omega.$$

Now we summarize several results of Fujitani [9] on commutators between operators on  $\Omega^*(M)$ , where  $M$  is a Sasakian manifold (N.B.: Fujitani uses  $\Phi$  for  $i_\phi$ ,  $\varphi$  for  $\Phi$ ,  $\lambda$  for  $i_\xi$ , and  $l$  for  $\epsilon_\eta$ ).

**Theorem 2.1.** *Let  $M$  be a compact Sasakian manifold of dimension  $2n+1$ . Then the operator  $i_\phi$  commutes with  $L$  and  $\Lambda$ . The Lie derivative  $\mathcal{L}_\xi$  commutes with  $d$ ,  $\delta$ ,  $\epsilon_\eta$ ,  $i_\xi$ ,  $L$ ,  $\Lambda$ , and  $i_\phi$ . Furthermore*

$$(2.5) \quad [d, \Lambda] = [i_\phi, \delta] - 2(n - \deg) i_\xi,$$

$$(2.6) \quad [\Delta, i_\xi] = 2[i_\phi, \delta] - 4(n - \deg) i_\xi,$$

$$(2.7) \quad [\Delta, i_\phi] = -2(\mathcal{L}_\xi - i_\xi d + \epsilon_\eta \delta),$$

$$(2.8) \quad [\Delta, \epsilon_\eta] = -2\mathcal{L}_\phi + 4\epsilon_\eta (n - \deg).$$

*Proof.* See Propositions 1.1, 1.2, 3.3 and Theorem 3.2 in [9]. q.e.d.

Using the equalities (2.5)–(2.8), Fujitani reobtained a few results of Tachibana [24] in [9, Theorem 4.1] and complemented them with the dual ones in [9, Corollary 4.2]. Below is the summary of these results.

**Proposition 2.2.** *Let  $M$  be a compact Sasakian manifold of dimension  $2n+1$  and  $\omega$  a harmonic  $p$ -form on  $M$ . Then  $i_\phi \omega$  is also harmonic and, moreover,*

- (i) if  $p \leq n$ , then  $i_\xi \omega = 0$ ;
- (ii) if  $p \geq n+1$ , then  $\epsilon_\eta \omega = 0$ ;
- (iii) if  $p \leq n+1$ , then  $\Lambda \omega = 0$ ;
- (iv) if  $p \geq n$ , then  $L \omega = 0$ .

In the following proposition we prove some other useful identities.

**Proposition 2.3.** *In any Sasakian manifold we have*

$$(2.9) \quad [i_\phi, \epsilon_\eta] = 0,$$

$$(2.10) \quad [i_\phi, i_\xi] = 0,$$

$$(2.11) \quad \mathcal{L}_\phi^2 = -2L\mathcal{L}_\xi,$$

$$(2.12) \quad \{\delta, \epsilon_\eta\} = -\mathcal{L}_\xi.$$

*Proof.* Since  $i_\phi$  is a derivation of degree zero, we have  $[i_\phi, \epsilon_\eta] = \epsilon_{i_\phi \eta} = 0$ , as  $i_\phi \eta = 0$  by (2.1). Next, it is easy to check that for any  $\psi: TM \rightarrow TM$  and any vector field  $X$  we have  $[i_\psi, i_X] = -i_{\psi X}$ . Thus  $[i_\phi, i_\xi]$  is zero by (2.1). Further, according to Frölicher-Nijenhuis calculus, we have

$\{\mathcal{L}_\phi, \mathcal{L}_\phi\} = \mathcal{L}_{[\phi, \phi]_{FN}}$ . Thus, from the normality condition for Sasakian manifolds, we get

$$\begin{aligned}\mathcal{L}_\phi^2 &= \frac{1}{2}\mathcal{L}_{[\phi, \phi]_{FN}} = -\mathcal{L}_{d\eta \otimes \xi} = -\{i_{d\eta \otimes \xi}, d\} = -\{d\eta \wedge i_\xi, d\} \\ &= -d\eta \wedge \mathcal{L}_\xi = -2L\mathcal{L}_\xi.\end{aligned}$$

Finally, it was shown on page 109 of [12] that, for any Killing vector field  $X$ , one has  $\mathcal{L}_X + \{\delta, \epsilon_{g(X, -)}\} = 0$ . Since in any Sasakian manifold the Reeb vector field  $\xi$  is Killing and  $\eta = g(\xi, -)$ , the equation (2.12) holds. q.e.d.

### 3. Hard Lefschetz isomorphism for harmonic forms

In this section we establish the Hard Lefschetz isomorphism between the spaces of harmonic  $p$ -forms and harmonic  $(2n+1-p)$ -forms in a compact Sasakian manifold  $M$  of dimension  $2n+1$ .

We start by introducing some notation. Let  $A_1, \dots, A_k$  be operators on  $\Omega^*(M)$ . We will denote by  $\Omega_{A_1, \dots, A_k}^p(M)$  the set of  $p$ -forms  $\omega$  such that  $A_1\omega = \dots = A_k\omega = 0$  and we will use  $\Omega_{A_1, \dots, A_k}^{p,\nu}(M)$  as a shorthand for  $\Omega_{\Delta-\nu I, A_1, \dots, A_k}^p(M)$ , where  $\nu$  is a real number. Of course these spaces are empty if  $\nu < 0$  and  $M$  is compact, since all  $\Delta$ -eigenvalues are non-negative in this case.

We will be mainly concerned with two families of spaces of differential forms on a compact Sasakian manifold, namely,  $\Omega_{d, i_\xi, \epsilon_{\eta\delta}}^{p,\nu}(M)$  and  $\Omega_{\delta, \epsilon_\eta, i_\xi d}^{p,\nu}(M)$ , where  $0 \leq p \leq 2n+1$  and  $\nu$  are non-negative real numbers. It follows from (2.12) that the spaces of both families are included in  $\Omega_{\mathcal{L}_\xi}^*(M)$ . Let us show that these two families are related to each other by the Hodge star operator  $*$ .

**Proposition 3.1.** *Let  $M$  be a compact Sasakian manifold of dimension  $2n+1$ . Then for any  $p$ -form  $\omega$ , we have  $\omega \in \Omega_{d, i_\xi, \epsilon_{\eta\delta}}^{p,\nu}(M)$  if and only if  $*\omega \in \Omega_{\delta, \epsilon_\eta, i_\xi d}^{2n+1-p,\nu}(M)$ .*

*Proof.* It is easy to check that for any  $p$ -form  $\omega$ , one has  $\delta * \omega = (-1)^{p+1} * d\omega$  and  $\epsilon_\eta * \omega = (-1)^{p+1} * i_\xi \omega$ . Hence  $i_\xi d * \omega = -* \epsilon_\eta \delta \omega$ . Moreover,  $\Delta * \omega = *\Delta \omega$ . q.e.d.

Now we show that the spaces of harmonic forms on a compact Sasakian manifold are included in the above families of subspaces.

**Proposition 3.2.** *Let  $M$  be a compact Sasakian manifold of dimension  $2n+1$ .*

(i) *For  $p \leq n$ , we have*

$$\Omega_{\Delta}^p(M) = \Omega_{d, i_\xi, \epsilon_{\eta\delta}}^{p,0}(M) \text{ and } \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p,0}(M) = 0.$$

(ii) For  $p \geq n + 1$ , we have

$$\Omega_{\Delta}^p(M) = \Omega_{\delta, \epsilon_{\eta}, i_{\xi}d}^{p,0}(M) \text{ and } \Omega_{d, i_{\xi}, \epsilon_{\eta}\delta}^{p,0}(M) = 0.$$

*Proof.* It is obvious that  $\Omega_{d, i_{\xi}, \epsilon_{\eta}\delta}^{p,0}(M) \subset \Omega_{\Delta}^p(M)$ . Let  $\omega \in \Omega_{\Delta}^p(M)$  with  $p \leq n$ . Since  $\omega$  is harmonic, we have  $d\omega = 0$  and  $\epsilon_{\eta}\delta\omega = 0$ . Moreover, by Proposition 2.2 we have that  $i_{\xi}\omega = 0$ , since  $p \leq n$ . Thus  $\Omega_{d, i_{\xi}, \epsilon_{\eta}\delta}^{p,0}(M) = \Omega_{\Delta}^p(M)$ .

In order to prove that  $\Omega_{\delta, \epsilon_{\eta}, i_{\xi}d}^{p,0}(M) = 0$ , notice that

$$\Omega_{\delta, \epsilon_{\eta}, i_{\xi}d}^{p,0}(M) \subset \Omega_{\Delta}^p(M) = \Omega_{d, i_{\xi}, \epsilon_{\eta}\delta}^{p,0}(M)$$

implies  $\epsilon_{\eta}\omega = 0$  and  $i_{\xi}\omega = 0$  for any  $\omega \in \Omega_{\delta, \epsilon_{\eta}, i_{\xi}d}^{p,0}(M)$ . Therefore using (2.2), we get

$$\omega = i_{\xi}\epsilon_{\eta}\omega + \epsilon_{\eta}i_{\xi}\omega = 0.$$

Hence  $\Omega_{\delta, \epsilon_{\eta}, i_{\xi}d}^{p,0}(M) = 0$ . The second part of the proposition can be proved by duality considerations, using Proposition 3.1. q.e.d.

**Proposition 3.3.** *Let  $M$  be a compact Sasakian manifold.*

- (i) *If  $\omega \in \Omega_{\delta, \epsilon_{\eta}, i_{\xi}d}^{p,4\nu}(M)$  then  $d\omega \in \Omega_{d, i_{\xi}, \epsilon_{\eta}\delta}^{p+1,4\nu}(M)$ . If moreover  $\nu \neq 0$ , then  $d\omega \neq 0$ .*
- (ii) *If  $\omega \in \Omega_{d, i_{\xi}, \epsilon_{\eta}\delta}^{p,4\nu}(M)$ , then  $\delta\omega \in \Omega_{\delta, \epsilon_{\eta}, i_{\xi}d}^{p-1,4\nu}(M)$ . If moreover  $\nu \neq 0$ , then  $\delta\omega \neq 0$ .*

*Proof.* By the duality property established in Proposition 3.1, it is enough to prove just (i). Let  $\omega \in \Omega_{\delta, \epsilon_{\eta}, i_{\xi}d}^{p,4\nu}(M)$ . Then

$$d(d\omega) = 0, \quad i_{\xi}d\omega = 0, \quad \Delta d\omega = d\Delta\omega = 4\nu d\omega.$$

It is left to show that  $\epsilon_{\eta}\delta d\omega = 0$ . Since  $\delta\omega = 0$  and  $\epsilon_{\eta}\omega = 0$ , we get

$$\epsilon_{\eta}\delta d\omega = \epsilon_{\eta}\delta d\omega + \epsilon_{\eta}d\delta\omega = \epsilon_{\eta}\Delta\omega = 4\nu\epsilon_{\eta}\omega = 0.$$

If moreover  $\nu \neq 0$ , then  $\delta d\omega = \Delta\omega = 4\nu\omega \neq 0$ . Thus also  $d\omega \neq 0$ . q.e.d.

Proposition 3.3 shows that for  $\nu \neq 0$  we have the pair of isomorphisms

$$(3.1) \quad \Omega_{\delta, \epsilon_{\eta}, i_{\xi}d}^{p,4\nu}(M) \xrightleftharpoons[\delta]{d} \Omega_{d, i_{\xi}, \epsilon_{\eta}\delta}^{p+1,4\nu}(M),$$

for any  $0 \leq p \leq 2n$ .

**Theorem 3.4.** *Let  $M$  be a compact Sasakian manifold of dimension  $2n + 1$ .*

- (i) *If  $\omega \in \Omega_{\delta, \epsilon_{\eta}, i_{\xi}d}^{p,4\nu}(M)$  then  $i_{\xi}\omega \in \Omega_{d, i_{\xi}, \epsilon_{\eta}\delta}^{p-1,4(\nu+p-n-1)}(M)$ .*
- (ii) *If  $\omega \in \Omega_{d, i_{\xi}, \epsilon_{\eta}\delta}^{p,4\nu}(M)$  then  $\epsilon_{\eta}\omega \in \Omega_{\delta, \epsilon_{\eta}, i_{\xi}d}^{p+1,4(\nu-p+n)}(M)$ .*

*Proof.* We will prove just (i) as (ii) can be obtained from (i) by using Hodge duality and Proposition 3.1. Let  $\omega \in \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p, 4\nu}(M)$ . We write  $\nu'$  for  $\nu + p - n - 1$ . We have to show that  $i_\xi \omega \in \Omega_{d, i_\xi, \epsilon_\eta \delta}^{p-1, 4\nu'}(M)$ . First of all, from (2.12), we get

$$(3.2) \quad di_\xi \omega = \mathcal{L}_\xi \omega - i_\xi d\omega = -\{\epsilon_\eta, \delta\} \omega = 0.$$

Next, it is obvious that  $i_\xi i_\xi \omega = 0$ . Moreover, by using (2.12), (2.2) and (3.2) we get

$$\epsilon_\eta \delta i_\xi \omega = -\mathcal{L}_\xi i_\xi \omega - \delta \epsilon_\eta i_\xi \omega = -i_\xi di_\xi \omega - \delta(\omega - i_\xi \epsilon_\eta \omega) = 0.$$

Thus  $i_\xi \omega \in \Omega_{d, i_\xi, \epsilon_\eta \delta}^{p-1}(M)$ . It is left to prove that  $\Delta i_\xi \omega = 4\nu' i_\xi \omega$ . From the equation (2.6) for  $[\Delta, i_\xi]$ , using that  $\delta \omega = 0$ , we get

$$\Delta i_\xi \omega - i_\xi \Delta \omega = -2\delta i_\phi \omega - 4(n-p+1)i_\xi \omega.$$

Since  $\Delta \omega = 4\nu \omega$  and  $\nu - n + p - 1 = \nu'$ , we get

$$(3.3) \quad \Delta i_\xi \omega = 4\nu' i_\xi \omega - 2\delta i_\phi \omega.$$

To get that  $\delta i_\phi \omega = 0$ , we proceed as follows. First we apply  $d\delta$  to (3.3). Using that  $d\delta$  commutes with  $\Delta$ , we get

$$\Delta d\delta i_\xi \omega = 4\nu' d\delta i_\xi \omega.$$

As  $di_\xi \omega = 0$  by (3.2), we have  $d\delta i_\xi \omega = \Delta i_\xi \omega$ . Therefore

$$(3.4) \quad \Delta \Delta i_\xi \omega = 4\nu' \Delta i_\xi \omega.$$

Comparing (3.3) and (3.4), we see that  $\Delta \delta i_\phi \omega = 0$ , thence also

$$(3.5) \quad \delta \Delta i_\phi \omega = 0.$$

Further, using the formula (2.7) for  $[\Delta, i_\phi]$ , we get

$$(3.6) \quad \Delta i_\phi \omega = i_\phi \Delta \omega - 2\mathcal{L}_\xi \omega + 2i_\xi d\omega - 2\epsilon_\eta \delta \omega = 4\nu i_\phi \omega,$$

since  $\omega \in \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p, 4\nu}(M)$  and hence  $\mathcal{L}_\xi \omega = 0$ . From (3.5) and (3.6) it follows that  $4\nu \delta i_\phi \omega = \delta \Delta i_\phi \omega = 0$ . If  $\nu \neq 0$ , this implies that  $\delta i_\phi \omega = 0$ . On the other hand if  $\nu = 0$ , then  $\omega$  is harmonic. Thus by Proposition 2.2 the form  $i_\phi \omega$  is also harmonic. Hence  $\delta i_\phi \omega = 0$ . Therefore from (3.3) we finally get that  $\Delta i_\xi \omega = 4\nu' i_\xi \omega$ . q.e.d.

Since  $\{\epsilon_\eta, i_\xi\} = I$ , we get from Theorem 3.4 that  $\epsilon_\eta$  and  $i_\xi$  induce the pair of inverse isomorphisms

$$(3.7) \quad \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p, 4\nu}(M) \xrightleftharpoons[\epsilon_\eta]{i_\xi} \Omega_{d, i_\xi, \epsilon_\eta \delta}^{p-1, 4\nu'}(M),$$

where  $\nu' = \nu + p - n - 1$  and  $1 \leq p \leq 2n + 1$ . In fact, if  $\omega$  is on the left hand side, then  $\epsilon_\eta \omega = 0$  and  $\epsilon_\eta i_\xi \omega = \omega - i_\xi \epsilon_\eta \omega = \omega$ . Similarly, if  $\omega$  is on the right hand side, then  $i_\xi \omega = 0$  and  $i_\xi \epsilon_\eta \omega = \omega$ .

**Corollary 3.5.** *Let  $M$  be a compact Sasakian manifold of dimension  $2n + 1$ . Then for  $\nu \neq 0$  and  $p \leq 2n - 1$*

$$(3.8) \quad \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p, 4\nu}(M) \xrightleftharpoons[\Lambda]{L} \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p+2, 4(\nu-p-1+n)}(M),$$

*is a pair of isomorphisms such that  $\Lambda L = \nu I$  and  $L\Lambda = \nu I$ . Moreover, for  $\nu \neq p - n$*

$$(3.9) \quad \Omega_{d, i_\xi, \epsilon_\eta \delta}^{p, 4\nu}(M) \xrightleftharpoons[\Lambda]{L} \Omega_{d, i_\xi, \epsilon_\eta \delta}^{p+2, 4(\nu-p+n)}(M),$$

*is a pair of isomorphisms such that  $L\Lambda = (\nu - p + n)I$  and  $\Lambda L = (\nu - p + n)I$ .*

*Proof.* Notice that from (2.3), we have that for every  $\omega \in \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p, 4\nu}(M)$

$$(3.10) \quad 2L\omega = \epsilon_\eta d\omega.$$

Similarly from (2.4), for every  $\omega \in \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p+2, 4(\nu-p-1+n)}(M)$ , we get

$$(3.11) \quad 2\Lambda\omega = \delta i_\xi \omega.$$

Therefore, using the isomorphisms (3.1) and (3.7), we can construct the diagram

$$\begin{array}{ccccc} & & 2L & & \\ & \swarrow & & \searrow & \\ \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p, 4\nu}(M) & \xrightleftharpoons[\delta]{d} & \Omega_{d, i_\xi, \epsilon_\eta \delta}^{p+1, 4\nu}(M) & \xrightleftharpoons[i_\xi]{\epsilon_\eta} & \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p+2, 4(\nu-p-1+n)}(M). \\ & \searrow & & \swarrow & \\ & & 2\Lambda & & \end{array}$$

This shows that  $L$  and  $\Lambda$  induce isomorphisms between the spaces in (3.8). For every  $\alpha \in \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p, 4\nu}(M)$  and every  $\beta \in \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p+2, 4(\nu-p-1+n)}(M)$ , we have

$$\begin{aligned} \Lambda L\alpha &= \frac{1}{4}\delta i_\xi \epsilon_\eta d\alpha = \frac{1}{4}\delta d\alpha = \frac{1}{4}\Delta\alpha = \nu\alpha, \\ L\Lambda\beta &= \frac{1}{4}\epsilon_\eta d\delta i_\xi \beta = \frac{1}{4}\epsilon_\eta \Delta i_\xi \beta = \nu\epsilon_\eta i_\xi \beta = \nu\beta. \end{aligned}$$

The second part of the corollary is proved by the same line of reasoning from the diagram

$$\begin{array}{ccccc} & & 2L & & \\ & \swarrow & & \searrow & \\ \Omega_{d, i_\xi, \epsilon_\eta \delta}^{p, 4\nu}(M) & \xrightleftharpoons[i_\xi]{\epsilon_\eta} & \Omega_{\delta, \epsilon_\eta, i_\xi d}^{p+1, 4(\nu-p+n)}(M) & \xrightleftharpoons[\delta]{d} & \Omega_{d, i_\xi, \epsilon_\eta \delta}^{p+2, 4(\nu-p+n)}(M). \\ & \searrow & & \swarrow & \\ & & 2\Lambda & & \end{array}$$

q.e.d.

Now, we are prepared to prove the Hard Lefschetz Theorem at the level of harmonic forms.

**Theorem 3.6.** *Let  $M$  a compact Sasakian manifold of dimension  $2n+1$  and  $p \leq n$ . Then the map*

$$\epsilon_\eta L^{n-p}: \Omega^p(M) \rightarrow \Omega^{2n+1-p}(M)$$

*induces an isomorphism  $F_p: \Omega_\Delta^p(M) \rightarrow \Omega_\Delta^{2n+1-p}(M)$ . Similarly, the map*

$$\Lambda^{n-p} i_\xi: \Omega^{2n+1-p}(M) \rightarrow \Omega^p(M)$$

*induces an isomorphism  $G_p: \Omega_\Delta^{2n+1-p}(M) \rightarrow \Omega_\Delta^p(M)$ . Moreover,*

$$F_p G_p = (n-p)!^2 \cdot I, \quad G_p F_p = (n-p)!^2 \cdot I.$$

*Proof.* We have by Proposition 3.2 that  $\Omega_\Delta^p(M) = \Omega_{d,i_\xi,\epsilon_\eta\delta}^{p,0}(M)$  and  $\Omega_\Delta^{2n+1-p}(M) = \Omega_{\delta,\epsilon_\eta,i_\xi d}^{2n+1-p,0}(M)$ . Let us define the numbers  $\nu_{p,k}$  by

$$(3.12) \quad \nu_{p,k} := k(n-p-k+1), \quad k \in \mathbb{Z}.$$

To make the notation less heavy, we will write  $\nu_k$  instead of  $\nu_{p,k}$  along the proof. It is easy to check that  $\nu_{k+1} = \nu_k - (p+2k) + n$ , and that  $\nu_k \neq 0$  for  $1 \leq k \leq n-p$ . Thus by Corollary 3.5, the operators  $L$  and  $\Lambda$  induce isomorphisms between  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{p+2k,4\nu_k}(M)$  and  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{p+2(k+1),4\nu_{k+1}}(M)$ .

Moreover, for  $\omega \in \Omega_{d,i_\xi,\epsilon_\eta\delta}^{p+2k,4\nu_k}(M)$ , it holds

$$(3.13) \quad \Lambda L \omega = \nu_{k+1} \omega.$$

Note that  $\nu_0 = 0$  and  $\nu_{n-p} = n-p$ . Then we get the chain of isomorphisms

$$(3.14)$$

$$\Omega_{d,i_\xi,\epsilon_\eta\delta}^{p,0}(M) \xrightarrow{L} \Omega_{d,i_\xi,\epsilon_\eta\delta}^{p+2,4\nu_1}(M) \xrightarrow{L} \dots \xrightarrow{L} \Omega_{d,i_\xi,\epsilon_\eta\delta}^{2n-p,4(n-p)}(M).$$

Thus  $L^{n-p}$  induces an isomorphism between the vector spaces  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{p,0}(M)$  and  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{2n-p,4(n-p)}(M)$ . From (3.7), we have that  $\epsilon_\eta$  and  $i_\xi$  induce two mutually inverse maps between  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{2n-p,4(n-p)}(M)$  and  $\Omega_{\delta,\epsilon_\eta,i_\xi d}^{2n-p+1,0}(M) = \Omega_\Delta^{2n-p+1}(M)$ . This shows, that  $F_p$  is an isomorphism between  $\Omega_\Delta^p(M)$  and  $\Omega_\Delta^{2n-p+1}(M)$ . Similarly, one can check that also  $G_p: \Omega_\Delta^{2n-p+1}(M) \rightarrow \Omega_\Delta^p(M)$  is an isomorphism. Iterating (3.13), we see that for all  $\omega \in \Omega_\Delta^p(M)$

$$\begin{aligned} G_p F_p \omega &= \Lambda^{n-p} L^{n-p} \omega = \left( \prod_{k=1}^{n-p} \nu_k \right) \omega = \left( \prod_{k=1}^{n-p} k(n-p-k+1) \right) \omega \\ &= (n-p)!^2 \omega. \end{aligned}$$

Similarly, for  $\omega \in \Omega_\Delta^{2n-p+1}(M)$ , we have  $F_p G_p \omega = (n-p)!^2 \omega$ . q.e.d.

From (3.14) it follows that the spaces  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{p+2k,4\nu_{p,k}}(M)$ , where  $\nu_{p,k}$  are defined by (3.12), are isomorphic to  $\Omega_\Delta^p(M)$  for all  $1 \leq k \leq n-p$ . Thus, if  $H^p(M) \neq 0$ , the same is true for  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{p+2k,4\nu_{p,k}}(M)$ . The following proposition shows that all the pairs  $(p, \nu)$  such that  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{p,4\nu}(M) \neq 0$  are necessarily of the form  $(q+2k, \nu_{q,k})$ .

**Proposition 3.7.** *Let  $M$  be a compact Sasakian manifold of dimension  $2n+1$ .*

(i) *If  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{p,4\nu}(M) \neq 0$  then  $\nu = \nu_{p-2k,k}$  for some integer  $k$  such that*

$$\max\{0, (p-n)/2\} \leq k \leq p/2.$$

(ii) *If  $\Omega_{\delta,\epsilon_\eta,i_\xi d}^{p,4\nu}(M) \neq 0$  then  $\nu = \nu_{p+1-2k,k}$  for some integer  $k$  such that*

$$\max\{0, (p+1-n)/2\} \leq k \leq (p+1)/2.$$

*Proof.* Note that the second part of the proposition follows from the first part and the isomorphism (3.1). We will prove the first part by induction on  $p$ . Suppose  $p=0$  and  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{0,4\nu}(M) \neq 0$ . Let  $f \in \Omega_{d,i_\xi,\epsilon_\eta\delta}^{0,4\nu}(M)$ ,  $f \neq 0$ . Then  $df=0$  and thus  $f$  is a harmonic function. Since  $4\nu$  is the  $\Delta$ -eigenvalue of  $f$ , we get that necessarily  $\nu=0$ . Thus  $\nu=\nu_{0,0}$ .

Suppose we proved the claim for all  $p' < p$  and  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{p',4\nu}(M) \neq 0$ . If  $\nu=0$ , then by Proposition 3.2 we get that  $p \leq n$ . Therefore 0 is in the allowed range of values for  $k$ , and we are done since  $\nu_{p,0}=0$ . Now consider the case  $\nu \neq 0$ . From Corollary 3.5, we get that the spaces  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{p-2,4(\nu+p-2-n)}(M)$  and  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{p,4\nu}(M)$  are isomorphic. From the induction assumption, we get that

$$\nu + p - 2 - n = \nu_{(p-2)-2k',k'}$$

for some  $k'$  between  $\max\{0, (p-2-n)/2\}$  and  $(p-2)/2$ . Define  $k=k'+1$ . Then

$$\begin{aligned} \nu &= \nu_{p-2k,k-1} - p + 2 + n \\ &= (k-1)(n-p+k+2) + n - p + 2 \\ &= k(n-p+k+1) \\ &= \nu_{p-2k,k}. \end{aligned}$$

Moreover  $k$  lies between  $\max\{1, (p-n)/2\}$  and  $p/2$ . This completes the induction argument. q.e.d.

In Figure 1, we give an illustration of what happens for a compact Sasakian manifold  $M$  of dimension 11. In the picture the circles mark the places  $(p, \nu)$  such that  $\Omega_{\delta,\epsilon_\eta,i_\xi d}^{p,4\nu}(M)$  can be non-zero. Similarly, the squares mark the points  $(p, \nu)$  such that the spaces  $\Omega_{d,i_\xi,\epsilon_\eta\delta}^{p,4\nu}(M)$  can be non-zero. The horizontal segments with a circle at the left edge




Figure 1

and a square at the right edge represent the isomorphisms of the type (3.1), and all the other segments correspond to the isomorphisms of the type (3.7). Thus, we can see that if we start with a harmonic  $p$ -form  $\omega$  for  $p \leq n$  and move it along the segments representing the isomorphisms of the types (3.1) and (3.7), we get in the intermediate steps  $\Delta$ -eigenforms with non-zero eigenvalues, and we will eventually end up with the harmonic  $(2n + 1 - p)$ -form  $F_p\omega$ .

#### 4. Hard Lefschetz isomorphism in de Rham cohomology

Let  $M$  be a compact Sasakian manifold of dimension  $2n + 1$ . Let us denote by  $\Pi_\Delta$  the orthogonal projection from  $\Omega^*(M)$  onto  $\Omega_\Delta^*(M)$ . Then we can define

$$(4.1) \quad \begin{aligned} Lef_p: H^p(M) &\rightarrow H^{2n+1-p}(M) \\ [\beta] &\mapsto [F_p \Pi_\Delta \beta]. \end{aligned}$$

Due to Hodge theory and Theorem 3.6 the map  $Lef_p$  is an isomorphism.

Suppose  $(\eta, g')$  is another Sasakian structure on  $M$  with the same contact form  $\eta$ . Denote by  $\Delta'$  the corresponding Laplacian. For a closed  $p$ -form  $\beta$  it can happen that

$$\Pi_{\Delta}\beta \neq \Pi_{\Delta'}\beta.$$

Thus it is not *a priori* clear whether  $Lef_p$  depends on the full Sasakian structure or just on its contact form. Note that this is rather different from the Kähler case, where it is obvious from the definition of the Lefschetz map that it is fully determined by the symplectic structure and does not depend on chosen Kähler metric.

The aim of this section is to show that  $Lef_p$  is uniquely determined by the contact structure of  $M$ . Note that from Proposition 2.2 it follows that for any Sasakian metric  $g'$  on  $(M, \eta)$  and any closed  $p$ -form  $\beta$  with  $p \leq n$ , we have

$$\Pi_{\Delta'}\beta \in \Omega_{i_\xi, d}^p(M).$$

Moreover, since  $d$  commutes with  $L$ , by applying (2.3) and Theorem 3.6 to  $(M, g')$  we get

$$(4.2) \quad L^{n-p+1}\Pi_{\Delta'}\beta = \frac{1}{2}d(\epsilon_\eta L^{n-p}\Pi_{\Delta'}\beta) = \frac{1}{2}d(F_p\Pi_{\Delta'}\beta) = 0.$$

Indeed, we will prove that given an arbitrary  $p$ -form  $\gamma$  with  $p \leq n$  such that

$$(4.3) \quad i_\xi\gamma = 0, \quad d\gamma = 0, \quad L^{n-p+1}\gamma = 0,$$

the cohomology classes of  $\epsilon_\eta L^{n-p}\gamma$  and of  $\epsilon_\eta L^{n-p}\Pi_{\Delta}\gamma$  are equal. As a consequence we will get that

$$[\epsilon_\eta L^{n-p}\Pi_{\Delta'}\beta] = [\epsilon_\eta L^{n-p}\Pi_{\Delta}\Pi_{\Delta'}\beta] = [\epsilon_\eta L^{n-p}\Pi_{\Delta}\beta],$$

since  $\Pi_{\Delta'}\beta$  satisfies (4.3) and  $\Pi_{\Delta}\Pi_{\Delta'}\beta = \Pi_{\Delta}\beta$  as  $[\Pi_{\Delta'}\beta] = [\beta]$ .

We start by recalling some facts on Hodge theory for compact manifolds. Let  $M$  be a compact Riemannian manifold. Denote by  $G$  the Green operator for  $\Delta$  acting on the de Rham complex  $\Omega^*(M)$  of  $M$ . Then

$$(4.4) \quad I - \Delta G = \Pi_{\Delta}, \quad I - G\Delta = \Pi_{\Delta}, \quad dG = Gd, \quad \delta G = G\delta.$$

**Proposition 4.1.** *Let  $(M, g)$  be a compact Riemannian manifold and  $\xi$  a Killing vector field on  $M$ . Then  $\mathcal{L}_\xi$  commutes with the Green operator  $G$ .*

*Proof.* By Theorem 3.7.1 in [12], we have that  $\mathcal{L}_\xi\omega = 0$  for every harmonic form  $\omega$ . Thus  $\mathcal{L}_\xi\Pi_{\Delta} = 0$ . Let us show that  $\Pi_{\Delta}\mathcal{L}_\xi = 0$ . Using (2.12) and that  $d^* = \delta$ ,  $i_\xi^* = \epsilon_\eta$ , we get

$$\mathcal{L}_\xi^* = \{d, i_\xi\}^* = \{\epsilon_\eta, \delta\} = -\mathcal{L}_\xi.$$

Hence for any  $\beta \in \Omega^k(M)$  and  $\omega \in \Omega_{\Delta}^k(M)$  we have

$$(\mathcal{L}_\xi\beta, \omega) = -(\beta, \mathcal{L}_\xi\omega) = 0,$$

where  $(\ , \ )$  denotes the usual global scalar product of differential forms on a compact Riemannian manifold. This shows that  $\mathcal{L}_\xi \beta$  is orthogonal to the subspace  $\Omega_\Delta^k(M)$  of  $\Omega^k(M)$  and thus  $\Pi_\Delta \mathcal{L}_\xi \beta = 0$ . Now, from (4.4) we get

$$\mathcal{L}_\xi - \mathcal{L}_\xi \Delta G = \mathcal{L}_\xi \Pi_\Delta = 0, \quad \mathcal{L}_\xi - G \Delta \mathcal{L}_\xi = \Pi_\Delta \mathcal{L}_\xi = 0.$$

We know from Theorem 2.1 that  $\mathcal{L}_\xi$  commutes with  $d$  and  $\delta$ , and thus also with  $\Delta = \{d, \delta\}$ . Hence

$$G \mathcal{L}_\xi = G \mathcal{L}_\xi \Delta G = G \Delta \mathcal{L}_\xi G = \mathcal{L}_\xi G.$$

q.e.d.

Let us introduce the auxiliary map

$$\begin{aligned} \mathcal{A}_p: \Omega^{p-1}(M) &\rightarrow \Omega^{2n-p+1}(M) \\ \alpha &\mapsto (n-p+1)L^{n-p}di_\phi d\alpha + L^{n-p+1}\Delta\alpha. \end{aligned}$$

We will study the interplay between  $\mathcal{A}_p$  and  $\delta$ ,  $d$ ,  $\Delta$ . The following technical lemma is needed.

**Lemma 4.2.** *Let  $M$  be a Sasakian manifold. Then for any  $k \geq 1$ , we have*

$$(4.5) \quad [\delta, L^k] = -kL^{k-1}\mathcal{L}_\phi + 2k\epsilon_\eta L^{k-1}(n - \deg - (k-1)).$$

*Proof.* For any two linear endomorphisms  $a$  and  $b$  of an arbitrary vector space  $V$ , we denote by  $\text{ad}_b(a)$  their commutator  $[b, a]$ . It can be checked (e.g. by induction) that

$$(4.6) \quad ab^k = \sum_{j=0}^k (-1)^j \binom{k}{j} b^{k-j} \text{ad}_b^j(a).$$

We will apply formula (4.6) to the case  $a = \delta$  and  $b = L$ . Dualizing (2.5), since  $i_\phi^* = -i_\phi$ , we get

$$\text{ad}_L(\delta) = [L, \delta] = [d, -i_\phi] - 2\epsilon_\eta(n - \deg) = \mathcal{L}_\phi - 2\epsilon_\eta(n - \deg).$$

Moreover, since  $\mathcal{L}_\phi$  is a derivation and  $L = \epsilon_\Phi$ , we get

$$[\mathcal{L}_\phi, L] = \epsilon_{\mathcal{L}_\phi}\Phi = 0.$$

Therefore

$$\text{ad}_L^2(\delta) = [L, \mathcal{L}_\phi - 2\epsilon_\eta n + 2\epsilon_\eta \deg] = 2\epsilon_\eta [L, \deg] = -4\epsilon_\eta L.$$

Thus  $\text{ad}_L^j(\delta) = 0$  for all  $j \geq 3$ . Now the claim of the lemma follows from (4.6). q.e.d.

**Theorem 4.3.** *Let  $(M, \eta, g)$  be a compact Sasakian manifold of dimension  $2n+1$ ,  $1 \leq p \leq n$ , and  $\alpha \in \Omega_{\mathcal{L}_\xi, \delta}^{p-1}(M)$ . Then  $\mathcal{A}_p \alpha$  is coclosed and*

$$(4.7) \quad \Delta \mathcal{A}_p \alpha = \mathcal{A}_p \Delta \alpha.$$

*Proof.* To check that

$$(4.8) \quad \mathcal{A}_p\alpha = (n-p+1)L^{n-p}di_\phi d\alpha + L^{n-p+1}\Delta\alpha$$

is coclosed, we will repeatedly use Lemma 4.2. We have

$$(4.9) \quad \delta(L^{n-p}di_\phi d\alpha) = [\delta, L^{n-p}] di_\phi d\alpha + L^{n-p}\delta di_\phi d\alpha.$$

From Lemma 4.2 we get

$$[\delta, L^{n-p}] di_\phi d\alpha = -(n-p)L^{n-p-1}\mathcal{L}_\phi di_\phi d\alpha,$$

since the second summand in (4.5) vanishes in this case due to the degree of  $\alpha$ . Moreover, by using (2.11) we have  $\mathcal{L}_\phi di_\phi d\alpha = \mathcal{L}_\phi^2 d\alpha = -2Ld\mathcal{L}_\xi\alpha = 0$ . Therefore we get

$$(4.10) \quad [\delta, L^{n-p}] di_\phi d\alpha = 0.$$

Now we compute the second summand of (4.9). We have

$$(4.11) \quad \delta di_\phi d\alpha = \Delta i_\phi d\alpha - d\delta i_\phi d\alpha.$$

From formula (2.7) for  $[\Delta, i_\phi]$  we get

$$\begin{aligned} (4.12) \quad \Delta i_\phi d\alpha &= -2(\mathcal{L}_\xi - i_\xi d + \epsilon_\eta \delta) d\alpha + i_\phi d\Delta\alpha \\ &= -2\epsilon_\eta \delta d\alpha + i_\phi d\Delta\alpha \\ &= -2\epsilon_\eta \Delta\alpha + i_\phi d\Delta\alpha, \end{aligned}$$

since  $\alpha \in \Omega_{\mathcal{L}_\xi, \delta}^{p-1}(M)$  implies  $\delta d\alpha = \Delta\alpha$ . Now we compute the second summand of (4.11). From (2.6) we get

$$\begin{aligned} (4.13) \quad d\delta i_\phi d\alpha &= d[\delta, i_\phi] d\alpha + di_\phi \delta d\alpha \\ &= -\frac{1}{2}d[\Delta, i_\xi] d\alpha - 2(n-(p-1))di_\xi d\alpha + di_\phi \Delta\alpha. \end{aligned}$$

Note that  $d[\Delta, i_\xi] d\alpha = -[\Delta, i_\xi] d^2\alpha = 0$ , since

$$\{d, [\Delta, i_\xi]\} = \{[d, \Delta], i_\xi\} + [\Delta, \mathcal{L}_\xi] = 0.$$

Also  $di_\xi d\alpha = -d^2 i_\xi \alpha + d\mathcal{L}_\xi \alpha = 0$ . Thus (4.13) becomes

$$(4.14) \quad d\delta i_\phi d\alpha = di_\phi \Delta\alpha.$$

Substituting (4.12) and (4.14) in (4.11), we get

$$(4.15) \quad \delta di_\phi d\alpha = -2\epsilon_\eta \Delta\alpha + i_\phi d\Delta\alpha - di_\phi \Delta\alpha = -2\epsilon_\eta \Delta\alpha + \mathcal{L}_\phi \Delta\alpha.$$

Thus, in view of (4.10), the formula (4.9) becomes

$$(4.16) \quad \delta(L^{n-p}di_\phi d\alpha) = -2\epsilon_\eta L^{n-p}\Delta\alpha + L^{n-p}\mathcal{L}_\phi \Delta\alpha.$$

Now we compute the value of  $\delta$  on the second summand of  $\mathcal{A}_p\alpha$ . Since  $\delta\Delta\alpha = \Delta\delta\alpha = 0$ , from Lemma 4.2 it follows that

$$\begin{aligned} \delta(L^{n-p+1}\Delta\alpha) &= [\delta, L^{n-p+1}] \Delta\alpha \\ (4.17) \quad &= -(n-p+1)L^{n-p}\mathcal{L}_\phi\Delta\alpha \\ &\quad + 2(n-p+1)\epsilon_\eta L^{n-p}\Delta\alpha \\ &= -(n-p+1)(L^{n-p}\mathcal{L}_\phi\Delta\alpha - 2\epsilon_\eta L^{n-p}\Delta\alpha). \end{aligned}$$

From (4.8), (4.16) and (4.17), we get that  $\delta\mathcal{A}_p\alpha = 0$ , that is  $\mathcal{A}_p\alpha$  is coclosed.

Now we will prove that  $\Delta\mathcal{A}_p\alpha = \mathcal{A}_p\Delta\alpha$ . Since  $\mathcal{A}_p\alpha$  is coclosed, we have  $\Delta\mathcal{A}_p\alpha = \delta d\mathcal{A}_p\alpha$ . As  $d$  commutes with  $L$ , we get from (4.8) that

$$d\mathcal{A}_p\alpha = L^{n-p+1}d\Delta\alpha.$$

Thus

$$\Delta\mathcal{A}_p\alpha = \delta d\mathcal{A}_p\alpha = [\delta, L^{n-p+1}] d\Delta\alpha + L^{n-p+1}\delta d\Delta\alpha.$$

By Lemma 4.2 we get

$$\begin{aligned} [\delta, L^{n-p+1}] d\Delta\alpha &= -(n-p+1)L^{n-p}\mathcal{L}_\phi d\Delta\alpha \\ &= (n-p+1)L^{n-p}di_\phi d\Delta\alpha. \end{aligned}$$

Next,

$$L^{n-p+1}\delta d\Delta\alpha = L^{n-p+1}\Delta\delta d\alpha = L^{n-p+1}\Delta^2\alpha.$$

Thus,

$$\Delta\mathcal{A}_p\alpha = (n-p+1)L^{n-p}di_\phi d\Delta\alpha + L^{n-p+1}\Delta^2\alpha = \mathcal{A}_p\Delta\alpha.$$

q.e.d.

**Corollary 4.4.** *Let  $(M, \eta, g)$  be a compact Sasakian manifold of dimension  $2n+1$  and  $\alpha \in \Omega_{\mathcal{L}_\xi, \delta}^{p-1}(M)$  with  $p \leq n$ , such that  $L^{n-p+1}d\Delta\alpha = 0$ . Then  $\mathcal{A}_p\alpha = 0$ .*

*Proof.* It is easy to check that  $\mathcal{A}_p\alpha$  is closed. It is also coclosed by Theorem 4.3. Therefore  $\mathcal{A}_p\alpha$  is harmonic. Now we consider the form  $\mathcal{A}_pG\alpha$ . It follows from (4.4) and Proposition 4.1 that  $G\alpha \in \Omega_{\mathcal{L}_\xi, \delta}^{p-1}(M)$ . By Theorem 4.3, we have

$$\Delta\mathcal{A}_pG\alpha = \mathcal{A}_p\Delta G\alpha = \mathcal{A}_p\alpha - \mathcal{A}_p\Pi_\Delta\alpha.$$

It is immediate from the definition of  $\mathcal{A}_p$  that  $\mathcal{A}_p\omega = 0$  for any harmonic  $(p-1)$ -form  $\omega$ , in particular for  $\Pi_\Delta\alpha$ . Thus  $\Delta\mathcal{A}_pG\alpha = \mathcal{A}_p\alpha$ . As  $\mathcal{A}_p\alpha$  is harmonic, we obtain

$$0 = (\Delta\mathcal{A}_p\alpha, \mathcal{A}_pG\alpha) = (\mathcal{A}_p\alpha, \Delta\mathcal{A}_pG\alpha) = (\mathcal{A}_p\alpha, \mathcal{A}_p\alpha).$$

Thus  $\mathcal{A}_p\alpha = 0$ .

q.e.d.

Now we will prove the main result of the article.

**Theorem 4.5.** *Let  $(M, \eta, g)$  be a compact Sasakian manifold,  $p \leq n$ , and  $\beta$  a closed  $p$ -form. For any  $\beta' \in [\beta]$  such that*

$$(4.18) \quad i_\xi \beta' = 0, \quad L^{n-p+1} \beta' = 0,$$

*we have  $Lef_p([\beta]) = [\epsilon_\eta L^{n-p} \beta']$ . In particular, the Lefschetz map  $Lef_p$  does not depend on the choice of a compatible Sasakian metric on  $(M, \eta)$ .*

*Proof.* Let us define  $\gamma = \delta G(\beta' - \Pi_\Delta \beta)$ . Then, since  $d$  and  $\delta$  commute with  $G$  by (4.4) and  $\beta'$  is closed, we get

$$(4.19) \quad \begin{aligned} d\gamma &= Gd\delta(\beta' - \Pi_\Delta \beta) \\ &= Gd\delta\beta' = G\Delta\beta' = \beta' - \Pi_\Delta\beta' = \beta' - \Pi_\Delta\beta. \end{aligned}$$

As  $Lef_p([\beta]) = [\epsilon_\eta L^{n-p} \Pi_\Delta \beta]$ , we have to prove that  $\epsilon_\eta L^{n-p} \beta'$  and  $\epsilon_\eta L^{n-p} \Pi_\Delta \beta$  are in the same cohomology class. Since

$$\epsilon_\eta L^{n-p} d\gamma = \epsilon_\eta L^{n-p} \beta' - \epsilon_\eta L^{n-p} \Pi_\Delta \beta,$$

it is enough to show that  $\epsilon_\eta L^{n-p} d\gamma$  is exact. Now, as

$$d(\epsilon_\eta L^{n-p} \gamma) = 2L^{n-p+1} \gamma - \epsilon_\eta L^{n-p} d\gamma,$$

the form  $\epsilon_\eta L^{n-p} d\gamma$  is exact if and only if  $L^{n-p+1} \gamma$  is exact.

Let  $\alpha = G\gamma$ , that is  $\alpha = \delta G^2(\beta' - \Pi_\Delta \beta)$ . We will now check that  $\alpha$  satisfies the hypotheses of Corollary 4.4, namely that  $\mathcal{L}_\xi \alpha$ ,  $\delta \alpha$ , and  $L^{n-p+1} d\Delta \alpha$  are zero. Since  $\mathcal{L}_\xi$  commutes with  $\delta$  by Theorem 2.1 and with  $G$  by Proposition 4.1, we get

$$\mathcal{L}_\xi \alpha = \delta G^2 \mathcal{L}_\xi (\beta' - \Pi_\Delta \beta).$$

As  $\Pi_\Delta \beta$  is harmonic and  $\xi$  is Killing, we have  $\mathcal{L}_\xi \Pi_\Delta \beta = 0$  by [12, Theorem 3.7.1]. Moreover,  $\beta'$  is closed and  $i_\xi \beta' = 0$  by assumption. Thus  $\mathcal{L}_\xi \beta' = 0$ . Therefore we get that  $\mathcal{L}_\xi \alpha = 0$ . As  $\alpha$  is in the image of  $\delta$ , we also have  $\delta \alpha = 0$ . Now

$$(4.20) \quad \Delta \alpha = \Delta G\gamma = \gamma - \Pi_\Delta \gamma = \gamma,$$

as  $\gamma$  is coexact, and this implies  $\Pi_\Delta \gamma = 0$ . Thus, by (4.18) and (4.19), we get

$$L^{n-p+1} d\Delta \alpha = L^{n-p+1} d\gamma = L^{n-p+1} \beta' - L^{n-p+1} \Pi_\Delta \beta = -L^{n-p+1} \Pi_\Delta \beta.$$

By Theorem 3.6 we know that  $\epsilon_\eta L^{n-p} \Pi_\Delta \beta$  is harmonic, therefore

$$L^{n-p+1} \Pi_\Delta \beta = \frac{1}{2} d(\epsilon_\eta L^{n-p} \Pi_\Delta \beta) = 0.$$

We conclude that  $L^{n-p+1} d\Delta \alpha = 0$ . Thus all conditions of Corollary 4.4 are satisfied for  $\alpha$ . Therefore  $\mathcal{A}_p \alpha = 0$  and thus by (4.20) we get that

$$\begin{aligned} L^{n-p+1} \gamma &= L^{n-p+1} \Delta \alpha = -(n-p+1) L^{n-p} d i_\phi d \alpha \\ &= d(-(n-p+1) L^{n-p} i_\phi d \alpha) \end{aligned}$$

is an exact form.

q.e.d.

**Corollary 4.6.** *Let  $(M, \eta)$  be a compact contact manifold of dimension  $2n+1$ . Suppose  $g$  and  $g'$  are two different Sasakian metrics on  $M$ , both compatible with  $\eta$ . Then, for any closed  $p$ -form  $\beta$  with  $p \leq n$ , it holds  $[\epsilon_\eta L^{n-p} \Pi_\Delta \beta] = [\epsilon_\eta L^{n-p} \Pi_{\Delta'} \beta]$ .*

*Proof.* We obviously have  $\Pi_{\Delta'} \beta \in [\beta]$ . By Proposition 2.2 applied to the Sasakian manifold  $(M, \eta, g')$ , we get  $i_\xi \Pi_{\Delta'} \beta = 0$ . Next by (4.2) we have  $L^{n-p+1} \Pi_{\Delta'} \beta = 0$ . Thus the form  $\beta' = \Pi_{\Delta'} \beta$  satisfies all the conditions of Theorem 4.5 for the Sasakian manifold  $(M, \eta, g)$ . Therefore  $[\epsilon_\eta L^{n-p} \Pi_{\Delta'} \beta] = \text{Lef}_p([\beta]) = [\epsilon_\eta L^{n-p} \Pi_\Delta \beta]$ . q.e.d.

## 5. Lefschetz contact manifolds

Let  $(M, \eta)$  be a compact contact manifold of dimension  $2n+1$  such that there exists a Sasakian metric on  $M$  compatible with the contact form  $\eta$ . Then by choosing an arbitrary such metric  $g$ , we can define the maps  $\text{Lef}_p$  for  $p \leq n$  as in (4.1) and these maps are isomorphisms.

Note that there is no obvious way to define similar maps between cohomology spaces of a general contact manifold  $(M, \eta)$  of dimension  $2n+1$ . To introduce the notion of Hard Lefschetz property for a contact manifold, we define the *Lefschetz relation* between cohomology groups  $H^p(M)$  and  $H^{2n+1-p}(M)$  of  $(M, \eta)$  to be

$$\mathcal{R}_{\text{Lef}_p} = \{([\beta], [\epsilon_\eta L^{n-p} \beta]) \mid \beta \in \Omega^p(M), d\beta = 0, i_\xi \beta = 0, L^{n-p+1} \beta = 0\}.$$

Thus if  $(M, \eta)$  admits a compatible Sasakian metric, from Theorem 4.5 it follows that  $\mathcal{R}_{\text{Lef}_p}$  is the graph of the isomorphism  $\text{Lef}_p$ . This justifies the following definition.

**Definition 5.1.** We say that a compact contact manifold  $(M, \eta)$  has the *Hard Lefschetz property* if for every  $p \leq n$  the relation  $\mathcal{R}_{\text{Lef}_p}$  is the graph of an isomorphism  $\text{Lef}_p : H^p(M) \longrightarrow H^{2n+1-p}(M)$ . Such manifolds will be called *Lefschetz contact manifolds*.

There is a simple extension to Lefschetz contact manifolds of the well-known property that the odd Betti numbers  $b_{2k+1}$  ( $0 \leq 2k+1 \leq n$ ) of compact Sasakian manifolds are even ([9]).

**Theorem 5.2.** *Let  $(M, \eta)$  be a Lefschetz contact manifold of dimension  $2n+1$ . Then for every odd  $p \leq n$  the Betti number  $b_p$  is even.*

*Proof.* Let  $p \leq n$ . Since  $\text{Lef}_p$  is an isomorphism, using Poincaré duality we can define a nondegenerate bilinear form  $B$  on the de Rham cohomology vector space  $H^p(M)$  by putting

$$B(x, x') = \int_M \text{Lef}_p(x) \smile x'.$$

Note that

$$(5.1) \quad B(x, x') = \int_M (\epsilon_\eta L^{n-p} \omega) \wedge \omega' = \int_M \eta \wedge \Phi^{n-p} \wedge \omega \wedge \omega',$$

where  $\omega \in x$  and  $\omega' \in x'$  are closed  $p$ -forms such that  $i_\xi\omega = i_\xi\omega' = 0$  and  $L^{n-p+1}\omega = L^{n-p+1}\omega' = 0$ . Such  $\omega$  and  $\omega'$  always exist since  $\mathcal{R}_{Lef_p}$  is the graph of a map. Now, (5.1) implies that  $B(x, x') = (-1)^p B(x', x)$ . It follows that, when  $p$  is odd, the vector space  $H^p(M)$  is even dimensional.  $\text{q.e.d.}$

It would be interesting to find a characterization of Lefschetz contact manifolds in the spirit of [5, 22]. It would be also interesting to find explicit examples of Lefschetz contact manifolds which do not admit any Sasakian structure. We will address these matters in our future research.

**Note added.** After the paper was submitted, Yi Lin has considered some of the above questions in [18].

## References

- [1] D. E. Blair, *Riemannian geometry of contact and symplectic manifolds*, second ed., Progress in Mathematics, vol. 203, Birkhäuser Boston Inc., Boston, MA, 2010, MR 2682326, Zbl 1246.53001.
- [2] C. P. Boyer & K. Galicki, *Sasakian geometry*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2008, MR 2382957, Zbl 1155.53002.
- [3] C. P. Boyer, K. Galicki, & J. Kollar, *Einstein metrics on spheres*, Ann. of Math. (2) **162** (2005), no. 1, 557–580, MR 2178969, Zbl 1093.53044.
- [4] C. P. Boyer, K. Galicki, & S. R. Simanca, *Canonical Sasakian metrics*, Comm. Math. Phys. **279** (2008), no. 3, 705–733, MR 2386725, Zbl 1142.53036.
- [5] J.-L. Brylinski, *A differential complex for Poisson manifolds*, J. Differential Geom. **28** (1988), no. 1, 93–114, MR 950556, Zbl 0634.58029.
- [6] D. Chinea, M. de León, & J. C. Marrero, *Topology of cosymplectic manifolds*, J. Math. Pures Appl. (9) **72** (1993), no. 6, 567–591, MR 1249410, Zbl 0845.53025.
- [7] A. El Kacimi-Alaoui, *Opérateurs transversalement elliptiques sur un feuilletage riemannien et applications*, Compositio Math. **73** (1990), no. 1, 57–106, MR 1042454, Zbl 0697.57014.
- [8] A. Frölicher & A. Nijenhuis, *Theory of vector-valued differential forms. I. Derivations of the graded ring of differential forms*, Nederl. Akad. Wetensch. Proc. Ser. A. **59** = Indag. Math. **18** (1956), 338–359, MR 0082554, Zbl 0079.37502.
- [9] T. Fujitani, *Complex-valued differential forms on normal contact Riemannian manifolds*, Tôhoku Math. J. (2) **18** (1966), 349–361, MR 0212730, Zbl 0143.44802.
- [10] A. Futaki, H. Ono, & G. Wang, *Transverse Kähler geometry of Sasaki manifolds and toric Sasaki-Einstein manifolds*, J. Differential Geom. **83** (2009), no. 3, 585–635, MR 2581358, Zbl 1188.53042.
- [11] J. P. Gauntlett, D. Martelli, J. Sparks, & D. Waldram, *A new infinite class of Sasaki-Einstein manifolds*, Adv. Theor. Math. Phys. **8** (2004), no. 6, 987–1000, MR 2194373, Zbl 1095.53034.
- [12] S. I. Goldberg, *Curvature and homology*, Pure and Applied Mathematics, Vol. XI, Academic Press, New York, 1962, MR 0139098, Zbl 0105.15601.

- [13] W. V. D. Hodge, *The theory and applications of harmonic integrals*, Cambridge University Press, 1952, 2d ed.
- [14] I. Kolář, P. W. Michor, and J. Slovák, *Natural operations in differential geometry*, Springer-Verlag, Berlin, 1993, MR 1202431, Zbl 0782.53013.
- [15] J. Kollar, *Einstein metrics on connected sums of  $S^2 \times S^3$* , J. Differential Geom. **75** (2007), no. 2, 259–272, MR 2286822, Zbl 1118.53026.
- [16] H. K. Kunduri & J. Lucietti, *Extremal Sasakian horizons*, Phys. Lett. B **713** (2012), no. 3, 308–312, MR 2944928.
- [17] S. Lefschetz, *L'analysis situs et la géométrie algébrique*, Gauthiers-Villars, Paris, 1924, JFM 50.0663.01.
- [18] Y. Lin, *Lefschetz contact manifolds and odd dimensional symplectic geometry*, arXiv:1311.1431v2 (2013).
- [19] J. Maldacena, *The large  $N$  limit of superconformal field theories and supergravity*, Adv. Theor. Math. Phys. **2** (1998), no. 2, 231–252, MR 1633016, Zbl 0914.53047.
- [20] D. Martelli, J. Sparks, & S.-T. Yau, *The geometric dual of  $a$ -maximisation for toric Sasaki-Einstein manifolds*, Comm. Math. Phys. **268** (2006), no. 1, 39–65, MR 2249795, Zbl 1190.53041.
- [21] D. Martelli, J. Sparks, & S.-T. Yau, *Sasaki-Einstein manifolds and volume minimisation*, Comm. Math. Phys. **280** (2008), no. 3, 611–673, MR 2399609, Zbl 1161.53029.
- [22] O. Mathieu, *Harmonic cohomology classes of symplectic manifolds*, Comment. Math. Helv. **70** (1995), no. 1, 1–9, MR 1314938, Zbl 0831.58004.
- [23] S. Sasaki, *On differentiable manifolds with certain structures which are closely related to almost contact structure. I*, Tôhoku Math. J. (2) **12** (1960), 459–476, MR 0123263, Zbl 0192.27903.
- [24] S. Tachibana, *On harmonic tensors in compact Sasakian spaces*, Tôhoku Math. J. (2) **17** (1965), 271–284, MR 0190878, Zbl 0132.16203.

DIPARTIMENTO DI MATEMATICA E INFORMATICA  
UNIVERSITÀ DEGLI STUDI DI CAGLIARI  
VIA OSPEDALE 72  
09124 CAGLIARI, ITALY  
*E-mail address:* b.cappellettimontano@gmail.com

CMUC, DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF COIMBRA  
3001-501 COIMBRA  
PORTUGAL  
*E-mail address:* antondenicola@gmail.com

CMUC, DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF COIMBRA  
3001-501 COIMBRA  
PORTUGAL  
*E-mail address:* yudin@mat.uc.pt