平成 10 年 6 月 19 日 印刷 平成 10 年 6 月 30 日 発行

編集発行者 東京工業大学大学院 理工学研究科数学教室

152-8551 東京都目黒区大岡山 2-12-1

印 刷 所 学術図書印刷株式会社 176-0012 東京都練馬区豊玉北 2-13-1

Printed by GAKUZITSUTOSHO PRINTING CO., LTD. Nerima-ku, Tokyo, Japan

CONTENTS

Kodai Mathematical Journal Vol. 21, No. 2

| Toshiharu Matsubara, On log Hodge structures of higher direct images | 81 |
|---|-----|
| TAKASHI SAKAI, On Riemannian manifolds admitting a function whose gra- | |
| dient is of constant norm II | 102 |
| DOAN THE HIEU, Decomposition of a k-covector with respect to a vector and | |
| computing its comass | 125 |
| PING LI, Meromorphic functions sharing three values or sets CM | 138 |
| Yoshiaki Fukuma, On sectional genus of k -very ample line bundles on | |
| smooth surfaces with non-negative Kodaira dimension | 153 |
| QING CAI ZHANG, The uniqueness of meromorphic functions with their de- | |
| rivatives | 179 |
| Takeshi Kawachi, On d-spannedness of the adjoint bundles on polarized | |
| manifolds | 185 |
| TADASHI NAKAJIMA, Joint distribution of the first hitting time and first hit- | |
| ting place for a random walk | 192 |
| Mei-jiao Wang and Shi-jie Li, Submanifolds with parallel mean curvature | |
| vector in a sphere | 201 |
| HELMUT A. HAMM AND LÊ DŨNG TRÁNG, Vanishing theorems for constructible | |
| sheaves II | 208 |