

## ON SUBGROUPS OF A PSEUDO LATTICE ORDERED GROUP

J. JAKUBÍK

**The purpose of this note is to investigate some problems raised in a recent paper of Conrad and Teller concerning  $o$ -ideals and  $p$ -subgroups in an abelian pseudo lattice ordered group.**

The concept of a pseudo lattice ordered group (“ $p$ -group”) has been introduced by Conrad [1]. In recent papers by Teller [5] and Conrad and Teller [2] there is developed a systematic theory of  $p$ -groups. Let  $G$  be an abelian  $p$ -group. In § 3 it is proved that if  $M$  is a subgroup of  $G$  such that  $\{a, b\} \cap M \neq \emptyset$  for any pair of  $p$ -disjoint elements  $a, b \in G$ , then  $M$  contains a prime  $o$ -ideal; this generalizes a result from [2]. In § 4 we prove that the intersection of two  $p$ -subgroups of a  $p$ -group  $G$  need not be a  $p$ -subgroup of  $G$ . Moreover, if  $\Delta$  is a partially ordered set and for each  $\delta \in \Delta$   $H_\delta \neq \{0\}$  is a linearly ordered group, then for the mixed product  $G = V(\Delta, H_\delta)$  the following conditions are equivalent: (i) for any two  $p$ -subgroups  $A, B$  of  $G$  their intersection  $A \cap B$  is a  $p$ -subgroup of  $G$  as well; (ii)  $G$  is an  $l$ -group. If  $A$  is an  $o$ -ideal of a  $p$ -group  $G$  and  $B$  is a  $p$ -subgroup of  $G$ , then  $A + B$  is a  $p$ -subgroup of  $G$ .

2. Preliminaries. Let  $G$  be a partially ordered group.  $G$  is a Riesz group (cf. Fuchs [3], [4]) if it is directed and if from  $a_i, b_j \in G, a_i \leq b_j$  ( $i, j = 1, 2$ ) it follows that there exists  $c \in G$  satisfying  $a_i \leq c \leq b_j$  ( $i, j = 1, 2$ ).  $G$  is a  $p$ -group (cf. [1] and [5]) if it is Riesz and if each  $g \in G$  has a representation  $g = a - b$  such that  $a, b \in G, a \geq 0, b \geq 0$  and

$$(*) \quad x \in G, x \leq a, x \leq b \implies nx \leq a, nx \leq b$$

for any positive integer  $n$ .

Throughout the paper  $G$  denotes an abelian  $p$ -group. Elements  $a, b \in G, a \geq 0, b \geq 0$  satisfying (\*) are called  $p$ -disjoint. A subgroup  $M$  of  $G$  is a  $p$ -subgroup, if for each  $m \in M$  there are elements  $a, b \in M$  such that  $a, b$  are  $p$ -disjoint in  $G$  and  $m = a - b$ . A subgroup  $C$  of  $G$  is an  $o$ -ideal, if it is directed and if  $0 \leq g \leq c \in C, g \in G$  implies  $g \in C$ . Let  $O(G)$  be the system of all  $o$ -ideals of  $G$  (partially ordered by the set inclusion). An  $o$ -ideal  $C$  of  $G$  is called prime, if  $G/C$  is a linearly ordered group. For any pair  $a, b$  of  $p$ -disjoint elements  $H(a, b)$  denotes the subgroup of  $G$  generated by the set

$$\{0 \leq m \in G \mid m \leq a, m \leq b\}.$$

Then  $H(a, b) \in O(G)$  (cf. [2]).

Let  $\Delta$  be a partially ordered set and let  $H_\delta \neq \{0\}$  be a linearly ordered group for each  $\delta \in \Delta$ . Let  $V = V(\Delta, H_\delta)$  be the set of all  $\Delta$ -vectors  $v = (\dots, v_\delta, \dots)$  where  $v_\delta \in H_\delta$ , for which the support  $S(v) = \{\delta \in \Delta \mid v_\delta \neq 0\}$  contains no infinite ascending chain. An element  $v \in V$ ,  $v \neq 0$  is defined to be positive if  $v_\delta > 0$  for each maximal element  $\delta \in S(v)$ . Then ([2], Th. 5.1)  $V$  is a  $p$ -group;  $V$  is an 1-group if and only if  $\Delta$  is a root system (i.e.,  $\{\delta \in \Delta \mid \delta \geq \gamma\}$  is a chain for each  $\gamma \in \Delta$ ).

**3. Subgroups containing a prime  $o$ -ideal.** The following assertion has been proved in [2] (Proposition 4.3):

(A) For  $M \in O(G)$ , the following are equivalent: (1)  $M$  is prime; (2) the  $o$ -ideals of  $G$  that contain  $M$  form a chain; (3) if  $a$  and  $b$  are  $p$ -disjoint in  $G$ , then  $a \in M$  or  $b \in M$ .

Further it is remarked in [2] that each subgroup  $M$  of  $G$  fulfilling (3) is a  $p$ -subgroup and any subgroup containing a prime  $o$ -ideal satisfies (3); then it is asked whether a subgroup  $M$  of a  $p$ -group  $G$  satisfies (3) if and only if it contains a prime  $o$ -ideal (a similar assertion is known to be valid for lattice ordered groups). We shall prove that the answer is positive.

We need the following propositions (cf. [2] and [5]):

(B) Let  $g = a - b \in G$  where  $a$  and  $b$  be  $p$ -disjoint. Then  $g = x - y$ , where  $x$  and  $y$  are  $p$ -disjoint, if and only if  $x = a + m$  and  $y = b + m$  for some  $m \in H(a, b)$ .

(C) If  $a$  and  $b$  are  $p$ -disjoint, then  $na$  and  $nb$  are  $p$ -disjoint for any positive integer  $n$  and  $H(a, b) = H(na, nb)$  ([2], Proposition 3.1).

**LEMMA 1.** *Let  $M$  be a subgroup of  $G$  fulfilling (3) and let  $a, b$  be  $p$ -disjoint elements in  $G$ . Then  $H(a, b) \subset M$ .*

*Proof.* Let  $h \in H(a, b)$ . According to (3) we may assume without loss of generality that  $a \in M$ . Suppose (by way of contradiction) that  $h \notin M$ . Then  $a + h \notin M$ , hence by (B)  $b + h \in M$ , and analogously  $b - h \in M$ , thus  $2b \in M$ . Further  $2a + h \notin M$  and therefore according to (C) and (B)  $2b + h \in M$ , which implies  $h \in M$ .

**LEMMA 2.** *Let  $M$  be a subgroup of  $G$  satisfying (3) and let  $X = \{X_i\}$  be the system of all  $o$ -ideals of  $G$  such that  $X_i \subset M$ . Then the system  $X$  has a largest element.*

*Proof.* Let  $Y$  be the subgroup of  $G$  generated by the set  $\bigcup X_i$ .

Then  $Y \subset M$  and  $Y$  is the supremum of the system  $\{X_i\}$  in the lattice  $\mathcal{S}$  of all subgroups of  $G$ . Since  $O(G)$  is a complete sublattice of  $\mathcal{S}$  ([2], Th. 2.1),  $Y \in O(G)$  and thus  $Y \in X$ .

Let  $H$  be the subgroup of  $G$  generated by the set  $\bigcup H(a, b)$  where  $a, b$  runs over the system of all  $p$ -disjoint pairs of elements in  $G$ . Since each set  $H(a, b)$  is an  $o$ -ideal ([2]),  $H = \bigvee H(a, b)$  ( $a$  and  $b$   $p$ -disjoint in  $G$ ) where  $\bigvee$  denotes the supremum in the lattice  $O(G)$ . According to Lemma 1  $H \subset M$  whenever the subgroup  $M$  of  $G$  satisfies (3).

For any  $u, v \in G$ ,  $u \leq v$ , the interval  $[u, v]$  is the set

$$\{x \in G \mid u \leq x \leq v\}.$$

LEMMA 3. *Let  $M$  be a subgroup of  $G$  satisfying (3) and let  $N$  be the largest  $o$ -ideal of  $G$  that is contained in  $M$ . Let  $g \in G$ ,  $g > 0$ . Then*

$$[0, g] \subset M \implies g \in N.$$

*Proof.* According to Lemma 2 the largest  $o$ -ideal  $N$  in  $M$  exists. Assume that  $g \in G$ ,  $g > 0$ ,  $[0, g] \subset M$ . The set

$$Z = \bigcup_{n=1}^{\infty} [-ng, ng]$$

is clearly an  $o$ -ideal in  $G$ . Let  $z \in Z$ , hence  $z \in [-ng, ng]$  for a positive integer  $n$ . This implies  $0 \leq y \leq 2ng$  where  $y = z + ng$ . Since  $G$  is a Riesz group, according to [3, p. 158, Th. 27] there are elements  $g_1, \dots, g_{2n} \in G$ ,  $0 \leq g_i \leq g$  such that  $y = g_1 + \dots + g_{2n}$ . Thus  $g_i \in M$ , therefore  $y \in M$  and  $Z \subset M$ . Now we have  $Z \subset N$  and so  $g \in N$ .

LEMMA 4. *Let  $M$  be a subgroup of  $G$  fulfilling (3) and let  $N$  be the largest  $o$ -ideal of  $G$  contained in  $M$ . Then  $G/N$  is a linearly ordered group.*

*Proof.* Assume (by way of contradiction) that  $G/N$  is not linearly ordered. According to Lemma 1  $H \subset N$ , hence by [2], Theorem 4.1  $G/N$  is a lattice ordered group. Thus there exist elements  $X, Y \in G/N$  such that  $X \wedge Y = \bar{0}$ ,  $X > \bar{0}$ ,  $Y > \bar{0}$  ( $\bar{0}$  being the neutral element of  $G/N$ ). From [2] (Proposition 2.2, (ii)) it follows that there are elements  $x \in X$ ,  $y \in Y$  such that  $x$  and  $y$  are  $p$ -disjoint in  $G$  and hence  $x \in M$  or  $y \in M$ . Clearly  $x \notin N$ ,  $y \notin N$  and thus according to Lemma 3 there exist elements  $x_1, y_1 \in G$  such that

$$0 < x_1 \leq x, \quad 0 < y_1 \leq y, \quad x_1 \notin M, \quad y_1 \notin M.$$

Then in  $G/N$  we have  $\bar{0} < x_1 + N \leq x + N = X$ ,  $\bar{0} < y_1 + N \leq y + N = Y$ , whence

$$(x_1 + N) \wedge (y_1 + N) = \bar{0}.$$

Thus by using repeatedly [2], Proposition 2.2, we can choose elements  $x_2 \in x_1 + N$ ,  $y_2 \in y_1 + N$  such that  $x_2$  and  $y_2$  are  $p$ -disjoint in  $G$ . Therefore (without loss of generality) we may assume  $x_2 \in M$  and this implies  $x_1 \in x_1 + N = x_2 + N \subset M$ , a contradiction. The proof is complete.

**THEOREM 1.** *Let  $M$  be a subgroup of a  $p$ -group  $G$ . Then (3)  $\Rightarrow$  (2) and the condition (3) is equivalent to (1')  $M$  contains a prime  $o$ -ideal.*

*Proof.* According to Lemma 4 (3)  $\Rightarrow$  (1'). By [2] (1')  $\Rightarrow$  (3). Assume that  $M$  is a subgroup of  $G$  fulfilling (3). Let  $K_1, K_2$  be  $o$ -ideals of  $G$  such that  $M \subset K_1 \cap K_2$ . Let  $N$  have the same meaning as in Lemma 4. Since  $N \subset M$ ,

$$K_1 \subset K_2 \iff K_1/N \subset K_2/N.$$

$K_1/N$  and  $K_2/N$  are  $o$ -ideals of  $G/N$  and  $G/N$  is linearly ordered, hence  $K_1/N \subset K_2/N$  or  $K_2/N \subset K_1/N$ ; therefore (2) holds.

If  $M$  is an  $o$ -ideal of  $G$  satisfying (3), then by Theorem 1  $M$  contains a prime  $o$ -ideal  $N$ ; according to [2] (Corollary 1 to the Induced Homomorphism Theorem)  $G/M$  is isomorphic to  $(G/N)/(M/N)$  and hence ( $G/N$  being linearly ordered)  $G/M$  is a linearly ordered group and  $M$  is prime. Thus it follows from Theorem 1 that (3)  $\Rightarrow$  (1) for  $M \in O(G)$  (cf. (A)).

Let us remark that if  $M$  is a subgroup of  $G$  fulfilling (3) then  $M$  need not contain any nonzero  $o$ -ideal that is a lattice; further (3) is not implied by (2).

**EXAMPLE 1.** Let  $B$  be an infinite Boolean algebra that has no atoms and put  $\Delta = \{b \in B \mid b \neq 0\}$ . For each  $\delta \in \Delta$  let  $H_\delta = E$  where  $E$  is the additive group of all integers with the natural order,  $G = V(\Delta, H_\delta)$ . Let  $M = \{v \in G \mid v_1 = 0\}$  (by 1 we denote the greatest element of  $B$ ). Then  $M$  is a prime  $o$ -ideal of  $G$ , hence  $M$  satisfies (3) and  $M$  contains no lattice ordered  $o$ -ideal different from  $\{0\}$ .

**EXAMPLE 2.** Let  $\Delta = \{\delta_1, \delta_2, \delta_3\}$ , where  $\delta_1 < \delta_3$ ,  $\delta_2 < \delta_3$  and  $\delta_1, \delta_2$  are incomparable. Put  $H_{\delta_i} = E$  ( $i = 1, 2, 3$ ),  $G = V(\Delta, H_\delta)$ ,  $M = \{v \in G \mid v_{\delta_1} = v_{\delta_2} = 0\}$ . Then the only  $o$ -ideal that contains  $M$  is  $G$ , thus (2) holds. Let  $a, b \in G$  such that  $a_{\delta_1} = 1$ ,  $a_{\delta_2} = a_{\delta_3} = 0$ ,  $b_{\delta_2} = 1$ ,  $b_{\delta_1} = b_{\delta_3} = 0$ . The elements  $a$  and  $b$  are  $p$ -disjoint in  $G$  and  $a \notin M$ ,  $b \notin M$ , hence  $M$  does not fulfil (3).

4. **Intersections and sums of two  $p$ -subgroups.** Another problem formulated in [2] is whether the intersection of two  $p$ -subgroups of a  $p$ -group  $G$  must be a  $p$ -subgroup of  $G$ ; there is remarked in [2] that this conjecture seems rather dubious. The answer to this problem is negative.

**EXAMPLE 3.** Let  $\Delta = \{\delta_1, \delta_2, \delta_3\}$ , where  $\delta_1 > \delta_3$ ,  $\delta_2 > \delta_3$  and  $\delta_1, \delta_2$  are incomparable. Let  $H_{\delta_i} = E(i = 1, 2, 3)$ ,  $G = V(\Delta, H_\delta)$ . We write  $v(\delta_i)$  instead of  $v_{\delta_i}$ . Let  $c_i \neq 0$  ( $i = 1, 2$ ) be positive integers,  $c_1 \neq c_2$ . Denote

$$A_i = \{v \in G \mid v(\delta_3) = c_i[v(\delta_1) + v(\delta_2)]\}$$

( $i = 1, 2$ ). Let  $i \in \{1, 2\}$  be fixed. For proving that  $A_i$  is a  $p$ -subgroup of  $G$  we have to verify that to each  $v \in A_i$  we can choose  $a, b \in A_i$ ,  $a \geq 0$ ,  $b \geq 0$  such that (\*) holds and  $v = a - b$ . It is easy to verify that it suffices to consider the case when 0 and  $v$  are incomparable, hence we may assume  $v(\delta_1) > 0$ ,  $v(\delta_2) < 0$  (the case  $v(\delta_1) < 0$ ,  $v(\delta_2) > 0$  being analogous). Let  $a, b \in G$ ,

$$\begin{aligned} a(\delta_1) &= v(\delta_1), \quad a(\delta_2) = 0, \quad a(\delta_3) = c_i a(\delta_1), \\ b(\delta_1) &= 0, \quad b(\delta_2) = -v(\delta_2), \quad b(\delta_3) = -c_i v(\delta_2). \end{aligned}$$

Then  $a$  and  $b$  have the desired properties, hence  $A_i$  is a  $p$ -subgroup of  $G$ . Denote  $C = A_1 \cap A_2$ . If  $v \in C$ , we have

$$c_1[v(\delta_1) + v(\delta_2)] = v(\delta_3) = c_2[v(\delta_1) + v(\delta_2)]$$

and thus (since  $c_1 \neq c_2$ )  $v(\delta_3) = 0$ ,  $v(\delta_2) = -v(\delta_1)$ . Therefore any element  $v \in C$ ,  $v \neq 0$  is incomparable with 0 and  $C$  is not a  $p$ -subgroup of  $G$ .

The method used in this example can be employed for proving the following theorem:

**THEOREM 2.** *Let  $\Delta$  be a partially ordered set and for each  $\delta \in \Delta$  let  $H_\delta \neq \{0\}$  be a linearly ordered group,  $V = V(\Delta, H_\delta)$ . If  $V$  is not lattice ordered, then  $V$  contains infinitely many pairs of  $p$ -subgroups  $A_1, A_2$  such that  $A_1 \cap A_2$  is not a  $p$ -subgroup of  $V$ .*

*Proof.* Assume that  $V$  is not lattice ordered. Then  $\Delta$  is no root system, hence there exist elements  $\delta_1, \delta_2, \delta_3$  such that  $\delta_1 > \delta_3$ ,  $\delta_2 > \delta_3$  and  $\delta_1, \delta_2$  are incomparable. Choose  $e_i \in H_{\delta_i}$ ,  $e_i > 0$  and let  $c_1, c_2$  be positive integers,  $c_1 \neq c_2$ . Let  $V_1 = \{v \in V \mid v_\delta = 0 \text{ for each } \delta \notin \{\delta_1, \delta_2, \delta_3\}\}$ ,

$$A_i = \{v \in V_1 \mid v(\delta_1) = n_1 e_1, \quad v(\delta_2) = n_2 e_2, \quad v(\delta_3) = c_i(n_1 + n_2)e_3\}$$

where  $n_1$  and  $n_2$  run over the set of all integers ( $i = 1, 2$ ). Analo-

gously as in Example 3 we can verify that  $A_1$  and  $A_2$  are  $p$ -subgroups of  $V$ . Let  $v \in C = A_1 \cap A_2$ . Then  $c_1(n_1 + n_2) = c_2(n_1 + n_2)$ , thus  $n_2 = -n_1$  and  $v(\delta_3) = 0$ . Therefore no element of  $C$  is strictly positive and  $C$  is no  $p$ -subgroup of  $G$ . Since the positive integers  $c_1 \neq c_2$  are arbitrary there exist infinitely many such pairs  $A_1, A_2$ .

As a corollary, we obtain:

**PROPOSITION 1.** *Let  $V = V(\Delta, H_\delta)$ , where each  $H_\delta$  is linearly ordered. Then the following conditions are equivalent: (i)  $V$  is lattice ordered; (ii) if  $A$  and  $B$  are  $p$ -subgroups of  $V$ , then  $A \cap B$  is a  $p$ -subgroup of  $V$  as well.*

*Proof.* By Theorem 2 (ii) implies (i). Let  $V$  be lattice ordered. Then a subgroup  $A$  of  $V$  is a  $p$ -subgroup of  $V$  if and only if it is an 1-subgroup of  $V$ ; since the intersection of two 1-subgroups is an 1-subgroup, (ii) is valid.

**PROPOSITION 2.** *Let  $\Delta$  be a partially ordered set and for any  $\delta \in \Delta$  let  $H_\delta \neq \{0\}$  be a linearly ordered group. Assume that there exist  $\delta_1, \delta_2, \delta_3 \in \Delta$  such that  $\delta_1 < \delta_3$ ,  $\delta_2 < \delta_3$  and  $\delta_1, \delta_2$  are incomparable,  $V = V(\Delta, H_\delta)$ . Then there are infinitely many  $p$ -subgroups  $A, B$  of  $V$  such that  $A + B$  is not a  $p$ -subgroup of  $V$ .*

*Proof.* Denote  $V_1 = \{v \in V \mid v(\delta) = 0 \text{ for each } \delta \notin \{\delta_1, \delta_2, \delta_3\}\}$  and let  $c$  be a fixed positive integer,  $e_i \in H_{\delta_i}$ ,  $e_i > 0$  ( $i = 1, 2, 3$ ). Put

$$\begin{aligned} A &= \{v \in V_1 \mid v(\delta_1) = ne_1, v(\delta_2) = -cne_2, v(\delta_3) = ne_3\}, \\ B &= \{v \in V_1 \mid v(\delta_1) = v(\delta_2) = 0, v(\delta_3) = ne_3\} \end{aligned}$$

where  $n$  runs over the set of all integers.  $A$  and  $B$  are linearly ordered subgroups of  $V$ , hence they are  $p$ -subgroups of  $V$ . The set  $C = A + B$  is the system of all elements  $v \in V_1$  such that

$$v(\delta_1) = n_1e_1, \quad v(\delta_2) = -cn_1e_2, \quad v(\delta_3) = n_2e_3$$

where  $n_1, n_2$  are arbitrary integers. Hence there is  $g \in C$  satisfying

$$g(\delta_1) = e_1, \quad g(\delta_2) = -ce_2, \quad g(\delta_3) = 0.$$

If  $g = a - b$ ,  $a \in C$ ,  $b \in C$ ,  $a \geq 0$ ,  $b \geq 0$ , then  $a \neq 0 \neq b$  (since  $g \not\geq 0$ ,  $g \not\leq 0$ ), thus  $a(\delta_3) = b(\delta_3) \geq e_3$ . There exists  $v \in V_1$  such that  $v(\delta_3) = a(\delta_3)$ ,  $v(\delta_1) < a(\delta_1)$  and  $b(\delta_1)$ ,  $v(\delta_2) < a(\delta_2)$  and  $b(\delta_2)$ . Thus  $v < a$ ,  $v < b$ , but  $2v \not< a$ ,  $2v \not< b$ . Therefore  $a$  and  $b$  are not  $p$ -disjoint in  $G$  and  $C$  is no  $p$ -subgroup of  $G$ .

One of the problems raised in [2] is affirmatively solved by

**THEOREM 3.** *Let  $A$  be an  $o$ -ideal of  $G$  and let  $B$  be a  $p$ -subgroup of  $G$ . Then  $A + B$  is a  $p$ -subgroup of  $G$ .*

*Proof.* Let us denote  $G/A = \bar{G}$  and for any  $t \in G$  write  $t + A = \bar{t}$ . Let  $A + B = X$ ,  $x \in X$ . There are elements  $a \in A$ ,  $b \in B$  such that  $x = a + b$  and since  $B$  is a  $p$ -subgroup there exist  $b_1, b_2 \in B$  such that  $b = b_1 - b_2$  and  $b_1, b_2$  are  $p$ -disjoint in  $G$ . Further  $x = u - v$ ,  $u, v \in G$ , where  $u$  and  $v$  are  $p$ -disjoint in  $G$ . According to [2]  $\bar{G}$  is a  $p$ -group and by [2], Proposition 2.2,  $\bar{b}_1$  and  $\bar{b}_2$  ( $\bar{u}$  and  $\bar{v}$ ) are  $p$ -disjoint in  $G$ . Further we have

$$\bar{x} = \bar{b}_1 - \bar{b}_2 = \bar{u} - \bar{v},$$

hence if we apply (B) (§ 3) to the  $p$ -group  $\bar{G}$  it follows that there exists  $\bar{m} \in H(\bar{u}, \bar{v})$  fulfilling

$$\bar{b}_1 = \bar{u} + \bar{m}, \quad \bar{b}_2 = \bar{v} + \bar{m}.$$

Again, by Proposition 2.2 of [2], there is  $m_1 \in \bar{m}$  such that  $m_1 \in H(u, v)$ . Thus according to (B) the elements  $u_1 = u + m_1$  and  $v_1 = v + m_1$  are  $p$ -disjoint in  $G$  and  $x = u_1 - v_1$ . Since

$$u_1 \in \bar{u}_1 = \bar{u} + \bar{m}_1 = \bar{u} + \bar{m} = \bar{b}_1 = b_1 + A \subset A + B = X$$

and analogously  $v_1 \in X$ , the set  $X$  is a  $p$ -subgroup of  $G$ .

#### REFERENCES

1. P. Conrad, *Representations of partially ordered Abelian groups as groups of real valued functions*, Acta Math. **116** (1966), 199-221.
2. P. Conrad and J. R. Teller, *Abelian pseudo lattice ordered groups*, Publications Math. (to appear)
3. L. Fuchs, *Partially ordered algebraic systems*, Moskva, 1965.
4. ———, *Riesz groups*, Ann. Scuola Norm. Sup. Pisa **19** (1965), 1-34.
5. J. R. Teller, *On abelian pseudo lattice ordered groups*, Pacific J. Math. **27** (1968), 411-419.

Received July 1, 1969.

TECHNICAL UNIVERSITY  
KOŠICE, CZECHOSLOVAKIA

