

THE DEFORMATION BEHAVIOUR OF THE KODAIRA DIMENSION OF ALGEBRAIC MANIFOLDS

(WITH AN APPENDIX BY K. UENO)

TADASHI ASHIKAGA

(Received October 1, 1979)

0. **Introduction.** We consider the following:

PROBLEM. *Is the Kodaira dimension of algebraic manifolds invariant under deformation?*

For curves and surfaces, the answer is affirmative. In the former case the result is clear. In the latter case it was proved by Iitaka [I₂] using the classification of surfaces, whereas for non-algebraic complex manifolds of dimension three, Nakamura [N] produced a counterexample to this problem.

On the other hand, Lieberman-Sernesi [LS] introduced a notion of the relative Kodaira dimension $\kappa(X/Y)$ for a family $f: X \rightarrow Y$ of algebraic manifolds, and proved that the Kodaira dimension κ of fibers over a countable intersection of Zariski open sets of Y is equal to $\kappa(X/Y)$ and κ of other fibers are greater than $\kappa(X/Y)$. Using this notion, we formulate our problem in the following way.

CONJECTURE DF _{n,k} . *Let $f: X \rightarrow Y$ be a family of n -dimensional algebraic manifolds with $\kappa(X/Y) = k$. Then for any fiber $X_y = f^{-1}(y)$ ($y \in Y$), we have $\kappa(X_y) = k$.*

Note that DF _{n,n} is true by Lieberman-Sernesi's theorem. If all the Conjectures DF _{$n,-\infty$} , DF _{$n,0$} , \dots , DF _{$n,n-1$} are true, then the deformation invariance of the Kodaira dimension in the algebraic case will be settled.

In this paper, we study Conjecture DF _{n,k} for $1 \leq k \leq n-1$. First we describe the geometric structure of every fiber of such a family as follows:

THEOREM I. *Let $f: X \rightarrow Y$ be a family of n -dimensional algebraic manifolds with $1 \leq \kappa(X/Y) \leq n-1$. Then for any $y \in Y$, the fiber X_y has the following property: There exist a nonsingular model X_y^* of X_y , a variety T and a fiber space $\psi: X_y^* \rightarrow T$ such that*

- (1) $\dim T = \kappa(X/Y)$
- (2) *There is an open set T' of T such that for any $t \in T'$, the*

fiber $\psi^{-1}(t)$ is irreducible nonsingular and is the quasi-specialization of parabolic varieties in the sense of Definition 4.1, i.e., $\psi^{-1}(t)$ is one of the irreducible components of a fiber of a certain degenerating family of varieties of Kodaira dimension zero (of so-called parabolic type).

By this theorem, Conjecture $DF_{n,k}$ for $1 \leq k \leq n - 1$ is reduced to a problem on $(n - k)$ -dimensional algebraic manifolds of Kodaira dimension zero:

THEOREM II. *Let (n, k) be a pair of positive integers with $1 \leq k \leq n - 1$. Assume (1) Conjecture $DF_{n-k,0}$ and (2) the lower-semicontinuity of κ for degenerating families of $(n - k)$ -dimensional parabolic manifolds, i.e., the κ of each components of a degenerate fiber of a degenerating family of parabolic manifolds is not larger than zero (see Conjecture PDG_{n-k} in § 5). Then Conjecture $DF_{n,k}$ is true.*

COROLLARY. $DF_{n,n-1}$ is true.

In § 1, we summarize some known results about the Kodaira dimension and give the definition of the relative Kodaira dimension. § 2 deals with general properties of the relative Kodaira dimension. As a preparation for § 4, § 3 is devoted to the study of a graded C -subalgebra of the canonical ring of an algebraic variety. In § 4, we study families over a nonsingular curve and prove Theorem I in that case. Theorems I and II are completely proved in § 5.

The author would like to express his hearty thanks to Professors Tadao Oda and Masa-Nori Ishida for their valuable advice and encouragements. The author also expresses his hearty thanks to Professor Kenji Ueno for writing an appendix to this paper.

2. Definitions and notations. We work in the category of schemes defined over the field of complex numbers C . An algebraic variety is an irreducible reduced C -scheme of finite type. A point of an algebraic variety is a closed point. Open sets are Zariski open sets. Unless otherwise stated, an algebraic manifold is a nonsingular complete algebraic variety. For a surjective morphism $f: V \rightarrow W$ from a variety V to another W , a fiber over a point in a certain countable intersection of nonempty open sets of W is called a *general fiber*. Note that a countable intersection of nonempty open sets of W is dense in W . $f: V \rightarrow W$ is called a *fiber space* if f is a surjective morphism from an algebraic variety V to another W such that the general fiber of f are connected. By a family $f: X \rightarrow Y$ of algebraic manifolds of dimension n , we mean that f is a proper smooth surjective morphism of relative

dimension n between algebraic varieties X and Y such that every fiber of f is connected.

Let V be an algebraic manifold and ω_V be its canonical sheaf. We put

$$R(V) = \bigoplus_{i \in \mathbf{Z}_{\geq 0}} R(V)_i, \quad R(V)_i = H^0(V, \omega_V^{\otimes i}),$$

where $\mathbf{Z}_{\geq 0}$ is the set of non-negative integers. We call $R(V)$ the canonical ring of V . We set

$$R(V)^{[i]} = C[R(V)_i]$$

which is the graded C -subalgebra of $R(V)$ generated by $R(V)_i$. We also set

$$N(V) = \{i \in \mathbf{Z}_{>0}; R(V)_i \neq 0\}$$

where $\mathbf{Z}_{>0}$ is the set of positive integers. This is a semigroup.

DEFINITION 1.1. The Kodaira dimension of V is

$$\kappa(V) = \begin{cases} \max_{i \in N(V)} \dim \text{Proj } R(V)^{[i]} & \text{if } N(V) \neq \emptyset \\ -\infty & \text{if } N(V) = \emptyset. \end{cases}$$

The reader is referred to Ueno [U] for a general discussion of the Kodaira dimension. The proofs of the following theorems can be found in Iitaka [I₁] and [U].

THEOREM 1.2 (The Iitaka estimate). *There exist positive integers m_0, d and positive real numbers α, β such that the following inequality holds for any integer $m \geq m_0$:*

$$\alpha m^{\kappa(V)} \leq \dim_C H^0(V, \omega_V^{\otimes dm}) \leq \beta m^{\kappa(V)}.$$

THEOREM 1.3 (The Iitaka fibration). *Suppose $N(V) \neq \emptyset$. Then there exists an integer m_0 such that for any $m \geq m_0$ satisfying $m \in N(V)$, the m -th canonical rational map*

$$\varphi_m: V \dashrightarrow W = \text{Proj } R(V)^{[m]}$$

has the following properties: Let $\varphi: V^ \rightarrow W$ be an elimination of the points of indeterminacy of the rational map φ_m . Then we have*

- (a) $\dim W = \kappa(V)$.
- (b) *The general fibers of φ are connected.*
- (c) *There is a countable intersection W' of nonempty open sets of W such that for any $w \in W'$, the fiber $V_w = \varphi^{-1}(w)$ is irreducible, non-singular and $\kappa(V_w) = 0$.*

We call $\varphi: V^* \rightarrow W$ the Iitaka fibration of V and W' an Iitaka sub-

set of W .

THEOREM 1.4. *Let $\psi: V \rightarrow W$ be a fiber space. Then there exists a nonempty open set U of W such that we have an inequality*

$$\kappa(V) \leq \kappa(V_x) + \dim W,$$

where $V_x = \psi^{-1}(x)$, for every point $x \in U$.

We remark that W may be singular in this theorem.

Next we introduce the notion of the relative Kodaira dimension, which is a generalization of the Kodaira dimension to the relative case. This was originally introduced by Lieberman-Sernesi [LS]. Note that this definition differs from the more recent one of Viehweg [V].

Let $f: X \rightarrow Y$ be a family of algebraic manifolds, and $\omega_{X/Y}$ be its relative canonical sheaf. We set

$$N(X/Y) = \{i \in \mathbf{Z}_{>0} ; f_*\omega_{X/Y}^{\otimes i} \neq 0\}$$

which is a semigroup.

If $i \in N(X/Y)$, a rational map $g_i: X \rightarrow P(f_*\omega_{X/Y}^{\otimes i})$ over Y is defined by the homomorphism $f^*f_*\omega_{X/Y}^{\otimes i} \rightarrow \omega_{X/Y}^{\otimes i}$, where $P(f_*\omega_{X/Y}^{\otimes i})$ is the projective fiber space over Y associated with the coherent sheaf $f_*\omega_{X/Y}^{\otimes i}$ (see [EGA, Chapter 2]). Z_i is the image of X by g_i , i.e., the closure of $g_i(X \setminus \Sigma_{g_i})$ in $P(f_*\omega_{X/Y}^{\otimes i})$, where Σ_{g_i} is the indeterminacy of g_i . We denote by π_i the restriction to Z_i of the projection $P(f_*\omega_{X/Y}^{\otimes i}) \rightarrow Y$. Hence we get the following diagram:

$$(1.1) \quad \begin{array}{ccc} X & \xrightarrow{g_i} & Z_i \subset P(f_*\omega_{X/Y}^{\otimes i}) \\ f \downarrow & \swarrow \pi_i & \\ & & Y \end{array}$$

DEFINITION 1.5. The relative Kodaira dimension $\kappa(X/Y)$ of $f: X \rightarrow Y$ is

$$\kappa(X/Y) = \begin{cases} \max_{i \in N(X/Y)} \dim Z_i - \dim Y & \text{if } N(X/Y) \neq \emptyset \\ -\infty & \text{if } N(X/Y) = \emptyset. \end{cases}$$

Now we slightly generalize the above definition.

DEFINITION 1.6. Let $f: X \rightarrow Y$ be a generically smooth surjective morphism of algebraic varieties, i.e., f is smooth over a certain open set Y' of Y and assume the fibers over Y' are connected. Then the relative Kodaira dimension $\kappa(X/Y)$ of $f: X \rightarrow Y$ is defined by


$$\kappa(X/Y) = \kappa(f^{-1}(Y')/Y').$$

This is obviously independent of the choice of Y' .

REMARK 1.7. Let $f: X \rightarrow Y$ be as above and assume further that f is a Gorenstein morphism. Let $\omega_{X/Y}$ be its dualizing sheaf. Then the above definition of $\kappa(X/Y)$ coincides with the relative $\omega_{X/Y}$ -dimension of [LS].

2. **The properties of the relative Kodaira dimension.** First, we observe the asymptotic behavior of the diagram (1.1) with respect to m . The following lemma and its proof is essentially analogous to that in [U, p. 54].

LEMMA 2.1. *There is a positive integer m_0 such that, for any m satisfying $m \geq m_0$ and $m \in N(X/Y)$, there exists a birational map $h_{m,m_0}: Z_m \rightarrow Z_{m_0}$ which makes up a commutative diagram*



on the locus where all the rational maps are defined.

PROOF. As the problem is local on Y , we can assume $Y = \text{Spec } A$ for a C -algebra A . We put

$$P = \bigoplus_{i \in \mathbb{Z}_{\geq 0}} P_i, \quad P_i = \Gamma(Y, f_* \omega_{X/Y}^{\otimes i}).$$

This is a graded A -algebra. We set $P^{[i]} = A[P_i]$ which is the graded A -subalgebra of P generated by P_i . Then we have

$$Z_i = \text{Proj } P^{[i]}$$

and the rational function field $C(Z_i)$ of Z_i is


$$C(Z_i) = (S_{P^{[i]}}^{-1} P^{[i]})_0$$

which is the degree 0 part of the graded quotient ring of $P^{[i]}$ with respect to the multiplicatively closed set $S_{P^{[i]}} = \bigcup_{j \geq 0} (P_j^{[i]} \setminus \{0\})$.

Let $d > 0$ be the greatest common divisor of the integers belonging to $N(X/Y)$. As $N(X/Y)$ is a semigroup, there is a positive integer l_0 such that $l \geq l_0$ implies $ld \in N(X/Y)$. We set $k = l_0 d$. Now we take a nonzero element $\varphi \in P_k$. For any positive integers n and t , the homomorphism

(2.1)
$$P_{nk} \rightarrow P_{(n+t)k}$$

sending ψ to $\varphi^t \psi$ induces a homomorphism $P^{[nk]} \rightarrow P^{[(n+t)k]}$. This induces a rational map $h_{(n+t)k, nk}: Z_{(n+t)k} \dashrightarrow Z_{nk}$ which makes up a commutative diagram



by definition. Further, the inclusion map

$$(S_{P^{[nk]}}^{-1} P^{[nk]})_0 \hookrightarrow (S_{P^{[(n+1)k]}}^{-1} P^{[(n+1)k]})_0$$

is induced by (2.1) with $t = 1$. So there is a chain of function fields

$$C(Y) \subset C(Z_k) \subset \dots \subset C(Z_{nk}) \subset C(Z_{(n+1)k}) \subset \dots \subset C(X).$$

The field extension $C(X)$ over $C(Y)$ is finitely generated. Thus there exists a positive integer n_0 such that

$$C(Z_{n_0 k}) = C(Z_{(n_0+1)k}) = C(Z_{(n_0+2)k}) = \dots$$

Now we set $m_0 = (n_0 + 1)k$. Then we can easily check that for any $m \geq m_0$ satisfying $m \in N(X/Y)$, $C(Z_m) = C(Z_{m_0})$ holds. Hence h_{m, m_0} is a birational map. q.e.d.

We immediately get the following:

COROLLARY 2.2. *There is a positive integer m_0 such that, for any $m \geq m_0$ satisfying $m \in N(X/Y)$, we have*

$$\dim Z_m - \dim Y = \kappa(X/Y).$$

Next for later use, we review the following consequence of Grothendieck's base change theorem.

THEOREM 2.3. *Let $f: X \rightarrow Y$ be a proper morphism of (reduced connected) algebraic varieties, F be a coherent sheaf on X which is \mathcal{O}_Y -flat, and q be any nonnegative integer. Then,*

- (1) *The function*

$$d_q: Y \rightarrow Z$$

$$\begin{matrix} \omega & & \omega \\ y \mapsto \dim_c H^q(X_y, F_y) \end{matrix}$$

is upper-semicontinuous on Y , where $F_y = F \otimes_{\mathcal{O}_Y} C_y$ and C_y is the residue field of the local ring $\mathcal{O}_{Y,y}$. Especially, the subset U of Y on which d_q takes the smallest value is open.

(2) The following conditions are equivalent:

(a) d_q is a constant function.

(b) $R^q f_* F$ is locally free on Y and the natural map

$$R^q f_* F \otimes_{\mathcal{O}_Y} C_y \rightarrow H^q(X_y, F_y)$$

is an isomorphism for any $y \in Y$.

For the proof of this theorem, see for instance Mumford [M, p. 50].

We study the relation between the relative Kodaira dimension and the Kodaira dimension of its fibers. The following theorem is a special case of Lieberman-Sernesi's theorem [LS, p. 83], although there seems to be a gap in their proof [LS, p. 84]. Hence we give here the proof of the required special case.

THEOREM 2.4. *Let $f: X \rightarrow Y$ be a family of algebraic manifolds. Then for any $y \in Y$, we have*

$$\kappa(X_y) \geq \kappa(X/Y).$$

Moreover, set

$$LS(X/Y) = \{y \in Y; \kappa(X_y) = \kappa(X/Y)\}.$$

Then $LS(X/Y)$ is a countable intersection of nonempty open sets of Y .

Before the proof of the above theorem, we give a lemma.

LEMMA 2.5. *We consider the following commutative diagram of algebraic varieties X, Y and Z :*

$$\begin{array}{ccc} X & \overset{g}{\dashrightarrow} & Z \\ f \downarrow & & \swarrow \pi \\ & & Y \end{array}$$

where f and π are surjective morphisms and g is a generically surjective rational map. Let Σ be the indeterminacy of g . Then there exists a nonempty open set Y' of Y such that (1) π is flat over Y' and (2) for any $y \in Y'$, Σ does not contain $X_y = f^{-1}(y)$ and the closure of $g(X_y \setminus X_y \cap \Sigma)$ coincides with $Z_y = \pi^{-1}(y)$.

PROOF. We put $U = X \setminus \Sigma$ and consider the dominant morphism $g: U \rightarrow Z$. There is an open set Z' of Z such that $Z' \subset g(U)$. Let A_1, A_2, \dots, A_k be the reduced models of the irreducible components of $Z \setminus Z'$. We consider $\pi|_{A_i}: A_i \rightarrow \pi(A_i) \subset Y$ for each i ($1 \leq i \leq k$), and put

$$Y_i = \{y \in Y; \dim(\pi|_{A_i})^{-1}(y) < \dim Z - \dim Y\}$$

which is clearly a nonempty open set of Y . Now we take an open set Y'' of Y such that π is flat over Y'' , and put

$$Y' = Y'' \cap \left(\bigcap_{i=1}^k Y_i \right).$$

This is also a nonempty open set of Y . Since the dimension of any irreducible component of Z_y is equal to $\dim Z - \dim Y$ for any $y \in Y''$, the closure of $Z'_y = (\pi|_{Z'})^{-1}(y)$ coincides with Z_y for any $y \in Y'$. q.e.d.

PROOF OF THEOREM 2.4. *Step 1.* Let Y^* be the set of points $y \in Y$ such that $\dim H^0(X_y, \omega_{X_y}^{\otimes m})$ is minimal for each $m \in \mathbf{Z}_{>0}$. Then by the base change theorem, Y^* is a countable intersection of nonempty open sets of Y . Furthermore,

$$f_* \omega_{X/Y}^{\otimes m} \otimes_{\mathcal{O}_Y} \mathcal{C}_y \cong H^0(X_y, \omega_{X_y}^{\otimes m})$$

holds for any $y \in Y^*$ and any $m \in \mathbf{Z}_{>0}$. Then we first claim

$$\kappa(X_y) = \kappa(X/Y)$$

for any $y \in Y^*$.

When $\kappa(X/Y) = -\infty$, the assertion is clear. Thus we may assume $\kappa(X/Y) \geq 0$. By Lemma 2.5, there exists a point $y \in Y^*$ such that for any $m \in \mathbf{N}(X/Y)$, the following conditions hold.

(1) The closure of $g_m(X_{y_0} \setminus X_{y_0} \cap \Sigma_{g_m})$ coincides with $\pi_m^{-1}(y_0)$.

(2) $\dim \pi_m^{-1}(y_0) = \dim Z_m - \dim Y$

where the notations are as in (1.1). Thus by letting m sufficiently large, we obtain

$$\kappa(X_{y_0}) = \dim \text{Proj } R(X_{y_0})^{[m]} = \pi_m^{-1}(y_0) = \dim Z_m - \dim Y = \kappa(X/Y).$$

Next we take any point $y \in Y^*$. The Iitaka estimates with respect to X_y and X_{y_0} say that there are positive real numbers $\alpha, \alpha', \beta, \beta'$ and a positive integer d such that for any sufficiently large integer m , we have

$$\begin{aligned} \alpha m^{\kappa(X_y)} &\leq \dim H^0(X_y, \omega_{X_y}^{\otimes dm}) \leq \beta m^{\kappa(X_y)} \\ \alpha' m^{\kappa(X_{y_0})} &\leq \dim H^0(X_{y_0}, \omega_{X_{y_0}}^{\otimes dm}) \leq \beta' m^{\kappa(X_{y_0})}. \end{aligned}$$

On the other hand, by the definition of Y^* ,

$$\dim H^0(X_y, \omega_{X_y}^{\otimes m}) = \dim H^0(X_{y_0}, \omega_{X_{y_0}}^{\otimes m})$$

holds for any $m \in \mathbf{Z}_{>0}$. Thus we have

$$\kappa(X_y) = \kappa(X_{y_0}).$$

Hence combining this with the previous equality, we obtain

$$\kappa(X_y) = \kappa(X/Y) \text{ for any } y \in Y^*.$$

Step 2. We secondly claim $\kappa(X_y) \geq \kappa(X/Y)$ for any $y \in Y \setminus Y^*$. By the definition of Y^* ,

$$\dim H^0(X_y, \omega_{X_y}^{\otimes m}) \geq \dim H^0(X_{y_0}, \omega_{X_{y_0}}^{\otimes m})$$

is satisfied for any $y \in Y \setminus Y^*$, $y_0 \in Y^*$ and any $m \in \mathbf{Z}_{>0}$. Thus by the same argument using the Iitaka estimates with respect to X_y and X_{y_0} , we obtain

$$\kappa(X_y) \geq \kappa(X_{y_0}) = \kappa(X/Y).$$

Consequently, we have the first assertion of the theorem.

Step 3. Now we prove the latter statement of the theorem by induction on the dimension of the base space Y . If $\dim Y = 0$, the assertion is obvious. Next suppose the statement is true if the dimension of the base space is smaller than n , and consider the case of $\dim Y = n > 0$. By Step 1, there exists a countable intersection Y^* of nonempty open sets of Y such that

$$\kappa(X_y) = \kappa(X/Y)$$

is satisfied for $y \in Y^*$. $Y \setminus Y^*$ can be written set-theoretically as the union of countably many irreducible reduced lower dimensional closed subvarieties W_i ($i \in I$) for a countable set I . Then the base change

$$f_i : X \times_Y W_i \rightarrow W_i$$

is a family of algebraic manifolds with $\dim W_i < n$. Thus by the induction assumption, the set

$$\mathbf{LS}_{f_i}(W_i) = \{w \in W_i; \kappa(X_w) = \kappa(X \times_Y W_i/W_i)\}$$

is a countable intersection of nonempty open sets of W_i for each i . Let I' be the subset of I consisting of $i \in I$ with $\kappa(X \times_Y W_i/W_i) = \kappa(X/Y)$. Then we obtain

$$\mathbf{LS}(X/Y) = Y^* \cup \left\{ \bigcup_{i \in I'} \mathbf{LS}_{f_i}(W_i) \right\} = Y \setminus \left\{ \left(\bigcup_{i \in I \setminus I'} W_i \right) \cup \left[\bigcup_{j \in I'} \{W_j \setminus \mathbf{LS}_{f_j}(W_j)\} \right] \right\}.$$

This is a countable intersection of nonempty open sets of Y . q.e.d.

For later use, we slightly generalize the definition of $\mathbf{LS}(X/Y)$.

DEFINITION 2.6. Let $f: X \rightarrow Y$ be a generically smooth surjective morphism of algebraic varieties of relative dimension n such that the general fibers of f are connected. Define the Lieberman-Sernesi set $\text{LS}(X/Y)$, or denoted $\text{LS}_f(Y)$, of Y with respect to f to be the set of points $y \in Y$ such that the fiber X_y is irreducible, nonsingular, n -dimensional and $\kappa(X_y) = \kappa(X/Y)$.

Now we come back to the diagram (1.1). We set m sufficiently large, set $Z = Z_m$, $g = g_m$, $\pi = \pi_m$ and consider an elimination of the points of indeterminacy of g ,

$$(2.3) \quad \begin{array}{ccc} X^\# & & \\ \tau \downarrow & \searrow h & \\ X & \dashrightarrow & Z \\ f \downarrow & \nearrow \pi & \\ Y & & \end{array}$$

where $X^\#$ is nonsingular and h is the induced surjective morphism (see [H]). In this situation,

$$\dim Z - \dim Y = \kappa(X/Y)$$

is satisfied. Let

$$(2.2) \quad \begin{array}{ccc} X_y^\# & & \\ \tau_y \downarrow & \searrow h_y & \\ X_y & \dashrightarrow & Z_y \\ f_y \downarrow & \nearrow \pi_y & \\ \{y\} & & \end{array}$$

be the restriction of (2.2) over $y \in Y$, where $X_y^\# = (f \cdot \tau)^{-1}(y)$, $Z_y = \pi^{-1}(y)$, $h_y = h|_{X_y}$ and so on. g_y is defined if the indeterminacy Σ_g of g does not contain X_y .

Then we have simultaneous Iitaka fibrations for general fibers as follows:

PROPOSITION 2.7. Let $f: X \rightarrow Y$ be a family of algebraic manifolds of dimension n with $1 \leq \kappa(X/Y) \leq n - 1$. And consider (2.2). Then

- (a) The fibers of h over normal points of Z are connected.
- (b) There exists a subset Y^\flat in Y which contains a countable intersection of nonempty open sets of Y such that we have $\kappa(X_y) = \kappa(X/Y)$

for any $y \in Y^\flat$. Furthermore,

$$h_y: X_y^\sharp \rightarrow Z_y$$

is the Iitaka fibration of X_y for any $y \in Y^\flat$.

PROOF. Let Y^\flat be the set of points $y \in Y$ which satisfies the following conditions.

(1) For any $i \in \mathbb{Z}_{>0}$,

$$f_* \omega_{X/Y}^{\otimes i} \otimes_{\mathcal{O}_Y} \mathcal{C}_y \cong H^0(X_y, \omega_{X_y}^{\otimes i}).$$

(2) For any $i \in N(X/Y)$ satisfying $i \geq m_0$ with m_0 as in Lemma 2.1, the birational map $h_{i,m_0}: Z_i \dashrightarrow Z_{m_0}$ induces a birational map between the fibers $\pi_i^{-1}(y)$ and $\pi_{m_0}^{-1}(y)$.

(3) $f \circ \tau$ is a smooth morphism over a neighborhood of y .

Y^\flat clearly is a countable intersection of nonempty open sets of Y . And for $y \in Y^\flat$, $\kappa(X_y) = \kappa(X/Y)$ is clear by (1) and Step 1 of the proof of Theorem 2.4. Further for any $y \in Y^\flat$, $h_y: X_y^\sharp \rightarrow Z_y$ is birationally equivalent to the m -th canonical rational map of X_y by (1) and (3). Hence by (2), h_y is the Iitaka fibration of X_y . Thus the statement (b) is true.

Next let $X^\sharp \xrightarrow{h} Z' \xrightarrow{\mu} Z$ be the Stein factorization of h . Then by (b) and Theorem 1.3 (b), μ is generically one to one. Thus by Zariski's main theorem, μ is an isomorphism over normal points of Z . This proves (a). q.e.d.

We remark that h is a generically smooth morphism by Sard's theorem applied to the nonsingular variety X^\sharp . Thus $\kappa(X^\sharp/Z)$ is well-defined.

COROLLARY 2.8. $\kappa(X/Z^\sharp) = 0$.

PROOF. Since $\pi^{-1}(Y^\flat)$ and $\text{LS}(X^\sharp/Z)$ are countable intersections of nonempty open sets of Z , there exists a $y \in Y^\flat$ such that Z_y intersect with $\text{LS}(X^\sharp/Z)$. Let Z'_y be an Iitaka subset of Z_y (see Theorem 1.3). As $Z'_y \cap \text{LS}(X/Z^\sharp)$ is clearly a countable intersection of nonempty open sets of Z_y , this is not empty. So if we take a point $z \in Z'_y \cap \text{LS}(X^\sharp/Z)$, then

$$\kappa(X^\sharp/Z) = \kappa(h^{-1}(z)) = 0. \quad \text{q.e.d.}$$

3. A graded subring of the canonical ring. For later use, we study graded rings. We fix an algebraic manifold V . Let R be the canonical ring of V and let

$$P = \bigoplus_{i \in \mathbf{Z}_{\geq 0}} P_i, \quad (\text{where } P_0 = C)$$

be a graded C -subalgebra of R . We also denote $P^{[i]} = C[P_i]$ in the same way as in § 1. We set

$$N(P) = \{i \in \mathbf{Z}_{>0}; P_i \neq 0\}.$$

This is a semigroup.

DEFINITION 3.1. The number $\kappa(P)$ is defined by

$$\kappa(P) = \begin{cases} \max_{i \in N(P)} \dim \text{Proj } P^{[i]} & \text{if } N(P) \neq \emptyset \\ -\infty & \text{if } N(P) = \emptyset. \end{cases}$$

REMARK 3.2. If $P = R$ itself, $\kappa(P)$ is equal to the Kodaira dimension of V .

Now we study the properties of $\kappa(P)$.

LEMMA 3.3. *If $N(P) \neq \emptyset$, then*

$$\kappa(P) = \text{tr.deg}_C P - 1.$$

LEMMA 3.4. *There is a positive integer m_0 such that, for any $m \geq m_0$ satisfying $m \in N(P)$, we have*

$$\dim \text{Proj } P^{[m]} = \kappa(P).$$

The above two lemmas are easily seen in the same method as in the case of the Kodaira dimension. Our aim of this section is the following:

PROPOSITION 3.5. *Assume that there exist a non-negative integer t , positive real numbers α, β and a positive integer d such that for any sufficiently large m ,*

$$\alpha m^t \leq \dim_C P_{dm} \leq \beta m^t$$

holds. Then we have $t = \kappa(P)$.

PROOF. We first prove $t \geq \kappa(P)$. We put $k = \kappa(P)$. Let x_0, x_1, \dots, x_k be a transcendental basis of P over C . We may assume that every x_i ($0 \leq i \leq k$) belongs to P_{n_0} for some positive integer n_0 . We put

$$P^{(n_0)} = \bigoplus_{i \in \mathbf{Z}_{\geq 0}} P_{in_0},$$

where the homogeneous part $P_i^{(n_0)}$ of degree i is P_{in_0} . Then the polynomial ring $C[x_0, x_1, \dots, x_k]$ is a graded C -subalgebra of $P^{(n_0)}$. Thus we have

$$\dim_C P_{n_0 m} \geq m^k/k!.$$

Hence $t \geq k$ is clear.

It remains to prove $t \leq \kappa(P)$. We put $r = \kappa(R) - \kappa(P) = \text{tr.deg}_c R - \text{tr.deg}_c P$ and let y_1, y_2, \dots, y_r be a transcendental basis of R over P . We may assume that every y_i ($1 \leq i \leq r$) belongs to $R_{n'}$ for some n' . We put

$$R' = R^{(n')} = \bigoplus_{i \in \mathbb{Z}_{\geq 0}} R_{in'}, \quad P' = P^{(n')} = \bigoplus_{i \in \mathbb{Z}_{\geq 0}} P_{in'}.$$

Then it is clear that $P'[y_1, y_2, \dots, y_r] \subset R'$.

First we consider the graded ring $P'[y_1]$. By the grading of $P'[y_1]$, we have $\dim P'[y_1]_m = \dim P'_m + \dim P'_{m-1} + \dots + \dim P'_1 + 1$. Thus by assumption, we have $\alpha' m^{t+1} \leq \dim P'[y_1]_m$ for any sufficiently large m and some $\alpha' > 0$. Repeating the above argument r -times, we obtain

$$\alpha'' m^{t+r} \leq \dim P'[y_1, y_2, \dots, y_r]_m \leq \dim R'_m$$

for $m \gg 0$ and some $\alpha'' > 0$.

On the other hand, the Iitaka estimate of the Kodaira dimension says

$$\dim R'_{d'm} \leq \beta' m^{\kappa(R)}$$

for $m \gg 0$, some $\beta' > 0$ and $d' > 0$. Thus we have $t + r \leq \kappa(R)$ and we are done. q.e.d.

4. Families over a nonsingular curve. In §§ 4 and 5, nonsingular curves may not be complete.

In this section, we study families over a nonsingular curve. First, we give a definition.

DEFINITION 4.1. For a variety V , we say that V is the quasi-specialization of parabolic varieties if there exist a nonsingular variety M , a nonsingular curve C and a proper surjective morphism $\varphi: M \rightarrow C$ such that

(1) $\kappa(M/C) = 0$.

(2) V is equal to the associated reduced scheme of one of the irreducible components of $\varphi^{-1}(0)$ for some $0 \in C$.

REMARK 4.2. A parabolic variety V (see [U, § 11]) trivially is a quasi-specialization of parabolic varieties since the second projection $p: V \times C \rightarrow C$ satisfies the required property.

Our aim of this section is to prove the following:

THEOREM 4.3. *Let $f: X \rightarrow S$ be a family of n -dimensional algebraic manifolds over a nonsingular curve S such that $1 \leq \kappa(X/S) \leq n - 1$.*

Then for any $s \in S$, the fiber X_s has the following property: There exist a nonsingular model X_s^* of X_s , a variety T and a fiber space $\psi: X_s^* \rightarrow T$ such that

(1) $\dim T = \kappa(X/S)$.

(2) There is an open set T' of T such that, for any $t \in T'$, the fiber $\psi^{-1}(t)$ is irreducible nonsingular and is the quasi-specialization of parabolic varieties.

In the situation of Theorem 4.3, we consider the following diagram putting N sufficiently large.

$$(4.1) \quad \begin{array}{ccc} U & & \\ \downarrow & & \\ X^\# & \xrightarrow{t} & W \\ \tau \downarrow & \searrow h & \downarrow \mu \\ X & \dashrightarrow & Z \subset P(f_* \omega_{X/S}^{\otimes N}) \\ f \downarrow & \swarrow g & \\ S & & \end{array}$$

where $X^\# \xrightarrow{t} W \xrightarrow{\mu} Z$ is the Stein factorization of h , U is the pull back of the defining locus of g by τ and others are the same as in (2.2) with $Y = S$. We may assume each component of any fiber of $f \circ \tau$ is nonsingular by Hironaka's theorem.

From now on, we fix a point $s \in S$. We consider the fiber space

$$(4.2) \quad \tilde{t}_s: \tilde{X}_s^\# \rightarrow \tilde{W}_s,$$

where $\tilde{X}_s^\#$ is the strict transform of X_s with respect to τ , i.e., the closure of $U_s = U \cap X_s^\#$, \tilde{t}_s is the restriction of t to $\tilde{X}_s^\#$ and \tilde{W}_s is the image of $\tilde{X}_s^\#$ by \tilde{t}_s . As $\tilde{X}_s^\#$ is one of the irreducible components of $X_s^\#$, $\tilde{X}_s^\#$ is nonsingular.

Now as a preparation for Lemma 4.5, we review the following, which was proved by Lieberman-Sernesi [LS, p. 79].

PROPOSITION 4.4. *Let X be a variety, S be a nonsingular curve, $f: X \rightarrow S$ be a proper morphism and F be an \mathcal{O}_S -flat coherent sheaf on X . Then,*

- (1) $f_* F$ is a locally free sheaf of finite rank.
- (2) For any $s \in S$, the natural map

$$t_s^0: f_* F \otimes_{\mathcal{O}_s} C_s \rightarrow H^0(X_s, F_s)$$

is injective.

In the situation before Proposition 4.4, we have

LEMMA 4.5. $\dim \tilde{W}_s = \kappa(X/S)$.

PROOF. For any point $u \in S$, we set

$$P(X_u) = \bigoplus_{i \in \mathbb{Z}_{\geq 0}} (f_* \omega_{X/Y}^{\otimes i} \otimes_{\mathcal{O}_s} C_u).$$

This is a graded C -subalgebra of the canonical ring $R(X_u)$ by Proposition 4.4 (2).

Now we set $\tilde{Z}_s = h(\tilde{X}_s^*)$. Since by Proposition 4.4 it is easy to show that the indeterminacy Σ_g does not contain X_s , \tilde{Z}_s coincides with the closure of $g(X_s \setminus X_s \cap \Sigma_g)$. Then we have $\tilde{Z}_s = \text{Proj } P(X_s)^{[N]}$. As N is sufficiently large, we have

$$\dim \tilde{Z}_s = \kappa(P(X_s))$$

where $\kappa(P(X_s))$ is as defined in § 3.

On the other hand, for a general point $u \in S$, we have $R(X_u) = P(X_u)$ and $\kappa(X_u) = \kappa(X/S)$. As $\dim P(X_u)_i = \dim P(X_s)_i$ holds for any $i \in \mathbb{Z}_{\geq 0}$ by Proposition 4.4 (1), the Iitaka estimate with respect to X_u says

$$\alpha m^{\kappa(X/S)} \leq \dim P(X_s)_{dm} \leq \beta m^{\kappa(X/S)}$$

for some $\alpha > 0, \beta > 0, d \in \mathbb{Z}_{>0}$ and any sufficiently large m . Hence by Proposition 3.5, we have

$$\kappa(P(X_s)) = \kappa(X/S).$$

Thus we have $\dim \tilde{Z}_s = \kappa(X/S)$. As $\dim \tilde{W}_s = \dim \tilde{Z}_s$ is clear, we are done. q.e.d.

We consider (4.2). As \tilde{X}_s^* is nonsingular, there is an open set \tilde{W}'_s of \tilde{W}_s such that (1) \tilde{t}_s is a smooth morphism over \tilde{W}'_s , (2) $\tilde{W}'_s \subset t(U)$ and (3) the closure of $U_w = (t|_U)^{-1}(w)$ coincides with $(\tilde{X}_s^*)_w$ for any $w \in \tilde{W}'_s$. The assertion (3) is possible by Lemma 2.5. We take and fix any point $w \in \tilde{W}'_s$.

LEMMA 4.6. *There exists an irreducible curve C on W passing through w such that*

(a) $\text{LS}_t(W) \cap C$ is a countable intersection of nonempty open sets of C .

(b) *There is an irreducible component \tilde{X} of $t^{-1}(C)$ such that*

(1) *the induced morphism $\tilde{t} = t|_{\tilde{X}}: \tilde{X} \rightarrow C$ is surjective.*

(2) $(\tilde{X}_s^*)_w = (\tilde{t}_s)^{-1}(w)$ is one of the irreducible components of the fiber $\tilde{X}_w = \tilde{t}^{-1}(w)$.

PROOF. We take points $p \in \text{LS}_t(W) \cap t(U)$ satisfying $p \neq w, x \in t^{-1}(w) \cap U$ and $y \in t^{-1}(p) \cap U$. By a lemma in Mumford [M, p. 56], we can take an irreducible curve D on X passing through x and y . We set $C = t(D)$. Since C is a curve passing through p , the desired property (a) is clearly satisfied.

Next we consider the proper morphism $t|_{t^{-1}(C)}: t^{-1}(C) \rightarrow C$. As the general fibers of $t|_{t^{-1}(C)}$ are irreducible by (a), there exists only one irreducible component \tilde{X} of $t^{-1}(C)$ which dominates C . We set $\tilde{U} = \tilde{X} \cap U$ and consider the dominant morphism

$$t|_{\tilde{U}}: \tilde{U} \rightarrow C.$$

$t|_{\tilde{U}}$ is an equidimensional morphism of relative dimension $n - \kappa(X/S)$ since \tilde{U} is irreducible, C is a curve and by (a). Then the closure of the fiber $U_w = (t|_{\tilde{U}})^{-1}(w)$ coincides with $(\tilde{X}_s^*)_w$, because \tilde{U}_w is contained in $U_w = (t|_U)^{-1}(w)$, the closure of U_w coincides with $(\tilde{X}_s^*)_w$, $\dim \tilde{U}_w = \dim (\tilde{X}_s^*)_w = n - \kappa(X/S)$ and $(\tilde{X}_s^*)_w$ is irreducible. As the closure of \tilde{U} coincides with \tilde{X} , the closure of \tilde{U}_w is contained in the fiber \tilde{X}_w . Thus the property (b) is clearly satisfied. q.e.d.

Now we prove Theorem 4.3. For any $s \in S$, we show that the fiber space $\tilde{t}_s: \tilde{X}_s^* \rightarrow \tilde{W}_s$ has the desired property. The statement (1) was proved in Lemma 4.5. For any $w \in \tilde{W}'_s$, we consider $\tilde{t}: \tilde{X} \rightarrow C$. Let $\hat{C} \rightarrow C$ be the normalization, $\hat{X} \rightarrow \tilde{X}$ be a desingularization and $\hat{t}: \hat{X} \rightarrow \hat{C}$ be the induced fiber space. By Corollary 2.8, Lemma 4.6 (a) and our construction, we have $\kappa(\hat{X}/\hat{C}) = 0$. Thus Lemma 4.6 (b) says that $\tilde{t}_s^{-1}(w)$ is the quasi-specialization of parabolic varieties for any $w \in \tilde{W}'_s$. q.e.d.

5. Proofs of main theorems. We prove Theorem I. We take any point $y \in Y$. By a lemma in [M, p. 56], we can take an irreducible curve C on Y passing through y such that $C \cap \text{LS}(X/Y)$ is a countable intersection of nonempty open sets of C . Let $\tau: \hat{C} \rightarrow C$ be the normalization and $\hat{f}: \hat{X} = X \times_Y \hat{C} \rightarrow \hat{C}$ be the induced family. Since $\tau\{\text{LS}(\hat{X}/\hat{C})\} \cap \text{LS}(X/Y) \neq \emptyset$, we have

$$\kappa(\hat{X}/\hat{C}) = \kappa(X/Y).$$

As X_y is isomorphic to $\hat{f}^{-1}(\hat{y})$ for some $\hat{y} \in \hat{C}$, our assertion is clear by Theorem 4.3. q.e.d.

Next we formulate the lower-semicontinuity of the Kodaira dimension of a degenerating family of parabolic varieties.

CONJECTURE PDG_n. *Let $f: M \rightarrow S$ be a proper surjective morphism with connected fibers of relative dimension n from an algebraic mani-*

fold M to a nonsingular curve S . Assume that there is a set $P = \{p_1, p_2, \dots, p_r\}$ of points of S such that (1) f is a smooth morphism over $S \setminus P$. (2) For any $s \in S \setminus P$, we have $\kappa(M_s) = 0$. Let

$$M_{p_i} = \bigcup_j m_{ij} M_{p_i}^{(j)}$$

be the decomposition into irreducible components of the fiber $M_{p_i} = f^{-1}(p_i)$ for $p_i \in P$. Then we have

$$\kappa(M_{p_i}^{(j)}) \leq 0$$

for any $p_i \in P$ and j .

REMARK. When $n = 1$, Conjecture PDG_1 is true because any component of a degenerate fiber of a degenerating family of elliptic curves is elliptic or rational (see Kodaira [K]).

When $n = 2$, every known example of degenerating families of surfaces of parabolic type satisfies the statement of Conjecture PDG_2 (see for example Persson [P]).

We prove Theorem II. For any $y \in Y$, let $\psi: X_y^* \rightarrow T$ be the fibration defined in Theorem I. For a point $t \in T$, let $\varphi: M \rightarrow C$ be the fiber space which induces on $\psi^{-1}(t)$ the quasi-specialization of parabolic varieties. Then Conjectures $\text{DF}_{n-k,0}$ and PDG_{n-k} clearly imply $\kappa(\psi^{-1}(t)) \leq 0$. Thus by Theorem 1.4, we have

$$\kappa(X_y) = \kappa(X_y^*) \leq \kappa(\psi^{-1}(t)) + \dim T \leq \kappa(X/Y).$$

On the other hand, $\kappa(X_y) \geq \kappa(X/Y)$ is also true by Theorem 2.4. Hence we obtain $\kappa(X_y) = \kappa(X/Y)$. q.e.d.

Since Conjectures $\text{DF}_{1,0}$ and PDG_1 are true, we have:

COROLLARY. Conjecture $\text{DF}_{n,n-1}$ is true.

REFERENCES

- [H] H. HIRONAKA, Resolution of singularities of an algebraic variety over a field of characteristic zero, I, II, Ann. of Math. 79 (1964), 109-326.
- [I₁] S. ITAKA, On D -dimensions of algebraic varieties, J. Math. Soc. Japan 23 (1971), 356-373.
- [I₂] S. ITAKA, Deformation of compact complex surfaces, I, II, and III, in Global analysis, papers in honor of K. Kodaira, Princeton Univ. Press and Univ. of Tokyo Press (1969), 267-272; J. Math. Soc. Japan 22 (1970), 247-261 and *ibid.* 23 (1971), 692-705.
- [K] K. KODAIRA, On compact analytic surfaces II, Ann. of Math. 77 (1963), 563-626.
- [LS] D. LIEBERMAN AND E. SERNESI, Semicontinuity of L -dimension, Math. Ann. 225 (1977), 77-88.
- [M] D. MUMFORD, Abelian Varieties, Oxford Univ. Press, London, 1970.
- [N] I. NAKAMURA, Complex parallelisable manifolds and their small deformations, J. Diff. Geometry 10 (1975), 85-112.

- [P] U. PERSSON, On degenerations of algebraic surfaces, *Memoirs of Amer. Math. Soc.* 189, 1977.
- [U] K. UENO, *Classification Theory of Algebraic Varieties and Compact Complex Spaces*, Lecture Notes in Math. 439, Springer-Verlag, Berlin-New York-Heidelberg, 1975.
- [EGA] A. GROTHENDIECK AND J. DIEUDONNÉ, *Eléments de géométrie algébrique*, *Publ. Math. I.H.E.S.*, 4, 8, 11, 17.
- [V] E. VIEHWEG, *Klassifikationstheorie algebraischer Varietäten der Dimension drei*, *Compositio Math.* 41 (1980), 361-400.

MIYAGI TECHNICAL COLLEGE
NATORI, 981-12
JAPAN

Appendix

KENJI UENO

(Received September 9, 1980)

In this appendix we shall show that PDG_2 is true. Since $\text{DF}_{2,0}$ is true by virtue of the classification of algebraic surfaces, Theorem II implies the validity of $\text{DF}_{n,n-2}$.

The conjecture PDG_2 is true by the following.

PROPOSITION. *Let $f: M \rightarrow D$ be a proper surjective morphism of a three dimensional complex manifold M to a disk $D = \{z \in \mathbf{C} \mid |z| < \varepsilon\}$ with connected fibers. Assume that f is smooth at each point on $f^{-1}(D^*)$, $D^* = D - \{0\}$ and f is projective. Then for each irreducible component E of the fiber $f^{-1}(0)$, we have*

$$\kappa(E) \leq \kappa(M_x),$$

where $M_x = f^{-1}(x)$ is a general fiber of f .

Note that κ is invariant under smooth deformations for surfaces. The above proposition was proved by Persson [P] except a few cases. By virtue of Kulikov [Ku], Persson-Pinkham [P²], we can prove the proposition in these cases. For reader's convenience, we give a detailed proof.

PROOF OF PROPOSITION. If $\kappa(M_x) = 2$, the proposition is trivially true. Therefore, first assume $\kappa(M_x) = 1$, that is, M_x is an elliptic surface of general type. Then, all fibers M_y , $y \in D^*$ are elliptic surfaces of general type and there is a positive integer m such that the m -canonical mapping $\Phi_{mK}: M_y \rightarrow C_y = \Phi_{mK}(M_y)$ associated with the m -canonical system $|mK_{M_y}|$ of M_y is a morphism and gives the structure of an elliptic surface for each M_y , $y \in D^*$. (See [K] and [I₂].) Let

$$\begin{array}{ccc}
 M & \overset{g}{\dashrightarrow} & Z \subset P(f_* \omega_{M/D}^{\otimes m}) \\
 \downarrow f & \swarrow & \\
 D & &
 \end{array}$$

be a diagram similar to that in § 1. The meromorphic mapping g is holomorphic on $M^* = f^{-1}(D^*)$. Let $h: \widehat{M} \rightarrow M$ be obtained by a succession of monoidal transformations along non-singular centers such that $\widehat{g} = g \circ h: \widehat{M} \rightarrow Z$ is a morphism. If the proposition is true for the family $\widehat{f} = f \circ h: \widehat{M} \rightarrow D$, then it is also true for the original family $f: M \rightarrow D$. Hence, we can assume that g is a morphism. For the same reason, we can assume that Z is a two dimensional complex manifold. By our construction, every regular fiber of g is a non-singular elliptic curve. Let us consider an irreducible component E of the fiber $f^{-1}(0)$. By taking a finite succession of monoidal transformations of M and Z along non-singular centers, we can assume that E is non-singular and $g(E)$ is a non-singular curve on Z . Since general fibers of g are connected, every fiber of $g|_E: E \rightarrow g(E)$ is connected. Let $p \in g(E)$ be a general smooth analytic curve in Z passing through p such that $g^{-1}(C)$ is non-singular. Then each general fiber of $g_C: g^{-1}(C) \rightarrow C$ is a non-singular elliptic curve. Since $g|_E^{-1}(p)$ is contained in $g_C^{-1}(p)$ and any irreducible component of $g_C^{-1}(p)$ is an elliptic curve or a rational curve, a general fiber of $g|_E: E \rightarrow g(C)$ is an elliptic curve or a rational curve. Hence, by Theorem 1.4, we have

$$\kappa(E) \leq 1.$$

This is the desired result.

Note that in the proof we need not assume that f is projective so that the proposition is valid for any fibration $f: M \rightarrow D$ with $\kappa(M_x) = 1$.

Next we consider the case in which $\kappa(M_x) = 0$. First we will show that we can assume that all fibers over D^* are minimal.

Let ω be a non-zero element in $H^0(M, \omega_{M/D}^{\otimes 12})$. (Since $f_* \omega_{M/D}^{\otimes 12}$ is torsion free, hence free over D , we can always find such an element.) If a regular fiber M_x contains an exceptional curve of the first kind, by [W, Proposition 2.4, p. 291], all regular fibers over D^* contain exceptional curves of the first kind. Moreover, all exceptional curves of the first kind on each regular fiber do not intersect each other. Therefore, if necessary, taking a finite ramified covering \widehat{D} of D ramified at the origin and a non-singular model \widehat{M} of $M \times_D \widehat{D}$, we may assume that there exists an irreducible divisor E appearing in the divisor (ω) such that the intersection $E \cap M_x$ is an exceptional curve of the first kind on

each fiber $M_x, x \in D^*$. (Note that by the argument given below if the proposition is valid for $\hat{f}: \hat{M} \rightarrow D$, it is also valid for our original family $f: M \rightarrow D$.) Let F be a relatively ample divisor of $f: M \rightarrow D$. We let m be the intersection number $(F \cap M_x) \cdot (E \cap M_x)$ on M_x . Let us consider an invertible sheaf $O_M(n(F + mE))$ for a sufficiently large positive integer n and consider a diagram

$$\begin{array}{ccc}
 M & \xrightarrow{h} & M' \subset P(f_* O_M(n(F + mE))) \\
 & \searrow f & \downarrow f' \\
 & & M
 \end{array}$$

Then h is a morphism on $M^* = f^{-1}(D^*)$, $h_y: M_y \rightarrow M'_y, y \in D^*$ is a contraction morphism (that is $h_y(E \cap M_y)$ is a point) and M' is smooth over D^* . We let M_1 be a nonsingular model of M' obtained by a finite succession of monoidal transformations along non-singular centers contained in the fibers over the origin. Since h is bimeromorphic, the proposition is true for the family $f: M \rightarrow D$ if and only if it is true for $f_1: M_1 \rightarrow D$. Applying this process finitely many times, we can assume that all regular fibers of $f: M \rightarrow D$ do not contain exceptional curves of the first kind.

Next we will show that it is enough to consider the case in which $p_g(M_x) = 1$ for a regular fiber. Assume $p_g(M_x) = 0$. Then there exists a positive integer $m \geq 2$ which is a divisor of 12 such that $P_m(M_x) = 1, P_l(M_x) = 0, l = 1, 2, \dots, m - 1$. Then, using a divisor (ω) for a non-zero element $\omega \in H^0(M, \omega_{M/D}^{\otimes m})$, we can construct an m -sheeted covering $g: \hat{M} \rightarrow M$ which may ramify along divisors contained in the fiber over the origin. (See, for example, [U, p. 176-177].) Note that $g_y: \hat{M}_y \rightarrow M_y$ is an m -sheeted covering with $p_g(\hat{M}_y) = 1$ for each $y \in D^*$. Let \tilde{M} be a non-singular model of \hat{M} obtained by a finite succession of monoidal transformations along non-singular centers contained in the fiber over the origin. Then for an irreducible component E of $f^{-1}(0)$, there is an irreducible component \tilde{E} of the fiber over the origin of $\tilde{f}: \tilde{M} \rightarrow D$ such that $\tilde{E} \rightarrow E$ is a generically finite morphism. Since $\kappa(\tilde{E}) \geq \kappa(E)$, it is enough to prove the proposition under the assumption that each regular fiber is a minimal surface with $\kappa = 0, p_g = 1$, hence an abelian surface or a $K3$ surface.

Now by [N] (for a family of abelian surfaces), [K] and [P²] (for a family of $K3$ surfaces), there exists a finite ramified covering $\hat{D} \rightarrow D$ ramified along the origin and a fibration $\hat{f}: \hat{M} \rightarrow \hat{D}$ which is bimeromorphically equivalent to $M \times_D \hat{D} \rightarrow \hat{D}$ such that our proposition is true for

the family $\hat{f}: \hat{M} \rightarrow \hat{D}$. Then, for the same reason as above, it is also true for our family $f: M \rightarrow D$.

Finally we consider the case in which $\kappa(M_x) = -\infty$ for a general fiber M_x . Since f is projective, M_x is a rational surface or a ruled surface. Therefore the surface M_x contains a rational curve l_x with self-intersection number $l_x^2 = 1$ or 0 in M_x . Moreover if $l_x^2 = 1$, then M_x is a rational surface. Let us consider the relative Hilbert scheme $\text{Hilb}_{M/D}$. By the stability theorem ([K₂]), the component F of $\text{Hilb}_{M/D}$ containing the point corresponding to l_x is of positive dimension. Since the natural morphism $p: F \rightarrow D$ is proper, there is a holomorphic map $\pi: E = \{t \in \mathbb{C} \mid |t| < \varepsilon'\} \rightarrow F$ such that $\pi \circ p: E \rightarrow D$ is a finite ramified covering. We may assume that $\pi \circ p$ is ramified only over the origin. Let $\hat{f}: \hat{M} \rightarrow E$ be a non-singular model of $M \times_D E \rightarrow E$ and L be the pull back of the universal family \mathcal{L} over F to E . Then the image \hat{L} of L to \hat{M} is a divisor on \hat{M} such that for a general fiber \hat{M}_y of \hat{f} , $\hat{L} \cap \hat{M}_y = \hat{l}_y$ is a non-singular rational curve with $\hat{l}_y^2 = 1$ or 0 in \hat{M}_y . Let us consider a diagram

$$\begin{array}{ccc} \hat{M} & \xrightarrow{g} & S \subset \mathbf{P}(\hat{f}_* O_{\hat{M}}(\hat{L})) \\ \hat{f} \downarrow & \swarrow & \\ \hat{E} & & \end{array}$$

For the same reason as above, we can assume that g is a morphism. If $\hat{l}_y^2 = 1$, then $\hat{f}_* O_{\hat{M}}(\hat{L})$ is a locally free sheaf of rank 3, since \hat{M}_y is rational. Moreover, in this case, it is easy to show that $S = \mathbf{P}(\hat{f}_* O_{\hat{M}}(\hat{L}))$ and g is bimeromorphic, since $\Phi_{\hat{l}_y}: \hat{M}_y \rightarrow \mathbf{P}^2$ is birational. Hence any component G of the fiber of f over the origin is a rational or ruled surface. Hence $\kappa(G) = -\infty$. This implies our proposition. Assume $\hat{l}_y^2 = 0$. Then, $\Phi_{\hat{l}_y}: \hat{M}_y \rightarrow C_y = \Phi_{L_y}(\hat{M}_y)$ gives the structure of a ruled surface, if we blow down all exceptional curves contained in fibers of $\Phi_{\hat{l}_y}$. Hence, in this case, S is a surface and general fibers of g are \mathbf{P}^1 's. Hence by an argument similar to that in the case of $\kappa(M_x) = 1$, we conclude that each component of the fibers of f over the origin is rational or ruled. Thus we obtain the desired result.

REFERENCES

[K₂] K. KODAIRA, On stability of compact submanifolds of complex manifolds, Amer. J. Math. 85 (1963), 79-94.
 [Ku] V. S. KULIKOV, Degenerations of $K 3$ surfaces and Enriques surfaces, Math. USSR Izvestija 11 (1977), 957-989.
 [N] Y. NAMIKAWA, Toroidal degeneration of abelian varieties, I, II, in Complex Analysis

and Algebraic Geometry, Iwanami Shoten and Cambridge Univ. Press 1977, 227-237,
and Math. Ann. 245 (1979), 117-150.

- [P²] U. PERSSON AND PINKHAM, Degeneration of surfaces with trivial canonical bundle, Ann.
of Math. 113 (1981), 45-66.
- [W] P.M.H. WILSON, The behavior of the plurigenera of surfaces under algebraic smooth
deformations, Inventiones Math. 47 (1978), 289-299.

DEPARTMENT OF MATHEMATICS
KYOTO UNIVERSITY
KYOTO, 606
JAPAN