

The $D(2)$ –problem for dihedral groups of order $4n$

SEAMUS O’SHEA

We give a full solution in terms of k –invariants of the $D(2)$ –problem for D_{4n} , assuming that $\mathcal{Z}[D_{4n}]$ satisfies torsion-free cancellation.

57M05; 55P15

1 Introduction

The following question was first posed by Wall in [12]:

$D(2)$ –problem. Let X be a finite connected 3–dimensional CW–complex, with universal cover \tilde{X} , such that

$$H_3(\tilde{X}; \mathcal{Z}) = H^3(X; \mathcal{B}) = 0$$

for all coefficient systems \mathcal{B} on X . Is it true that X is homotopy equivalent to a finite 2–dimensional CW–complex?

The $D(2)$ –problem is parametrized by the fundamental group of X ; we say that the $D(2)$ –property holds for a finitely presented group G if the above question is answered in the affirmative for every X with $\pi_1(X) \cong G$.

We shall be concerned with the $D(2)$ –problem for D_{4n} , the dihedral group of order $4n$. Johnson [7] has shown that the $D(2)$ –property holds for the groups D_{4n+2} for any $n \geq 1$; however his result relies on the fact that D_{4n+2} has periodic cohomology, a property not shared by D_{4n} . Mannan [9] has shown that the $D(2)$ –property holds for D_8 . We say that *torsion-free cancellation* holds for a group ring $\mathcal{Z}[G]$ if

$$X \oplus M \cong X \oplus N \Rightarrow M \cong N$$

for any $\mathcal{Z}[G]$ –lattices X , M and N . We shall show:

Theorem 1.1 *Suppose that $\mathcal{Z}[D_{4n}]$ satisfies torsion-free cancellation. Then the $D(2)$ –property holds for D_{4n} .*

The calculations of Swan [11] and Endo and Miyata [3] show that torsion-free cancellation holds for $\mathbf{Z}[D_{4p}]$ when p is prime and $3 \leq p \leq 31$, $p = 47, 179$ or 19379 . To date the only finite nonabelian, nonperiodic groups for which the $D(2)$ -property is known to hold are those of the form D_{4p} , where p is prime.

Let G be a group and set $\Lambda = \mathbf{Z}[G]$. Any finite 2-dimensional CW-complex K with $\pi_1(K) = G$ gives rise to an exact sequence of Λ -modules

$$(1) \quad 0 \rightarrow \pi_2(K) \rightarrow C_2(K) \xrightarrow{\partial_2} C_1(K) \xrightarrow{\partial_1} C_0(K) \rightarrow \mathbf{Z} \rightarrow 0,$$

where $C_r(K) = H_r(\tilde{K}_r, \tilde{K}_{r-1}; \mathbf{Z})$ is the free Λ -module with basis the r -cells of K . By an algebraic 2-complex over a group G , we mean an exact sequence of right Λ -modules of the form

$$(2) \quad 0 \rightarrow J \rightarrow F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\varepsilon} \mathbf{Z} \rightarrow 0,$$

where each F_i is finitely generated free. An algebraic 2-complex is said to be geometrically realizable if it is homotopy equivalent to a 2-complex of type (1). If every algebraic 2-complex over a group G is geometrically realizable we say that the realization property holds for G . The following result is due to Johnson [7] and Mannan [10]:

Theorem 1.2 *Let G be a finitely presented group. Then the $D(2)$ -property holds for G if and only if the realization property holds for G .*

We are grateful to the referee for pointing out a paper of Latiolais [8], in which it is proved that the homotopy type of a CW-complex with fundamental group D_{4n} is determined by the Euler characteristic. This result was extended by Hambleton and Kreck [6] to include those complexes whose fundamental groups are finite subgroups of $SO(3)$. Latiolais achieves this by realizing all values of the Browning obstruction group (see Browning [1], Gruenberg [4], Gutierrez and Latiolais [5]); combining this realization with Theorem 1.2, it seems possible to give a proof of Theorem 1.1 without assuming torsion-free cancellation.

We begin by briefly recalling the classification of algebraic complexes in terms of k -invariants — for a full treatment, see Johnson [7, Chapter 6]. Fix a finite group G and put $\Lambda = \mathbf{Z}[G]$. Let $\mathcal{P} = (0 \rightarrow J \rightarrow F_2 \rightarrow F_1 \rightarrow F_0 \rightarrow \mathbf{Z} \rightarrow 0)$ be an algebraic 2-complex over G and let $\mathcal{E} = (0 \rightarrow J \rightarrow E_2 \rightarrow E_1 \rightarrow E_0 \rightarrow \mathbf{Z} \rightarrow 0) \in \text{Ext}_{\Lambda}^3(\mathbf{Z}, J)$ be an arbitrary extension of \mathbf{Z} by J . Then by the universal property of projective

modules, there exists a commutative diagram:

$$\begin{array}{ccccccccccc}
 \mathcal{P} & = & (0 & \longrightarrow & J & \longrightarrow & F_2 & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & \mathbf{Z} & \longrightarrow & 0) \\
 \downarrow \alpha & & & & \downarrow \alpha_+ & & \downarrow \alpha_2 & & \downarrow \alpha_1 & & \downarrow \alpha_0 & & \downarrow \text{Id} & & \\
 \mathcal{E} & = & (0 & \longrightarrow & J & \longrightarrow & E_2 & \longrightarrow & E_1 & \longrightarrow & E_0 & \longrightarrow & \mathbf{Z} & \longrightarrow & 0)
 \end{array}$$

We may extend α_+ thus:

$$\begin{array}{ccccccccccc}
 0 & \longrightarrow & J & \longrightarrow & F_2 & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & \mathbf{Z} & \longrightarrow & 0 \\
 & & \downarrow \alpha_+ & & \downarrow \alpha'_2 & & \downarrow \alpha'_1 & & \downarrow \alpha'_0 & & \downarrow \tilde{\alpha} & & \\
 0 & \longrightarrow & J & \longrightarrow & F_2 & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & \mathbf{Z} & \longrightarrow & 0
 \end{array}$$

Then $\tilde{\alpha}$ is unique up to congruence modulo $|G|$ and we have a well-defined map $\kappa: \text{End}_\Lambda J \rightarrow \mathbf{Z}/|G|$ given by $\kappa(\alpha_+) = \tilde{\alpha}$. The k -invariant of the transition $\alpha: \mathcal{P} \rightarrow \mathcal{E}$ is defined to be $k(\mathcal{P} \rightarrow \mathcal{E}) = \kappa(\alpha_+)$. Given $\alpha \in \text{End}_\Lambda J$ we have a k -invariant $k(\mathcal{P} \rightarrow \alpha_*(\mathcal{P})) = \kappa(\alpha)k(\mathcal{P} \rightarrow \mathcal{P}) = \kappa(\alpha)$, where $\alpha_*(\mathcal{P})$ is the pushout extension. Since $\kappa(\alpha)$ is a unit if α is an isomorphism, this induces a mapping

$$\text{Aut}_\Lambda J \rightarrow (\mathbf{Z}/|G|)^*$$

called the Swan map, which is independent of the choice of algebraic complex in which J appears. We have (see Johnson [7, Theorems 54.6 and 54.7]):

Theorem 1.3 *Suppose that the Swan map $\text{Aut } J \rightarrow (\mathbf{Z}/|G|)^*$ is surjective. Then for each $n \geq 0$ there is, up to chain homotopy equivalence, a unique algebraic 2-complex of the form*

$$0 \rightarrow J \oplus \Lambda^n \rightarrow F_2 \rightarrow F_1 \rightarrow F_0 \rightarrow \mathbf{Z} \rightarrow 0.$$

2 The Swan map for D_{2n}

For any n the group D_{2n} may be described by the presentation

$$\langle x, y \mid x^n, y^2, y^{-1}xyx \rangle.$$

Write $\Lambda = \mathbf{Z}[D_{2n}]$ and $\Sigma = 1 + x + x^2 + \dots + x^{n-1}$. Applying the Cayley complex construction to this presentation gives the following 2-complex:

$$(3) \quad 0 \rightarrow J \rightarrow \Lambda^3 \xrightarrow{\partial_2} \Lambda^2 \xrightarrow{\partial_1} \Lambda \xrightarrow{\varepsilon} \mathbf{Z} \rightarrow 0,$$

where ε is the augmentation map, $\partial_1 = (x - 1, y - 1)$ and $\partial_2 = \begin{pmatrix} \Sigma & 0 & 1+yx \\ 0 & 1+y & x-1 \end{pmatrix}$. The following proposition is easily verified:

Proposition 2.1 Fix n and let k be any odd integer with $3 \leq k \leq n - 1$. If we write $m = (k - 1)/2$ then the following diagram commutes:

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & J & \longrightarrow & \Lambda^3 & \xrightarrow{\partial_2} & \Lambda^2 & \xrightarrow{\partial_1} & \Lambda & \xrightarrow{\varepsilon} & \mathbf{Z} & \longrightarrow & 0 \\
 & & \downarrow \theta & & \downarrow \alpha_2 & & \downarrow \alpha_1 & & \downarrow \alpha_0 & & \downarrow k & & \\
 0 & \longrightarrow & J & \longrightarrow & \Lambda^3 & \xrightarrow{\partial_2} & \Lambda^2 & \xrightarrow{\partial_1} & \Lambda & \xrightarrow{\varepsilon} & \mathbf{Z} & \longrightarrow & 0
 \end{array}$$

where $\partial_1 = (x - 1, y - 1)$, $\partial_2 = \begin{pmatrix} \Sigma & 0 & 1+yx \\ 0 & 1+y & x-1 \end{pmatrix}$,

$$\alpha_0 = (1 + x^{-1} + \dots + x^{-m} + x^{-1}y + \dots + x^{-m}y),$$

$$\alpha_1 = \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}, \quad \alpha_2 = \begin{pmatrix} a & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

$a = 1 + x^{-1} + \dots + x^{-m} - x^{-2}y - \dots - x^{-m-1}y$ and $\theta = \alpha_2|_J$.

Consider the commutative diagram above as a diagram of (free) \mathbf{Z} -modules and \mathbf{Z} -linear maps; taking determinants we have:

Proposition 2.2 $k \det \theta \det \alpha_1 = \det \alpha_2 \det \alpha_0$.

Proof Let v denote the restriction of α_0 to $\ker \varepsilon$ and let u denote the restriction of α_1 to $\ker \partial_1$. Then $v(\ker \varepsilon) \subset \ker \varepsilon$, $u(\ker \partial_1) \subset \ker \partial_1$ and we have an commutative diagram:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \ker \partial_1 & \longrightarrow & \Lambda^2 & \xrightarrow{\partial_1} & \ker \varepsilon & \longrightarrow & 0 \\
 & & \downarrow u & & \downarrow \alpha_1 & & \downarrow v & & \\
 0 & \longrightarrow & \ker \partial_1 & \longrightarrow & \Lambda^2 & \xrightarrow{\partial_1} & \ker \varepsilon & \longrightarrow & 0
 \end{array}$$

Considered as a diagram of (free) \mathbf{Z} -modules, both exact sequences split, and so there exists α'_1 such that

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \ker \partial_1 & \longrightarrow & \ker \partial_1 \oplus \ker \varepsilon & \longrightarrow & \ker \varepsilon & \longrightarrow & 0 \\
 & & \downarrow u & & \downarrow \alpha'_1 & & \downarrow v & & \\
 0 & \longrightarrow & \ker \partial_1 & \longrightarrow & \ker \partial_1 \oplus \ker \varepsilon & \longrightarrow & \ker \varepsilon & \longrightarrow & 0
 \end{array}$$

commutes with the obvious maps, and where $\det \alpha'_1 = \det \alpha_1$. Therefore we have $\det \alpha'_1 = \det \begin{pmatrix} u & w \\ 0 & v \end{pmatrix} = \det u \det v$. Similarly

$$\det \alpha_2 = \det \theta \det u \quad \text{and} \quad \det \alpha_0 = \det v \det k = k \det v.$$

Thus

$$\det \alpha_2 \det \alpha_0 = \det \theta \det u \det(v) k = k \det \theta \det \alpha_1$$

as required. □

Now, any Λ -homomorphism is a Λ -isomorphism if and only if it is an isomorphism as a \mathbf{Z} -linear map. Thus, in order to show that $[k]$ is in the image of the Swan map, it suffices to show that $\det \theta = \pm 1$.

Proposition 2.3 *Suppose that k is coprime to $2n$. Then $\det \alpha_0 = \pm k$.*

Proof Let $M(\alpha_0)$ be the matrix of the Λ -linear map given by $x \mapsto \alpha_0 x$ with respect to the \mathbf{Z} -basis $\{1, x, \dots, x^{n-1}, y, \dots, x^{n-1}y\}$, with the elements of Λ being interpreted as columns. Notice that $M(\alpha_0) = \begin{pmatrix} A & B \\ B & A \end{pmatrix}$ where $A = (a_{i,j})$ and $B = (b_{i,j})$ are $n \times n$ matrices. We know that $a_{i,1} = 1$ if α_0 contains an x^{i-1} term and $a_{i,1} = 0$ otherwise. Thus

$$a_{i,1} = \begin{cases} 1 & \text{if } i \in \{1, n - m + 1, n - m + 2, \dots, n\}, \\ 0 & \text{otherwise.} \end{cases}$$

Similarly

$$b_{i,1} = \begin{cases} 1 & \text{if } i \in \{n - m + 1, n - m + 2, \dots, n\}, \\ 0 & \text{otherwise.} \end{cases}$$

The other columns of A and B are obtained by cyclically permuting the first column; let $\sigma_+, \sigma_-: \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ be the permutations given by $\sigma_+(i) = i + 1 \pmod n$ and $\sigma_-(i) = i - 1 \pmod n$. We now have

$$a_{i,j} = a_{\sigma_-^{j-1}(i),1} \quad \text{and} \quad b_{i,j} = b_{\sigma_+^{j-1}(i),1}.$$

Now label the columns of $M(\alpha_0)$ by v_1, \dots, v_{2n} . Let N be the matrix with columns v'_1, \dots, v'_{2n} where $v'_i = v_i$ for $1 \leq i \leq n$ and $v'_{n+i} = v_{n+i} - v_{n+1-i}$ for $1 \leq i \leq n$. For example, if $n = 4$ and $k = 3$ (so that $m = 1$), we would have:

$$M(\alpha_0) = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \end{pmatrix}; \quad N = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 1 & 1 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 1 & -1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

If $N = \begin{pmatrix} A & C \\ B & D \end{pmatrix}$ for matrices $C = (c_{i,j})$ and $D = (d_{i,j})$ then $c_{i,j} = b_{i,j} - a_{i,n+1-j}$ and $d_{i,j} = a_{i,j} - b_{i,n+1-j}$. Now,

$$a_{i,n} = a_{\sigma_{-}^{n-1}(i),1} = \begin{cases} 1 & \text{if } i \in \{n-m, n-m+1, \dots, n\}, \\ 0 & \text{otherwise,} \end{cases}$$

and so

$$c_{i,1} = \begin{cases} -1 & \text{if } i = n-m, \\ 0 & \text{otherwise,} \end{cases}$$

Similarly,

$$d_{i,1} = \begin{cases} 1 & \text{if } i = n-m+1, \\ 0 & \text{otherwise.} \end{cases}$$

We also have

$$c_{i,j} = c_{\sigma_{+}^{j-1}(i),1} \quad \text{and} \quad d_{i,j} = d_{\sigma_{-}^{j-1}(i),1}.$$

There is precisely one -1 appearing in the i -th row of C ; fix i, j such that $c_{i,j} = -1$. Then $c_{\sigma_{+}^{j-1}(i),1} = -1 \Rightarrow \sigma_{+}^{j-1}(i) = n-m \Rightarrow j = \sigma_{-}^{i-1}(n-m)$. The row of D containing $+1$ in the j -th position is the k -th, where

$$\begin{aligned} d_{k,\sigma_{-}^{i-1}(n-m)} = 1 &\Rightarrow d_{\sigma_{-}^{\sigma_{-}^{i-1}(n-m)-1}(k),1} = 1 \\ &\Rightarrow \sigma_{-}^{\sigma_{-}^{i-1}(n-m)-1}(k) = n-m+1 \\ &\Rightarrow k - \sigma_{-}^{i-1}(n-m) + 1 = n-m+1 \pmod n \\ &\Rightarrow k - n + m + i = n-m+1 \pmod n \\ &\Rightarrow k = n - 2m - i + 1 \pmod n \\ &\Rightarrow k = \sigma_{-}^{i-1}(n-2m). \end{aligned}$$

Let the rows of N be labelled by w_1, \dots, w_{2n} . Put $w'_i = w_i$ for $n+1 \leq i \leq 2n$ and $w'_i = w_i + w_{n+\sigma_{-}^{i-1}(n-2m)}$ for $1 \leq i \leq n$. If we let P be the matrix with rows w'_1, \dots, w'_{2n} then by the preceding argument P is of the form $P = \begin{pmatrix} E & 0 \\ B & D \end{pmatrix}$. Here D is a permutation matrix, and so we have $\det M(\alpha_0) = \pm \det E$. In the case $n = 4, k = 3$ we have:

$$E = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \end{pmatrix}$$

If $E = (e_{i,j})$, then

$$e_{i,j} = a_{i,j} + b_{\sigma_{-}^{i-1}(n-2m),j}.$$

Consider

$$\begin{aligned} e_{i,j} - e_{\sigma_{-}^{j-1}(i),1} &= a_{i,j} + b_{\sigma_{-}^{i-1}(n-2m),j} - a_{\sigma_{-}^{j-1}(i),1} - b_{\sigma_{-}^{\sigma_{-}^{j-1}(i)-1}(n-2m),1} \\ &= b_{\sigma_{+}^{j-1}(\sigma_{-}^{i-1}(n-2m)),1} - b_{\sigma_{-}^{\sigma_{-}^{j-1}(i)-1}(n-2m),1}, \end{aligned}$$

where we have cancelled the a terms. Now,

$$\begin{aligned} \sigma_{-}^{\sigma_{-}^{j-1}(i)-1}(n-2m) &= n-2m - \sigma_{-}^{j-1}(i) + 1 = n-2m - (i-j+1) + 1 \pmod n \\ &= n-2m + j - i \pmod n. \end{aligned}$$

However,

$$\begin{aligned} \sigma_{+}^{j-1}(\sigma_{-}^{i-1}(n-2m)) &= \sigma_{-}^{i-1}(n-2m) + j - 1 = n-2m - i + 1 + j - 1 \pmod n \\ &= n-2m + j - i \pmod n, \end{aligned}$$

so the b terms also cancel, and we can conclude that $e_{i,j} = e_{\sigma_{-}^{j-1}(i),1}$.

Consider the first column of E : we know that

$$b_{\sigma_{-}^{i-1}(n-2m),1} = \begin{cases} 1 & \text{if } \sigma_{-}^{i-1}(n-2m) \in \{n-m+1, n-m+2, \dots, n\}, \\ 0 & \text{otherwise.} \end{cases}$$

However,

$$\sigma_{-}^{i-1}(n-2m) \in \{n-m+1, \dots, n\} \iff [n-2m-i+1] \in \{[n-m+1], \dots, [n]\},$$

where $[\]$ represents class modulo n . This is equivalent to

$$[-i] \in \{[2m-1], [2m-2], \dots, [m]\},$$

or $i \in \{n-2m+1, n-2m+2, \dots, n-m\}$. Comparing this with the $a_{i,1}$ s, we see that

$$e_{i,1} = \begin{cases} 1 & \text{if } i \in \{1, n-2m+1, \dots, n\}, \\ 0 & \text{otherwise,} \end{cases}$$

so that E has $2m+1 = k$ 1s in each column. We may cyclically permute the rows of E to form a new matrix $F = (f_{i,j})$ with $f_{i,j} = f_{\sigma_{-}^{j-1}(i),1}$ and

$$f_{i,1} = \begin{cases} 1 & \text{if } 1 \leq i \leq k, \\ 0 & \text{otherwise.} \end{cases}$$

The matrix F is the circulant matrix associated to the row vector $(v_0, v_1, \dots, v_{n-1})$ with $v_i = 1$ for $0 \leq i \leq k-1$ and $v_i = 0$ for $k-1 \leq i \leq n-1$. The determinant of

F is given by the well-known formula (see for example [2]):

$$\det F = \prod_{i=0}^{n-1} \sum_{j=0}^{n-1} \zeta^{ij} v_j,$$

where ζ is a primitive n -th root of unity. Write $\lambda_i = \sum_{j=0}^{n-1} \zeta^{ij} v_j$; clearly $\lambda_0 = k$. However, for each $i \geq 1$, we have

$$\lambda_i = \sum_{j=0}^{k-1} (\zeta^i)^j = \frac{\zeta^{ik} - 1}{\zeta^i - 1},$$

and hence

$$\det F = k \prod_{i=1}^{n-1} \frac{\zeta^{ik} - 1}{\zeta^i - 1}.$$

We note that since k is coprime to n , the sets $\{\zeta^{ik} \mid i \in \{1, 2, \dots, n-1\}\}$ and $\{\zeta^i \mid i \in \{1, 2, \dots, n-1\}\}$ coincide, and hence $\det \alpha_0 = \pm \det F = \pm k$. \square

Proposition 2.4 $\det \alpha_1 = \det \alpha_2 \neq 0$.

Proof The following commutes:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & J & \longrightarrow & \Lambda^3 & \xrightarrow{\partial_2} & \Lambda^2 & \xrightarrow{\partial_1} & \Lambda & \xrightarrow{\varepsilon} & \mathbf{Z} & \longrightarrow & 0 \\ & & \downarrow \theta' & & \downarrow \alpha'_2 & & \downarrow \alpha_1 & & \downarrow \alpha_0 & & \downarrow k & & \\ 0 & \longrightarrow & J & \longrightarrow & \Lambda^3 & \xrightarrow{\partial_2} & \Lambda^2 & \xrightarrow{\partial_1} & \Lambda & \xrightarrow{\varepsilon} & \mathbf{Z} & \longrightarrow & 0 \end{array}$$

where

$$\alpha'_2 = \begin{pmatrix} m+1-my & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

and θ' is the restriction of α'_2 to J . We proceed to calculate $\det \alpha'_2 = \det(m+1-my)$. If we represent $(m+1-my)$ with respect to the basis $\{1, x, \dots, x^{n-1}, y, \dots, x^{n-1}y\}$, then we form the matrix:

$$M = \begin{pmatrix} A & B \\ B & A \end{pmatrix}$$

Here A is diagonal with each diagonal entry equal to $m+1$, and B is equal to $-m$ times the permutation matrix associated to $\begin{pmatrix} 1 & 2 & 3 & \dots & n \\ 1 & n & n-1 & \dots & 2 \end{pmatrix}$. Label the rows of M by v_1, \dots, v_{2n} and let N be the matrix with rows v'_1, \dots, v'_{2n} , where $v'_1 = v_1 + v_{n+1}$, $v'_i = v_i + v_{2n-i+2}$ for $2 \leq i \leq n$, and $v'_i = v_i$ for $n+1 \leq i \leq 2n$. Now label the columns of M by w_1, \dots, w_{2n} and let L be the matrix with columns w'_1, \dots, w'_{2n}

where $w'_i = w_i$ for $1 \leq i \leq n$, $w'_{n+1} = w_{n+1} - w_1$ and $w'_{n+i} = w_{n+1} - w_{n-i+2}$ for $2 \leq i \leq n$. For example, if $n = 4$ and $k = 3$ (so that $m = 1$) we have:

$$N = \begin{pmatrix} 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 2 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 2 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 2 \end{pmatrix}; \quad L = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 3 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 3 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 3 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 3 \end{pmatrix}$$

It is easy to see that L is lower triangular with n diagonal entries equal to 1 and n diagonal entries equal to $2m + 1 = k$. Then $\det \alpha'_2 = \det(m + 1 - my) = \det L = k^n$. Using $k \det \theta' \det \alpha_1 = \det \alpha_0 \det \alpha'_2 = \pm k^{n+1}$ we see that $\det \alpha_1 = \det \alpha_2 \neq 0$. \square

Therefore by Propositions 2.2, 2.3 and 2.4:

Proposition 2.5 *If $3 \leq k \leq n - 1$ is coprime to $2n$ then $\det \theta = \pm 1$ and so θ is an isomorphism. Thus $[k]$ is in the image of the Swan map.*

Clearly $[-1]$ is in the image of the Swan map and so:

Corollary 2.6 *The Swan map $\text{Aut } J \rightarrow (\mathbf{Z}/2n)^*$ is surjective for each D_{2n} .*

Mannan [9] has previously shown that the Swan map is surjective for D_{2n} .

3 The $D(2)$ -property for $\mathbf{Z}[D_{4n}]$

We now restrict to the case D_{4n} . An application of Schanuel’s lemma shows that the module J appearing in (2) is determined up to stable equivalence; that is, if $0 \rightarrow J \rightarrow F_2 \rightarrow F_1 \rightarrow F_0 \rightarrow \mathbf{Z} \rightarrow 0$ and $0 \rightarrow J' \rightarrow F'_2 \rightarrow F'_1 \rightarrow F'_0 \rightarrow \mathbf{Z} \rightarrow 0$ are two algebraic 2-complexes, we have $J \oplus \Lambda^n \cong J' \oplus \Lambda^m$ for some n, m . Write $\Omega_3(\mathbf{Z})$ for the class of modules J' appearing in an algebraic 2-complex over D_{4n} . Now take $J = \ker \partial_2$ in (3); the following proposition is due to Mannan [9]:

Proposition 3.1 *J has minimal \mathbf{Z} -rank in $\Omega_3(\mathbf{Z})$.*

Let Γ be an order over a Dedekind domain R . We say that *torsion-free cancellation* holds for Γ if $X \oplus M \cong X \oplus N \implies M \cong N$ for lattices X, M and N over Γ (so that X, M and N are finitely generated as Γ -modules and torsion-free over R). There are very few finite groups G for which $\Gamma = \mathbf{Z}[G]$ has torsion-free cancellation; if G is nonabelian then the only possible candidates are A_4, A_5, S_4 and D_{2n} for certain values of n . Clearly we have:

Proposition 3.2 *Suppose that $\mathbf{Z}[D_{4n}]$ has torsion-free cancellation. Then every $J' \in \Omega_3(\mathbf{Z})$ is of the form $J' \cong J \oplus \Lambda^m$ for some $m \geq 0$.*

For a finite group G , the integral group ring $\mathbf{Z}[G]$ is a \mathbf{Z} -order in the semisimple algebra $\mathbf{Q}[G]$; we may choose a maximal \mathbf{Z} -order Γ in $\mathbf{Q}[G]$ containing $\mathbf{Z}[G]$, and define $D(\mathbf{Z}[G]) = \ker(\tilde{K}_0(\mathbf{Z}[G]) \rightarrow \tilde{K}_0(\Gamma))$. A necessary condition for $\mathbf{Z}[G]$ to possess torsion-free cancellation is $D(\mathbf{Z}[G]) = 0$. The following is due to Swan [11]:

Theorem 3.3 *Let p be a prime. Then D_{4p} satisfies torsion-free cancellation if and only if $D(\mathbf{Z}[D_{4p}]) = 0$.*

Endo and Miyata [3] calculate the order of $D(\mathbf{Z}[D_{2n}])$ for various values of n . In particular they show $D(\mathbf{Z}[D_{4p}]) = 0$ for prime p when $3 \leq p \leq 31, p = 47, 179$ or 19379 . However, there do exist values of n for which $D(\mathbf{Z}[D_{4n}]) \neq 0$, for example $n = 37$. Moreover, results of Swan show that $D(\mathbf{Z}[D_{4n}]) = 0$ is not a sufficient condition for torsion-free cancellation to hold. For example, $D(\mathbf{Z}[D_{2n}]) = 0$ for all n , yet torsion-free cancellation fails when $n \geq 7$ (see [11, Theorem 8.1]). Of course, although values of n exist for which $\mathbf{Z}[D_{4n}]$ does not have torsion-free cancellation, it may still be the case that cancellation of finitely generated free modules holds within $\Omega_3(\mathbf{Z})$ for such n .

If torsion-free cancellation holds for D_{4n} then, by Theorem 1.3, Corollary 2.6 and Proposition 3.2, up to congruence, the only algebraic 2-complexes over D_{4n} are of the form

$$\mathcal{E}_m = (0 \rightarrow J \oplus \Lambda^m \rightarrow \Lambda^3 \oplus \Lambda^m \xrightarrow{\partial_2 \pi_1} \Lambda^2 \xrightarrow{\partial_1} \Lambda \rightarrow \mathbf{Z} \rightarrow 0),$$

where $\pi_1: \Lambda^3 \oplus \Lambda^m \rightarrow \Lambda^3$ denotes projection onto the first factor. If a pair of algebraic 2-complexes are congruent then they are homotopy equivalent (see Johnson [7, page 182]), and so the \mathcal{E}_m represent all homotopy classes of algebraic 2-complexes over D_{4n} . However, \mathcal{E}_m is geometrically realized by the Cayley complex arising from the presentation

$$\mathcal{G}_m = \langle x, y \mid x^{2n}, y^2, y^{-1}xyx, 1, \dots, 1 \rangle,$$

where there are m trivial relators added to the standard presentation for D_{4n} . Therefore every homotopy class of algebraic 2-complex over D_{4n} is geometrically realized and hence by [Theorem 1.2](#) we have proved [Theorem 1.1](#). By [Theorems 1.1](#) and [3.3](#) we have:

Corollary 3.4 *Let p be a prime and suppose that $D(\mathbb{Z}[D_{4p}]) = 0$. Then the $D(2)$ -property holds for D_{4p} .*

References

- [1] **W J Browning**, *Homotopy types of certain finite CW-complexes with finite fundamental group*, PhD thesis, Cornell University (1978) [MR2628267](#) Available at <http://tinyurl.com/d5gtflx>
- [2] **P J Davis**, *Circulant matrices*, Wiley-Interscience, New York (1979) [MR543191](#)
- [3] **S Endô, T Miyata**, *On the class groups of dihedral groups*, J. Algebra 63 (1980) 548–573 [MR570730](#)
- [4] **K W Gruenberg**, *Homotopy classes of truncated projective resolutions*, Comment. Math. Helv. 68 (1993) 579–598 [MR1241473](#)
- [5] **M Gutierrez, M P Latiolais**, *Partial homotopy type of finite two-complexes*, Math. Z. 207 (1991) 359–378 [MR1115169](#)
- [6] **I Hambleton, M Kreck**, *Cancellation of lattices and finite two-complexes*, J. Reine Angew. Math. 442 (1993) 91–109 [MR1234837](#)
- [7] **F E A Johnson**, *Stable modules and the $D(2)$ -problem*, London Math. Soc. Lect. Note Series 301, Cambridge Univ. Press (2003) [MR2012779](#)
- [8] **M P Latiolais**, *When homology equivalence implies homotopy equivalence for 2-complexes*, J. Pure Appl. Algebra 76 (1991) 155–165 [MR1145864](#)
- [9] **W H Mannan**, *The $D(2)$ property for D_8* , Algebr. Geom. Topol. 7 (2007) 517–528 [MR2308955](#)
- [10] **W H Mannan**, *Realizing algebraic 2-complexes by cell complexes*, Math. Proc. Cambridge Philos. Soc. 146 (2009) 671–673 [MR2496351](#)
- [11] **R G Swan**, *Torsion free cancellation over orders*, Illinois J. Math. 32 (1988) 329–360 [MR947032](#)
- [12] **C T C Wall**, *Finiteness conditions for CW-complexes*, Ann. of Math. 81 (1965) 56–69 [MR0171284](#)

Department of Mathematics, University College London
Gower Street, London, WC1E 6BT, United Kingdom
s.o'shea@ucl.ac.uk

Received: 29 March 2012 Revised: 17 August 2012