

The eta-inverted sphere over the rationals

GLEN MATTHEW WILSON

We calculate the motivic stable homotopy groups of the two-complete sphere spectrum after inverting multiplication by the Hopf map η over fields of cohomological dimension at most 2 with characteristic different from 2 (this includes the p -adic fields \mathbb{Q}_p and the finite fields \mathbb{F}_q of odd characteristic) and the field of rational numbers; the ring structure is also determined.

14F42; 18G15, 55Q45, 55T15

1 Introduction

Guillou and Isaksen [7] laid the foundation for calculating $\pi_{**}(\mathbb{1}_2^\wedge)[\eta^{-1}]$, the motivic stable homotopy groups of the two-complete sphere spectrum after inverting multiplication by η , over the complex numbers using the h_1 -inverted motivic Adams spectral sequence. They conjectured a pattern of differentials in the h_1 -inverted motivic Adams spectral sequence and identified the E_∞ page of the spectral sequence assuming the conjecture. Shortly after Guillou and Isaksen's paper appeared, Andrews and Miller [2] proved Guillou and Isaksen's conjecture. All together, these results show $\pi_{**}(\mathbb{1}_2^\wedge)[\eta^{-1}] \cong \mathbb{F}_2[\eta^{\pm 1}, \mu, \varepsilon]/(\varepsilon^2)$ [7, Conjecture 1.3]. Guillou and Isaksen [8] then analyzed the h_1 -inverted motivic Adams spectral sequence over the real numbers and gave a complete calculation of the ring $\pi_{**}(\mathbb{1}_2^\wedge)[\eta^{-1}]$ over the base field \mathbb{R} .

The subject of this paper is the calculation of $\pi_{**}(\mathbb{1}_2^\wedge)[\eta^{-1}]$ over the field of rational numbers \mathbb{Q} and fields F with $\text{cd}_2(F) \leq 2$ and characteristic different from 2, such as the p -adic fields \mathbb{Q}_p and finite fields \mathbb{F}_q of odd characteristic. We write $\pi_{s,w}(\mathbb{1}_2^\wedge)(F)$ for the stable homotopy group $\mathcal{SH}(F)(\Sigma^{s,w}\mathbb{1}, \mathbb{1}_2^\wedge)$ and frequently abbreviate this to $\pi_{s,w}(\mathbb{1}_2^\wedge)$ if the base field F is clear from context.

We write $\mathfrak{M}(F)$ for the motivic Adams spectral sequence at the prime 2 over the field F at the motivic sphere spectrum. This spectral sequence has E_2 page given by $\mathfrak{M}(F)_2^{f,s,w} = \text{Ext}_{A_{**}(F)}^{f,s+f,w}(H_{**}(F), H_{**}(F))$ and conditionally converges:

$$\mathfrak{M}(F)^{f,s,w} \implies \pi_{s,w}(\mathbb{1}_2^\wedge)(F),$$

where $\mathbb{1}_H^\wedge$ is the H -nilpotent completion of the sphere spectrum defined by Bousfield [4, Section 5]. Hu, Kriz and Ormsby [9, Theorem 1] proved that $\mathbb{1}_H^\wedge$ is weakly equivalent to $\mathbb{1}_2^\wedge$ over a field F of characteristic 0 with $\text{cd}_2(F[\sqrt{-1}]) < \infty$. Wilson and Østvær [20, Proposition 5.10] note that the same argument works over fields of positive characteristic under the assumption that $\text{cd}_2(F[\sqrt{-1}]) < \infty$.

Given the conditionally convergent spectral sequence $\mathfrak{M}(F) \Rightarrow \pi_{**}(\mathbb{1}_2^\wedge)(F)$ and the fact that $\eta \in \pi_{1,1}(\mathbb{1}_2^\wedge)(F)$ is detected by $h_1 \in \mathfrak{M}(F)^{1,1,1}$, one can try to calculate $\pi_{**}(\mathbb{1}_2^\wedge)(F)[\eta^{-1}]$ using the h_1 -inverted spectral sequence, defined as the following colimit of spectral sequences:

$$\mathfrak{M}(F)[h_1^{-1}] = \text{colim}(\mathfrak{M}(F) \xrightarrow{h_1} \mathfrak{M}(F) \xrightarrow{h_1} \mathfrak{M}(F) \cdots).$$

It is not obvious that $\mathfrak{M}(F)[h_1^{-1}]$ converges to $\pi_{**}(\mathbb{1}_2^\wedge)(F)[\eta^{-1}]$. Guillou and Isaksen show that it does converge for the complex numbers \mathbb{C} in [7, Section 6] and the real numbers \mathbb{R} in [8, Section 5]. We address convergence for more general fields in Section 2.

The Milnor–Witt t -stem of $\mathbb{1}_2^\wedge$ over F is the group $\hat{\Pi}_t(F) = \bigoplus_{k \in \mathbb{Z}} \pi_{k+t,k}(\mathbb{1}_2^\wedge)(F)$. Note that $\hat{\Pi}_0(F)$ is a ring and $\hat{\Pi}_t(F)$ is a $\hat{\Pi}_0(F)$ -module. Our main results will be stated in terms of Milnor–Witt stems and the Witt group of quadratic forms $W(F)$. In many cases, the two-complete η -inverted Milnor–Witt 0-stem can be described in terms of the Witt ring of quadratic forms of the field.

Proposition 1 *If F is a field for which the Witt group of quadratic forms $W(F)$ is finitely generated or $W(F)$ has bounded 2-torsion exponent (if $F = \mathbb{Q}$, for example), then there is an isomorphism $\hat{\Pi}_0(F)[\eta^{-1}] \cong W(F)_2^\wedge[\eta^{\pm 1}]$.*

Proof Morel [14] has shown there is an isomorphism $\pi_{n,n}(\mathbb{1}) \cong W(F)$ for $n \geq 1$. For $n \geq 1$ the homotopy group $\pi_{n,n}(\mathbb{1}_2^\wedge)$ fits into the exact sequence

$$0 \rightarrow \text{Ext}(\mathbb{Z}/2^\infty, W(F)) \rightarrow \pi_{n,n}(\mathbb{1}_2^\wedge) \rightarrow \text{Hom}(\mathbb{Z}/2^\infty, \pi_{n-1,n}(\mathbb{1})) \rightarrow 0$$

by [9, Equation (2)]. But as $\pi_{n-1,n}\mathbb{1} = 0$ by Morel’s connectivity theorem [14, Theorem 1.18], there is an isomorphism $\text{Ext}(\mathbb{Z}/2^\infty, W(F)) \rightarrow \pi_{n,n}(\mathbb{1}_2^\wedge)$.

If $W(F)$ is finitely generated, there is an isomorphism $\text{Ext}(\mathbb{Z}/2^\infty, W(F)) \cong W(F)_2^\wedge$ by a result of Bousfield and Kan [5, Chapter VI, Section 2.1], hence $\pi_{n,n}(\mathbb{1}_2^\wedge) \cong W(F)_2^\wedge$. If $W(F)$ has bounded 2-torsion exponent, then ${}_nW(F) = {}_mW(F)$ for all n and m sufficiently large. The Mittag-Leffler condition is satisfied for the tower $\{{}_nW(F)\}$,

hence $\varprojlim^1 2^n W(F) \cong \varprojlim^1 \text{Hom}(\mathbb{Z}/2^n, W(F))$ vanishes. By the short exact sequence of Weibel [19, Application 3.5.10],

$$0 \rightarrow \varprojlim^1 \text{Hom}(\mathbb{Z}/2^n, W(F)) \rightarrow \text{Ext}(\mathbb{Z}/2^\infty, W(F)) \rightarrow W(F)_2^\wedge \rightarrow 0,$$

there is an isomorphism $\text{Ext}(\mathbb{Z}/2^\infty, W(F)) \cong W(F)_2^\wedge$, and so $\pi_{n,n}(\mathbb{1}_2^\wedge) \cong W(F)_2^\wedge$.

Finally, there is an isomorphism $\widehat{\Pi}_0(F)[\eta^{-1}] \cong W(F)_2^\wedge[\eta^{\pm 1}]$ since for any class $\alpha \in \widehat{\Pi}_0(F)$ and n sufficiently large, the class $\eta^n \alpha$ is an element of $\pi_{k,k} \mathbb{1} \cong W(F)$ with $k \geq 1$. □

For finite fields \mathbb{F}_q , the Milnor–Witt 0–stem is now determined by the calculation of the Witt group of finite fields, a standard reference being Scharlau [17, Chapter 2, Theorem 3.3]:

$$\widehat{\Pi}_0(\mathbb{F}_q)[\eta^{-1}] \cong W(\mathbb{F}_q)_2^\wedge[\eta^{\pm 1}] = \begin{cases} \mathbb{Z}/2[\eta^{\pm 1}, u]/u^2 & \text{if } q \equiv 1 \pmod{4}, \\ \mathbb{Z}/4[\eta^{\pm 1}] & \text{if } q \equiv 3 \pmod{4}. \end{cases}$$

We find in Theorem 9 that for a field F with $\text{cd}_2(F) \leq 2$ and characteristic different from 2, the two-complete η -inverted Milnor–Witt stems take the following form:

$$\widehat{\Pi}_t(F)[\eta^{-1}] \cong \begin{cases} W(F)_2^\wedge[\eta^{\pm 1}] & \text{if } t \geq 0 \text{ and either } t \equiv 3 \pmod{4} \text{ or } t \equiv 0 \pmod{4}, \\ 0 & \text{otherwise.} \end{cases}$$

This gives a complete calculation of $\widehat{\Pi}_*(\mathbb{F}_q)[\eta^{-1}]$ for the finite fields \mathbb{F}_q of odd characteristic.

Theorem 19 calculates the ring $\widehat{\Pi}_*(\mathbb{Q})[\eta^{-1}]$. In particular, the Milnor–Witt stems are

$$\widehat{\Pi}_t(\mathbb{Q})[\eta^{-1}] \cong \begin{cases} W(\mathbb{Q})_2^\wedge[\eta^{\pm 1}] & \text{if } t = 0, \\ W(\mathbb{Q})_2^\wedge[\eta^{\pm 1}]/2^{n+1} & \text{if } t \geq 0, t \equiv 3 \pmod{4}, n = \nu_2(t + 1), \\ M & \text{if } t \equiv 0 \pmod{4}, t \geq 4, \\ 0 & \text{otherwise,} \end{cases}$$

where $\nu_2(t + 1)$ is the 2–adic valuation of $t + 1$ and M is the submodule of $W(\mathbb{Q})_2^\wedge[\eta^{\pm 1}]$ defined in Definition 18.

The method of proof for the calculations over \mathbb{Q} of Theorem 19 follows the strategy employed by Ormsby and Østvær [15] to calculate the homotopy groups of $BP\langle n \rangle$ over \mathbb{Q} . First, for each completion \mathbb{Q}_ν of \mathbb{Q} one uses the ρ –Bockstein spectral sequence to calculate $\text{Ext}(\mathbb{Q}_\nu)[h_1^{-1}]$ and then the motivic Adams spectral sequence to calculate $\pi_{**}(\mathbb{1}_2^\wedge)(\mathbb{Q}_\nu)[\eta^{-1}]$. We next follow the motivic Hasse principle to identify the differentials in the ρ –Bockstein spectral sequence and the motivic Adams spectral

sequence over \mathbb{Q} by comparing these spectral sequences with the associated spectral sequences over the completions.

The result of Ormsby, Röndigs and Østvær [16, Proof of Theorem 1.5] shows that the vanishing $\widehat{\Pi}_t(\mathbb{Q})[\eta^{-1}] = 0$ when $t \equiv 1, 2 \pmod{4}$ occurs systematically for all formally real fields F with $\text{cd}_2(F[i]) < \infty$. Their calculation is used in this paper to show that $\widehat{\Pi}_1(\mathbb{Q})[\eta^{-1}]$ vanishes, as it is unclear whether or not the motivic Adams spectral sequence over \mathbb{Q} converges strongly in Milnor–Witt stem 1.

Ananyevskiy, Levine and Panin [1] investigate the η -inverted sphere spectrum $\mathbb{1}[\eta^{-1}]$ over fields F of characteristic different from 2. They find that the stable homotopy sheaf $\bigoplus_{n \in \mathbb{Z}} \pi_{n,n} \mathbb{1}[\eta^{-1}]$ is isomorphic to the sheaf $\underline{W}[\eta^{\pm 1}]$, where \underline{W} is the Nisnevich sheaf associated to the presheaf of Witt groups (the Witt group $W(X)$ of an algebraic variety X is defined by Knebusch [12, Chapter I, Section 5]). The consequence of this for calculating stable homotopy groups is that

$$\bigoplus_{n \in \mathbb{Z}} \pi_{n,n}(\mathbb{1}[\eta^{-1}](F)) \cong W(F)[\eta^{\pm 1}]$$

for all fields F of characteristic different from 2. In addition to this absolute statement about the η -inverted Milnor–Witt 0–stem, they identify the rationalization of $\mathbb{1}[\eta^{-1}]$ with an object in the heart of the homotopy t -structure on $\mathcal{SH}(F)$ [1, Theorem 3.4] and find that the sheaf $\pi_{s,w}(\mathbb{1}[\eta^{-1}]_{\mathbb{Q}})$ takes the following form:

$$\pi_{s,w}(\mathbb{1}[\eta^{-1}]_{\mathbb{Q}}) = \begin{cases} W_{\mathbb{Q}} & \text{if } s = w, \\ 0 & \text{otherwise.} \end{cases}$$

The calculations in this paper are about the η -inverted 2–complete sphere spectrum $\mathbb{1}_2^{\wedge}[\eta^{-1}]$ in contrast to Ananyevskiy, Levine and Panin’s results about $\mathbb{1}[\eta^{-1}]$ and $\mathbb{1}[\eta^{-1}]_{\mathbb{Q}}$.

We will follow the grading conventions for $\text{Ext}(F) = \text{Ext}_{\mathcal{A}_{**}(F)}(H_{**}(F), H_{**}(F))$ employed by Guillou and Isaksen [7, Section 2.1]. In particular, for a class $x \in \text{Ext}(F)$ in Adams filtration f , stem s , and weight w , the Milnor–Witt stem of x is $t = s - w$ and the Chow weight of x is $c = s + f - 2w$. We will write degrees as $\text{deg}(x) = (f, t, c)$ unless otherwise specified. We will frequently use the isomorphism $\text{Ext}(\mathbb{C})[h_1^{-1}] \cong \mathbb{F}_2[h_1^{\pm 1}, v_1^4, v_2, v_3, \dots]$ with $\text{deg}(v_1^4) = (4, 4, 4)$ and $\text{deg}(v_n) = (1, 2^n - 1, 1)$ established in [7, Theorem 3.4], and we adopt the convention of writing P for v_1^4 .

Acknowledgements I would like to thank Dan Isaksen for teaching me how to use Massey product shuffles, Elden Elmanto for the motivation to write up the convergence

result of Section 2, Jonas Kylling and Håkon Kolderup for their comments on an earlier version of this work, Oliver Röndigs for helpful suggestions, and Paul Arne Østvær for his support and encouragement that made this paper possible. This paper was partially completed while participating in the research program “Algebraic-geometric and homotopical methods” at the Institut Mittag-Leffler. I gratefully acknowledge support from the RCN project “Motivic Hopf Equations”, no. 250399.

2 Convergence of the h_1 -inverted motivic Adams spectral sequence

We refer the reader to Boardman’s notes on spectral sequences [3] for the terminology concerning their convergence. The notion of a map of filtered groups compatible with a map of spectral sequences is defined by Weibel in [19, page 126].

Consider a collection of cohomologically graded spectral sequences $({}^iE_r, {}^i d_r)$ for $i \in \mathbb{N}$, where ${}^iE_r^s = 0$ for all $s < 0$ and all r , and each spectral sequence iE converges strongly to an abelian group iG filtered by iF_s . Here r indicates the page of the spectral sequence and s indicates the internal degree. We will omit these subscripts and superscripts where it is convenient. These assumptions on our spectral sequences precisely mean the following conditions hold:

- (1) The filtration is exhaustive with ${}^iF_0 = {}^iG$:

$${}^iG = {}^iF_0 \supseteq {}^iF_1 \supseteq {}^iF_2 \supseteq \dots .$$

- (2) For all $s \in \mathbb{Z}$ there are isomorphisms ${}^iE_\infty^s \cong {}^iF_s / {}^iF_{s+1}$.

- (3) The filtration is Hausdorff: $\bigcap_j {}^iF_j = 0$.

- (4) The filtration is complete: $\varprojlim_j {}^iG / {}^iF_j \cong {}^iG$.

We assume the spectral sequences are \mathbb{Z}^k -graded in addition to the internal grading s . In practice, k is either 1 or 2.

Let ${}^i f: {}^iE \rightarrow {}^{i+1}E$ be a directed system of maps of spectral sequences compatible with ${}^i g: {}^iG \rightarrow {}^{i+1}G$ of degree $a \in \mathbb{Z}^k$. The colimit of the directed system ${}^i f: {}^iE \rightarrow {}^{i+1}E$ is again a spectral sequence $\tilde{E} = \text{colim } {}^iE$ that converges weakly to $\tilde{G} = \text{colim } {}^iG$ filtered by $\tilde{F}_j = \text{colim } {}^iF_j$. Under what circumstances can we guarantee \tilde{E} converges strongly to \tilde{G} ?

The filtration \tilde{F}_j of \tilde{G} may fail to be Hausdorff or complete. To see how the Hausdorff condition may fail, consider ${}^iE = \mathbb{Z}$, ${}^iG = \mathbb{Z}$, and ${}^iF_j = 0$ for $j \geq i$ and ${}^iF_j = {}^iG$ for $j < i$. If we define maps ${}^if = 0$ and ${}^ig(1) = 1$, then $\tilde{E} = 0$, $\tilde{G} = \mathbb{Z}$, and $\tilde{F}_j = \mathbb{Z}$ for all $j \geq 0$. The issue is that the class $1 \in {}^iG$ is in filtration i , which shows $1 \in \tilde{G}$ is in filtration i for all natural numbers i .

Completeness can fail if each iG has a finite filtration but the colimit \tilde{G} has an infinite filtration. Consider ${}^iG = \mathbb{Z}$, ${}^iF_j = 2^j\mathbb{Z}$ for $j \leq i$ and ${}^iF_j = 0$ for $j > i$ with the map ${}^ig(1) = 1$. Then $\tilde{G} = \mathbb{Z}$, yet $\tilde{E}^s \cong \tilde{F}_s/\tilde{F}_{s+1} \cong \mathbb{Z}/2$ for all $s \geq 0$, and $\varprojlim_s \tilde{G}/\tilde{F}_s$ is isomorphic to the 2-adic integers \mathbb{Z}_2 .

Definition 2 Consider a directed system of \mathbb{Z}^k -graded spectral sequences ${}^if: {}^iE \rightarrow {}^{i+1}E$ of degree a , that is, for all $i \in \mathbb{N}$ and $x \in {}^iE$ the degree of ${}^if(x)$ is $a + \text{deg}(x)$, and if does not change the internal degree s . The directed system if has a horizontal vanishing line of height N in the direction a if for any degree b there exists $K \in \mathbb{N}$ for which the groups ${}^iE^{s,b+ia}$ vanish for all $i > K$ and $s > N$.

The term horizontal vanishing line comes from the special case where for all i we have ${}^iE = E$ and $E = \bigoplus_{s,p} E^{s,p}$ is a \mathbb{Z} -graded spectral sequence in p with internal degree s . If one makes a chart for iE where the vertical axis is the internal degree s and the horizontal axis is p , a horizontal vanishing line of height N in the direction 1 says that $E^{s,p}$ vanishes when (s, p) is above the horizontal line $s = N$ and p is sufficiently large.

Proposition 3 Suppose ${}^if: {}^iE \rightarrow {}^{i+1}E$ is a directed system of \mathbb{Z}^k -graded spectral sequences of degree a with a horizontal vanishing line of height N in the direction a . The colimit spectral sequence \tilde{E} then converges strongly to \tilde{G} with respect to the filtration \tilde{F} .

Proof We first show the filtration \tilde{F} of \tilde{G} is complete. For $b \in \mathbb{Z}^k$, the degree- b component of \tilde{G} is

$$\tilde{G}^b = \text{colim}({}^0G^b \rightarrow {}^1G^{b+a} \rightarrow \dots \rightarrow {}^iG^{b+ia} \rightarrow \dots).$$

The assumption that there is a horizontal vanishing line of height N in the direction a implies for all $i > K$ the filtration of ${}^iG^{b+ia}$ is finite. This is because the filtration of iG is Hausdorff and ${}^iE^{s,b+ia}$ vanishes for $s > N$, so iF_j is trivial for $i > K$

and $j > N$. Since finite limits and directed colimits commute, it follows that

$$\begin{aligned} \varprojlim_j (\tilde{G}^b / \tilde{F}_j^b) &\cong \varprojlim_j (\operatorname{colim}_{i>K} {}^iG / {}^iF_j) \\ &\cong \operatorname{colim}_{i>K} (\varprojlim_j {}^iG / {}^iF_j) \\ &\cong \operatorname{colim}_{i>K} {}^iG^{b+ia} \\ &\cong \tilde{G}^b. \end{aligned}$$

We now show the filtration \tilde{F}_j of \tilde{G} is Hausdorff. Let $x \in \tilde{G}^b$ be a nonzero element. **Lemma 4** shows there is some ${}^i x \in {}^iG^{b+ia}$ which maps to $x \in \tilde{G}^b$ for which ${}^i x$ is detected by ${}^i y \in {}^iE_r^{s,b+ia}$ and ${}^{i+k+1}y$ is nonzero for all $k \in \mathbb{N}$. Since ${}^i f$ is compatible with ${}^i g$, it follows that ${}^{i+k+1}y$ detects ${}^{i+k+1}x = {}^{i+k}g \circ \dots \circ {}^i g({}^i x)$ for all $k \in \mathbb{N}$. Furthermore, ${}^{i+k}x \in {}^{i+k}G$ is nonzero for all $k \in \mathbb{N}$, and so ${}^{i+k}y$ survives to ${}^{i+k}E_\infty^{s,b+(i+k)a}$. Our assumption that the spectral sequences iE converge strongly to iG means that

$${}^{i+k}E_\infty^{s,b+(i+k)a} \cong {}^{i+k}F_s / {}^{i+k}F_{s+1}.$$

Hence every class ${}^{i+k}x$ is in filtration s but not $s+1$, so that $x \in \tilde{F}_s$ but $x \notin \tilde{F}_{s+1}$. It now follows that the filtration \tilde{F} of \tilde{G} is Hausdorff. \square

Lemma 4 Under the conditions of **Proposition 3**, consider a nonzero element $x \in \tilde{G}^b$. There exists some ${}^i x \in {}^iG^{b+ia}$ that maps to $x \in \tilde{G}^b$ for which ${}^i x$ is detected by ${}^i y \in {}^iE_r^{s,b+ia}$ and

$${}^{i+k+1}y = {}^{i+k}f \circ \dots \circ {}^i f({}^i y) \in {}^{i+k+1}E_r^{s,b+(i+k+1)a}$$

is nonzero for all $k \in \mathbb{N}$.

Proof There is some ${}^j x \in {}^jG$ which maps to $x \in \tilde{G}$. The classes

$${}^{j+k}x = {}^{j+k-1}g \circ \dots \circ {}^j g({}^j x)$$

are nonzero for all $k \geq 1$ and are therefore detected by some class ${}^{j+k}y \in {}^{j+k}E^{s_k,*}$. The vanishing line implies $s_k \leq N$ for k sufficiently large, and the compatibility of the maps ${}^j f$ with ${}^j g$ implies s_k is a nondecreasing function of k . Hence s_k is eventually constant, say for all $k + j \geq i$. Then ${}^i x$ has the desired property. \square

These results can be applied to inverting multiplication by h_1 in the motivic Adams spectral sequence at the prime 2 after reindexing the filtration. For a field F , write ${}^iE^f$

for $\mathfrak{M}(F)^{f+i}$. Here f is the internal degree of the spectral sequence (“ f ” for “filtration”). With this convention, the maps $h_1 \cdot : {}^i E \rightarrow {}^{i+1} E$ form a directed system of spectral sequences which is compatible with the maps $\eta : {}^i G \rightarrow {}^{i+1} G$, where ${}^i f_j = F_{j+i}(\pi_{**}(\mathbb{1}_2^\wedge))$. The degree of multiplication by η is $(s, w) = (1, 1)$, where s is the stem and w the weight. A horizontal vanishing line of height N in the direction $(1, 1)$ is equivalent to the following condition: for any (s, w) there exists k such that for all $i > k$ and $f > N$ the group $\mathfrak{M}(F)^{f+i, s+i, w+i}$ vanishes. In the usual manner of drawing a chart for $\mathfrak{M}(F)$, such as those made by Isaksen [10], the horizontal vanishing line for the system ${}^i E^f$ is transformed into a line of slope 1.

Such vanishing conditions occur over \mathbb{R} in positive Milnor–Witt stems as proved by Guillou and Isaksen [8, Lemma 5.1]. Over \mathbb{R} it suffices to take $N = 1$, but one must take larger values for other fields. For fields of cohomological dimension at most 2 and number fields, take $N = 3$ for the positive Milnor–Witt stems.

Corollary 5 *The h_1 -inverted motivic Adams spectral sequence over fields of cohomological dimension at most 2 and the field of rational numbers \mathbb{Q} converges strongly to $\pi_{s,w}(\mathbb{1}_2^\wedge)[\eta^{-1}]$ when $s - w > 1$.*

Proof Consider the directed system ${}^i E^f = \mathfrak{M}(F)^{f+i}$ with maps $h_1 \cdot : {}^i E \rightarrow {}^{i+1} E$ as described above. With $N = 3$, the vanishing conditions required for Proposition 3 are satisfied for fields F of cohomological dimension at most 2. The ρ -Bockstein spectral sequence for such a field has E_1 page $H_{**}(F) \otimes_{\mathbb{F}_2[\tau]} \text{Ext}(\mathbb{C})$ and converges off to $\text{Ext}(F)$. The E_1 page of the ρ -Bockstein spectral sequence has the claimed vanishing line in positive Milnor–Witt stem; hence $\text{Ext}(F)$ does too.

An argument similar to the one given by Guillou and Isaksen in [8, Lemma 5.1] establishes a vanishing line over \mathbb{Q} in positive Milnor–Witt stems with $N = 3$. Their choice of A works just as well over \mathbb{Q} (A corresponds to k when $s = 0$ in the notation above) because the ρ -inverted Hopf algebroid $(H_{**}(\mathbb{Q})[\rho^{-1}], \mathcal{A}_{**}(\mathbb{Q})[\rho^{-1}])$ is isomorphic to the ρ -inverted Hopf algebroid over \mathbb{R} . Their argument with the ρ -Bockstein spectral sequence must only be modified to account for y being of the form $y = \alpha \tilde{y}$ with $\tilde{y} \in \text{Ext}(\mathbb{C})$, and $\alpha \in H_{i,i}(\mathbb{Q})$ with $i \leq 2$ and α not divisible by ρ . The motivic Adams spectral sequence for fields F with $\text{cd}_2(F) \leq 2$ and \mathbb{Q} converges conditionally to $\pi_{**}(\mathbb{1}_2^\wedge)$ by [9, Theorem 1] of Hu, Kriz and Ormsby. The vanishing line described above ensures that it also converges strongly in Milnor–Witt stem at least 2, as in such degrees $d_r = 0$ for r sufficiently large. Hence we get the convergence result of the h_1 -inverted Adams spectral sequences. □

3 Fields of cohomological dimension at most 2

Let F be a field of 2-cohomological dimension at most 2. The mod 2 Milnor K -theory of such a field satisfies $k_n^M(F) = 0$ for $n \geq 3$. We first calculate $\text{Ext}(F)[h_1^{-1}]$ using the ρ -Bockstein spectral sequence and then observe that the structure of $\mathfrak{M}(F)_2 \cong \text{Ext}(F)[h_1^{-1}]$ forces the h_1 -inverted motivic Adams spectral sequence to collapse at the E_2 page. See Figures 1 and 2 for a depiction of the ρ -Bockstein spectral sequence E_1 and E_∞ pages up to Milnor–Witt stem 24.

Proposition 6 For F a field with $\text{cd}_2(F) \leq 2$, the E_2 page of $\mathfrak{M}(F)[h_1^{-1}]$ is

$$\text{Ext}(F)[h_1^{-1}] \cong k_*^M(F) \otimes \text{Ext}(\mathbb{C})[h_1^{-1}].$$

Proof If -1 is a square in F , it follows that $\text{Ext}(F) \cong H_{**}(F) \otimes_{\mathbb{F}_2[\tau]} \text{Ext}(\mathbb{C})$ by an argument similar to [20, Proposition 7.1]. The class τ is killed after inverting h_1 , so the result follows in this case.

If -1 is not a square in F , use the ρ -Bockstein spectral sequence. The E_1 page of the h_1 -inverted ρ -Bockstein spectral sequence is

$$E_1^{\epsilon,*,*,*} \cong \rho^\epsilon k_*^M(F) / \rho^{\epsilon+1} k_*^M(F) \otimes \text{Ext}(\mathbb{C})[h_1^{-1}],$$

and the d_r differential has degree $(r, 1, -1, 0)$ with respect to the grading (ϵ, f, t, c) .

The differentials d_r with $r \geq 1$ vanish on the generators $P = v_1^4$ and v_n for $n \geq 2$ of $\text{Ext}(\mathbb{C})[h_1^{-1}]$ for degree reasons. Any nonzero class $x \in \text{Ext}(\mathbb{C})[h_1^{-1}]$ has $t + c \equiv 0 \pmod{4}$, but the degree of $d_r(x)$ satisfies $t + c \equiv 3 \pmod{4}$. If F has cohomological dimension at most 2, then any nonzero class in the h_1 -inverted ρ -Bockstein spectral sequence satisfies $t + c \not\equiv 3 \pmod{4}$. Hence the h_1 -inverted ρ -Bockstein spectral sequence collapses. There is no possibility for hidden extensions, so the proposition follows. \square

Proposition 7 If \bar{F} is an algebraically closed field of characteristic different from 2, the η -inverted motivic homotopy groups of spheres over \bar{F} are given by

$$\pi_{**}(\mathbb{1}_2^\wedge)(\bar{F})[\eta^{-1}] \cong \pi_{**}(\mathbb{1}_2^\wedge)(\mathbb{C})[\eta^{-1}] \cong \mathbb{F}_2[\eta^{\pm 1}, \mu, \varepsilon]/(\varepsilon^2),$$

where $\mu \in \pi_{9,5}(\mathbb{1}_2^\wedge)$ is the unique homotopy class detected by Ph_1 and $\varepsilon \in \pi_{8,5}(\mathbb{1}_2^\wedge)$ is the unique homotopy class detected by c_0 .




Figure 1: The E_1 page of the h_1 -inverted ρ -Bockstein spectral sequence over a field F with $\text{cd}_2(F) = 2$ up to Milnor–Witt stem 24. Solid vertical lines indicate possible ρ -multiplications that depend on the field. Black dots represent the group $\mathbb{Z}/2[h_1^{\pm 1}]$, blue dots represent $k_1^M(F)[h_1^{\pm 1}]$, and red dots represent $k_2^M(F)[h_1^{\pm 1}]$. Solid lines of slope $\frac{1}{3}$ indicate multiplication by v_2 and arrows in this direction represent a tower of nonzero v_2 multiples. The horizontal axis t is the Milnor–Witt stem and the vertical axis c is the Chow weight, while the Adams filtration is suppressed.

Proof If \bar{F} has characteristic zero, there is an isomorphism $\mathfrak{M}(\bar{F}) \cong \mathfrak{M}(\mathbb{C})$ by the proof of [20, Lemma 6.4]. If \bar{F} has positive characteristic, the change of characteristic argument [20, Corollary 6.1] comparing $\mathfrak{M}(\bar{F})$ to $\mathfrak{M}(\mathbb{C})$ via the motivic Adams spectral sequence over the ring of Witt vectors of \bar{F} shows there is an isomorphism of




Figure 2: The E_∞ page of the h_1 -inverted motivic Adams spectral sequence over a field F with $cd_2(F) \leq 2$ up to Milnor–Witt stem 24. The notational conventions of Figure 1 apply here.

spectral sequences $\mathfrak{M}(\bar{F}) \cong \mathfrak{M}(\mathbb{C})$. This isomorphism propagates to an isomorphism after inverting multiplication by h_1 . The now resolved conjecture of Guillou and Isaksen in [7, Conjecture 1.3] gives the explicit description. □

Proposition 8 *The d_2 differentials for the h_1 -inverted motivic Adams spectral sequence for a field F with characteristic different from 2 and $cd_2(F) \leq 2$ follow from $d_2(v_n) = h_1^2 v_{n-1}^2$ for $n \geq 3$ and $d_2(x) = 0$ for $x \in k_*^M(F)$ by using the Leibniz rule. Furthermore, $\mathfrak{M}(F)[h_1^{-1}]$ collapses at the E_3 page.*

Proof The inclusion of F into its algebraic closure \bar{F} induces a map of spectral sequences

$$\Phi: \mathfrak{M}(F)[h_1^{-1}] \rightarrow \mathfrak{M}(\bar{F})[h_1^{-1}] \cong \mathfrak{M}(\mathbb{C})[h_1^{-1}].$$

Andrews and Miller [2, Theorem 9.15] have proved that in $\mathfrak{M}(\mathbb{C})[h_1^{-1}]$ there are differentials $d_2(v_n) = h_1^2 v_{n-1}^2$ for all $n \geq 3$. It follows that in $\mathfrak{M}(F)[h_1^{-1}]$ we must have $d_2(v_n) = h_1^2 v_{n-1}^2$ up to some element in the kernel of the comparison map Φ . A class $x \in \ker(\Phi)$ satisfies $t + c \equiv 1 \pmod{4}$ or $t + c \equiv 2 \pmod{4}$, whereas $d_2(v_n)$ satisfies $t + c \equiv 0 \pmod{4}$. Hence $d_2(v_n) = h_1^2 v_{n-1}^2$ is true on the nose. That the spectral sequence collapses at the E_3 page follows by degree reasons. \square

Theorem 9 For a field F with $\text{cd}_2(F) \leq 2$ and characteristic different from 2, the two-complete η -inverted Milnor–Witt stems of F are

$$\hat{\Pi}_t(F)[\eta^{-1}] \cong \begin{cases} W(F)_2^\wedge[\eta^{\pm 1}] & \text{for } t \geq 0 \text{ and either } t \equiv 3 \pmod{4} \text{ or } t \equiv 0 \pmod{4}, \\ 0 & \text{otherwise.} \end{cases}$$

$\hat{\Pi}_*(F)[\eta^{-1}]$ is the polynomial ring over $W(F)_2^\wedge[\eta^{\pm 1}]$ on two classes $\{v_2\}$ and $\{P\}$ in Milnor–Witt stems 3 and 4 respectively, subject to the relation $\{v_2\}^2 = 0$.

Proof $\hat{\Pi}_0(F)[\eta^{-1}]$ is shown to be $W(F)_2^\wedge[\eta^{\pm 1}]$ in Proposition 1. The remaining stems and ring structure follow from the calculation of the h_1 -inverted motivic Adams spectral sequence over F whose differentials are determined in Proposition 8. \square

We now identify some classes in $\pi_{**}(\mathbb{1}_2^\wedge)(\mathbb{F}_q)$ for finite fields \mathbb{F}_q using the analysis of the motivic Adams spectral sequence by Wilson and Østvær [20]. Over a finite field \mathbb{F}_q with $q \equiv 1 \pmod{4}$, define $\varepsilon \in \pi_{8,5}(\mathbb{1}_2^\wedge) \cong (\mathbb{Z}/2)^4$ to be a class detected by c_0 . The class ε is uniquely determined modulo $u\eta\varepsilon$. Write μ for a class in $\pi_{9,5}(\mathbb{1}_2^\wedge) \cong (\mathbb{Z}/2)^4$ detected by Ph_1 . The class μ is uniquely determined modulo $u\eta\mu$.

Over a finite field \mathbb{F}_q with $q \equiv 3 \pmod{4}$, there is an isomorphism $\pi_{8,5}(\mathbb{1}_2^\wedge) \cong \mathbb{Z}/4 \oplus \mathbb{Z}/4$. Recall there is a Hopf map $\sigma \in \pi_{7,4}(\mathbb{1}_2^\wedge)$ defined by Dugger and Isaksen in [6]. The class $\eta\sigma$ generates an order-four cyclic subgroup of $\pi_{8,5}(\mathbb{1}_2^\wedge)$; define $\varepsilon \in \pi_{8,5}(\mathbb{1}_2^\wedge)$ by the property that ε generates $\pi_{8,5}(\mathbb{1}_2^\wedge)/(\eta\sigma)$. The class ε is detected by c_0 and well defined up to an odd multiple. Further, there is an isomorphism $\pi_{9,5}(\mathbb{1}_2^\wedge) \cong \mathbb{Z}/4 \oplus (\mathbb{Z}/2)^2$. Define μ to be a class of order four that is detected by Ph_1 ; the class μ is uniquely defined up to an odd multiple.

Corollary 10 For a finite field \mathbb{F}_q with q odd, the η -inverted Milnor–Witt stems are as follows:

$$\widehat{\Pi}_n(\mathbb{F}_q)[\eta^{-1}] \cong \begin{cases} W(\mathbb{F}_q)_2^\wedge[\eta^{\pm 1}] & \text{if } n \geq 0 \text{ and either } n \equiv 3 \pmod 4 \text{ or } n \equiv 0 \pmod 4, \\ 0 & \text{otherwise.} \end{cases}$$

The classes μ and ε generate $\widehat{\Pi}_*(\mathbb{F}_q)[\eta^{-1}]$ as an algebra over $\widehat{\Pi}_0(\mathbb{F}_q)[\eta^{-1}]$, subject to the relation $\varepsilon^2 = 0$. When $q \equiv 3 \pmod 4$ this shows $\widehat{\Pi}_*(\mathbb{F}_q) \cong \mathbb{Z}/4[\eta^{\pm 1}, \mu, \varepsilon]/\varepsilon^2$, and for $q \equiv 1 \pmod 4$ we have $\widehat{\Pi}_*(\mathbb{F}_q) \cong \mathbb{Z}/2[\eta^{\pm 1}, u, \mu, \varepsilon]/(u^2, \varepsilon^2)$.

Proof The mod 2 Milnor K -theory of a finite field with odd characteristic is given by $k_*^M(\mathbb{F}_q) = \mathbb{F}_2[u]/u^2$, where u is the class of a nonsquare element of \mathbb{F}_q^\times . If $q \equiv 3 \pmod 4$ then $u = \rho = [-1]$. As $h_1\rho$ in $\mathfrak{M}(\mathbb{F}_q)[h_1^{-1}]$ detects multiplication by 2 in $\pi_{**}(\mathbb{1}_2^\wedge)$, we arrive at the claimed group structure. The product structure is clear given the products in the h_1 -inverted motivic Adams spectral sequence. \square

Corollary 11 The η -inverted Milnor–Witt stems for a p -adic field \mathbb{Q}_p are as follows:

$$\widehat{\Pi}_0(\mathbb{Q}_p)[\eta^{-1}] \cong W(\mathbb{Q}_p)_2^\wedge[\eta^{\pm 1}] \cong \begin{cases} \mathbb{Z}/2[\eta^{\pm 1}, u, \pi]/(u^2, \pi^2) & \text{if } p \equiv 1 \pmod 4, \\ (\mathbb{Z}/4 \oplus \mathbb{Z}/4)[\eta^{\pm 1}] & \text{if } p \equiv 3 \pmod 4, \\ (\mathbb{Z}/8 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/2)[\eta^{\pm 1}] & \text{if } p = 2, \end{cases}$$

$$\widehat{\Pi}_n(\mathbb{Q}_p)[\eta^{-1}] \cong \begin{cases} W(\mathbb{Q}_p)_2^\wedge[\eta^{\pm 1}] & \text{if } n \geq 0 \text{ and either } n \equiv 3 \pmod 4 \text{ or } n \equiv 0 \pmod 4, \\ 0 & \text{otherwise.} \end{cases}$$

Proof The mod 2 Milnor K -theory of the p -adic fields can be calculated from the result of Milnor [13, Lemma 4.6] in addition to the description of the Witt ring for p -adic fields which is discussed by Serre in [18]. Explicitly, the mod 2 Milnor K -theory of a p -adic field is

$$k_*^M(\mathbb{Q}_p) = \begin{cases} \mathbb{Z}/2[\pi, u]/(\pi^2, u^2) & \text{if } p \equiv 1 \pmod 4, \\ \mathbb{Z}/2[\pi, \rho]/(\rho^2, \rho\pi + \pi^2) & \text{if } p \equiv 3 \pmod 4, \\ \mathbb{Z}/2[\pi, \rho, u]/(\rho^3, u^2, \pi^2, \rho u, \rho\pi, \rho^2 + u\pi) & \text{if } p = 2, \end{cases}$$

where $\pi = [p]$, $\rho = [-1]$, u is the class of a lift of a nonsquare in \mathbb{F}_p^\times when $p \equiv 1 \pmod 4$, and $u = [5]$ when $p = 2$. \square

4 The field of rational numbers

We approach the calculation of $\widehat{\Pi}_*(\mathbb{Q})[\eta^{-1}]$ with the strategy suggested by the motivic Hasse principle, following the method of Ormsby and Østvær in [15]. That is, we

product	conditions
$[\ell][2] = 0$	$\ell = 2$ or $\ell = -1$
$[-1][\ell] = a_\ell$	$\ell = -1$ or ℓ prime and $\ell \equiv 3 \pmod{4}$
$[-1][\ell] = 0$	ℓ prime and $\ell \equiv 1 \pmod{4}$
$[\ell][q] = (q \mid \ell)a_\ell + (\ell \mid q)a_q$	ℓ and q odd primes
$[2][q] = (\frac{1}{8}(q^2 - 1) \pmod{2})a_q$	q odd prime

Table 1: Products in $k_*^M(\mathbb{Q})$

analyze the h_1 -inverted motivic Adams spectral sequence for \mathbb{Q} using our knowledge of the h_1 -inverted motivic Adams spectral sequence over the completions of \mathbb{Q} .

We fix our notation for $k_*^M(\mathbb{Q})$. The mod 2 Milnor K -theory of \mathbb{Q} is generated by the classes $[-1]$ and $[p]$ for p a prime. Milnor shows in [13, Lemma A.1] that there is a short exact sequence

$$0 \rightarrow k_2^M(\mathbb{Q}) \rightarrow \bigoplus k_2^M(\mathbb{Q}_v) \rightarrow \mathbb{Z}/2 \rightarrow 0,$$

where the summation is over all completions \mathbb{Q}_v of \mathbb{Q} . For every completion \mathbb{Q}_v of \mathbb{Q} there is an isomorphism $k_2^M(\mathbb{Q}_v) \cong \mathbb{Z}/2$; write e_v for the image of 1 under the canonical map $k_2^M(\mathbb{Q}_v) \rightarrow \bigoplus k_2^M(\mathbb{Q}_v)$. For ℓ an odd prime or -1 , write a_ℓ for the class in $k_2^M(\mathbb{Q})$ that maps to $e_\ell + e_2$ in $\bigoplus k_2^M(\mathbb{Q}_v)$. For $n \geq 3$ the class ρ^n generates $k_n^M(\mathbb{Q})$. The product structure in $k_*^M(\mathbb{Q})$ can be deduced from the products given in Table 1; we write $(q \mid \ell)$ for the Legendre symbol that takes values in the additive group $\mathbb{Z}/2$.

Lemma 12 *The E_1 page of the h_1 -inverted ρ -Bockstein spectral sequence over \mathbb{Q} is the $\mathbb{Z}/2$ -algebra*

$$\mathfrak{B}(\mathbb{Q})[h_1^{-1}] \cong \bigoplus_{n \in \mathbb{N}} \rho^n k_*^M(\mathbb{Q}) / \rho^{n+1} \otimes_{\mathbb{F}_2} \text{Ext}(\mathbb{C})[h_1^{-1}].$$

The class ρ^n is in filtration $\epsilon = n$ for all $n \in \mathbb{N}$, for $\ell \equiv 3 \pmod{4}$ a prime a_ℓ is in filtration 1, for $\ell \equiv 1 \pmod{4}$ a prime a_p is in filtration 0, and $[p]$ for p a prime is in filtration 0. The r^{th} differential d_r for the ρ -Bockstein spectral sequence has degree $(\epsilon, f, t, c) = (r, 1, -1, 0)$. See Figure 3 for a chart of the E_1 page up to Milnor–Witt stem 15.

Proof The ρ -Bockstein spectral sequence arises from filtering the cobar complex $\mathcal{C}^*(\mathbb{Q})$ by powers of ρ . The s^{th} term of the cobar complex is

$$\mathcal{C}^s(\mathbb{Q}) = H_{**}(\mathbb{Q}) \otimes A_{**}(\mathbb{Q})^{\otimes s},$$




Figure 3: The E_1 page of the h_1 -inverted ρ -Bockstein spectral sequence over \mathbb{Q} up to Milnor–Witt stem 15. Black dots represent the group $\mathbb{Z}/2[h_1^{\pm 1}]$, blue dots represent $\bigoplus_{p \equiv 3(4)} \mathbb{Z}/2[h_1^{\pm 1}]$, red dots represent $\bigoplus_{p \equiv 1(4), p=2} \mathbb{Z}/2[h_1^{\pm 1}]$, and green dots represent $\bigoplus_{p \equiv 1(4)} \mathbb{Z}/2[h_1^{\pm 1}]$. Solid vertical lines indicate multiplication by ρ , and a vertical arrow means that the tower of ρ -multiplications continues indefinitely. Every dot supports an infinite tower of v_2 -multiples, however we only indicate this with lines and arrows of slope $\frac{1}{3}$ on the classes of $\text{Ext}(\mathbb{C})[h_1^{-1}]$ and $k_*^M(\mathbb{Q})$. The horizontal axis t is the Milnor–Witt stem, and the vertical axis c is the Chow weight, while the Adams filtration is suppressed.

where the tensor products are taken over $H_{**}(\mathbb{Q})$, taking care to use the left and right actions of $H_{**}(\mathbb{Q})$ on $A_{**}(\mathbb{Q})$ arising from the left and right units η_L and η_R . Any class $a[x_1 | \cdots | x_s]$ can be reduced to a sum of monomials $b[y_1 | \cdots | y_s]$, where each y_i is a monomial in $\mathbb{Z}/2[\tau_0, \tau_1, \dots, \xi_1, \xi_2, \dots]$. The class τ is killed after inverting h_1 ,




Figure 4: The E_∞ page of $\mathfrak{B}(\mathbb{Q})[h_1^{-1}]$, which is also $\text{Ext}(\mathbb{Q})[h_1^{-1}]$, up to Milnor–Witt stem 15. Black dots represent the group $\mathbb{Z}/2[h_1^{\pm 1}]$, blue dots represent $\bigoplus_{p \equiv 3(4)} \mathbb{Z}/2[h_1^{\pm 1}]$, red dots represent $\bigoplus_{p \equiv 1(4), p=2} \mathbb{Z}/2[h_1^{\pm 1}]$, and green dots represent $\bigoplus_{p \equiv 1(4)} \mathbb{Z}/2[h_1^{\pm 1}]$. Solid vertical lines indicate multiplication by ρ , and a vertical arrow means that the tower of ρ -multiplications continues indefinitely. Multiplication by v_2 is indicated by lines of slope $\frac{1}{3}$, and an arrow of slope $\frac{1}{3}$ indicates that the class supports a tower of v_2 -multiplications. The d_2 differentials of $\mathfrak{M}(\mathbb{Q})[h_1^{-1}]$ are indicated with red lines of slope -1 .

hence every element of $\mathcal{C}^S(\mathbb{Q})$ is a sum of monomials $b[y_1 | \cdots | y_s]$, where each y_i is a monomial in $\mathbb{Z}/2[\tau_0, \tau_1, \dots, \xi_1, \xi_2, \dots]$ and $b \in k_*^M(\mathbb{Q})$. The filtration of the cobar complex now is determined by the filtration of $k_*^M(\mathbb{Q})$ by powers of ρ . \square

Proposition 13 *The differentials for the h_1 -inverted ρ -Bockstein spectral sequence over \mathbb{Q} are determined by $d_{2n-1}(v_1^{2^n}) = h_1^{2^n} \rho^{2^n-1} v_n$ for $n \geq 2$ and $d_r(v_n) = 0$ for $r \geq 1$ and $n \geq 2$. (See Figure 4 for a chart of the E_∞ page up to Milnor–Witt stem 15).*

class	(f, t, c)	ρ -torsion	conditions
$h_1^{\pm 1}$	$(\pm 1, 0, 0)$	∞	
ρ	$(0, 0, 1)$	∞	
$[2] + \rho$	$(0, 0, 1)$	1	
$[\ell]P^k$	$(0, 0, 1) + k(4, 4, 4)$	1	ℓ prime, $\ell \equiv 1 \pmod{4}$, $k \geq 0$
$[\ell]P^k$	$(0, 0, 1) + k(4, 4, 4)$	2	ℓ prime, $p \equiv 3 \pmod{4}$, $k \geq 0$
$[2]P^k$	$(0, 0, 1) + k(4, 4, 4)$	1	$k \geq 1$
$P^{2k}v_2$	$(1, 3, 1) + k(8, 8, 8)$	3	$k \geq 0$
$P^{4k}v_3$	$(1, 7, 1) + k(16, 16, 16)$	7	$k \geq 0$
$P^{8k}v_4$	$(1, 15, 1) + k(32, 32, 32)$	15	$k \geq 0$
\vdots	\vdots	\vdots	\vdots

Table 2: Generators of $\text{Ext}(\mathbb{Q})[h_1^{-1}]$

Proof The injection $k_*^M(\mathbb{Q}) \rightarrow \prod_v k_*^M(\mathbb{Q}_v)$ extends to an injection of h_1 -inverted ρ -Bockstein spectral sequences at the E_1 page:

$$\mathfrak{B}(\mathbb{Q})[h_1^{-1}] \rightarrow \prod_v \mathfrak{B}(\mathbb{Q}_v)[h_1^{-1}].$$

The differentials in the h_1 -inverted ρ -Bockstein spectral sequence over \mathbb{Q}_p vanish for all primes p . Only the differentials in $\mathfrak{B}(\mathbb{R})[h_1^{-1}]$ contribute to the differentials over \mathbb{Q} , and these were identified by Guillou and Isaksen in [8, Lemma 3.1]. \square

Proposition 14 *The h_1 -inverted ρ -Bockstein spectral sequence for \mathbb{Q} converges strongly to $\text{Ext}(\mathbb{Q})[h_1^{-1}]$, and there are no hidden extensions.*

Proof The h_1 -inverted ρ -Bockstein spectral sequence is isomorphic to the ρ -Bockstein spectral sequence obtained by filtering the $[\xi_1]$ -inverted cobar complex $\mathcal{C}^*(\mathbb{Q})$, hence it converges strongly to $\text{Ext}(\mathbb{Q})[h_1^{-1}]$. Guillou and Isaksen have shown that there are no hidden extensions in the h_1 -inverted ρ -Bockstein spectral sequence over \mathbb{R} [8, Proposition 4.9] and there are no hidden extensions in the h_1 -inverted ρ -Bockstein spectral sequence over the other completions of \mathbb{Q} by Proposition 6. We therefore conclude there are no hidden extensions since the Hasse map embeds $\mathfrak{B}(\mathbb{Q})[h_1^{-1}]_\infty \Rightarrow \text{Ext}(\mathbb{Q})[h_1^{-1}]$ into $\prod_v \mathfrak{B}(\mathbb{Q}_v)[h_1^{-1}]_\infty \Rightarrow \prod_v \text{Ext}(\mathbb{Q}_v)[h_1^{-1}]$. \square

Corollary 15 *$\text{Ext}(\mathbb{Q})[h_1^{-1}]$ is generated by the classes in Table 2. The relations among these generators over $k_*^M(\mathbb{Q})$ include: $[\ell]P^k \cdot [q]P^j = [\ell] \cdot [q]P^{k+j}$ for ℓ, q primes and $k, j \geq 0$; $[\ell]P^{2^{n-1}k} \cdot v_n = [\ell] \cdot P^{2^{n-1}k}v_n$ for ℓ a prime, $n \geq 2$, and $k \geq 0$; the*




Figure 5: The E_3 page of $\mathfrak{M}(\mathbb{Q})[h_1^{-1}]$ up to Milnor–Witt stem 15. The d_3 differentials are indicated with blue lines of slope -2 . The notational conventions of Figure 4 apply here.

vanishing of the product of three or more generators of the form $[\ell]P^k$; and the relations which set the ρ -torsion of the generators.

Proof The generators can be determined by comparing $\text{Ext}(\mathbb{Q})[h_1^{-1}]$ to $\text{Ext}(\mathbb{R})[h_1^{-1}]$, and the latter was determined by Guillou and Isaksen in [8, Theorem 4.10]. The relations stated are present in the ρ -Bockstein spectral sequence and persist to $\text{Ext}(\mathbb{Q})$. \square

The differentials in $\mathfrak{M}(\mathbb{Q})[h_1^{-1}]$, the h_1 -inverted Adams spectral sequence over \mathbb{Q} , are determined by the differentials obtained from the comparison to \mathbb{Q}_p and \mathbb{R} . See Figures 4 and 5 for a depiction of the E_2 and E_3 pages up to Milnor–Witt stem 15.

class	(f, t, c)	ρ -torsion	conditions
$h_1^{\pm 1}$	$(\pm 1, 0, 0)$	∞	
ρ	$(0, 0, 1)$	∞	
$[2] + \rho$	$(0, 0, 1)$	1	
$[\ell]P^k$	$(0, 0, 1) + k(4, 4, 4)$	1	ℓ prime, $\ell \equiv 1 \pmod{4}$, $k \geq 0$
$[\ell]P^k$	$(0, 0, 1) + k(4, 4, 4)$	2	ℓ prime, $\ell \equiv 3 \pmod{4}$, $k \geq 0$
$[2]P^k$	$(0, 0, 1) + k(4, 4, 4)$	1	$k \geq 1$
$P^{2k}v_2$	$(1, 3, 1) + k(8, 8, 8)$	3	$k \geq 0$
$\rho^3 P^{4k}v_3$	$(1, 7, 4) + k(16, 16, 16)$	4	$k \geq 0$
$\rho^7 P^{8k}v_4$	$(1, 15, 8) + k(32, 32, 32)$	8	$k \geq 0$
\vdots	\vdots	\vdots	\vdots
$P^{4(2j+1)}v_3^2$	$(2, 14, 2) + (2j + 1)(4, 4, 4)$	7	$j \geq 0$
$P^{8(2j+1)}v_4^2$	$(2, 30, 2) + (2j + 1)(4, 4, 4)$	15	$j \geq 0$
\vdots	\vdots	\vdots	\vdots

Table 3: Generators of $\mathfrak{M}(\mathbb{Q})[h_1^{-1}]_3$

Proposition 16 *The d_2 differential in $\mathfrak{M}(\mathbb{Q})[h_1^{-1}]$ is determined by the Leibniz rule from the equations $d_2(P^{2^{n-1}k}v_n) = P^{2^{n-1}k}v_{n-1}^2$ for $k \geq 0$ and $n \geq 3$ and the vanishing of d_2 on the remaining generators. For $r \geq 3$, the differential d_r in $\mathfrak{M}(\mathbb{Q})[h_1^{-1}]$ is determined by the Leibniz rule from the equations*

$$d_r(\rho^{2^n - 2^{n-r+2} - r + 2} P^{2^{n-1}k} v_n) = P^{2^{n-1}k + 2^{n-2} - 2^{n-r}} v_{n-r+1}^2$$

for $n \geq r + 1$ and the vanishing of d_r on the remaining generators.

Proof $\text{Ext}(\mathbb{Q})[h_1^{-1}]$ injects into the product $\prod_v \text{Ext}(\mathbb{Q}_v)[h_1^{-1}]$ under the base-change maps obtained from $\mathbb{Q} \rightarrow \mathbb{Q}_v$. The map is seen to be injective by the explicit calculation of $\text{Ext}(\mathbb{Q})[h_1^{-1}]$ given in [Corollary 15](#), $\text{Ext}(\mathbb{Q}_p)[h_1^{-1}]$ in [Proposition 6](#), and $\text{Ext}(\mathbb{R})[h_1^{-1}]$ in [\[8, Theorem 4.10\]](#). The differentials $d_2(v_n) = v_{n-1}^2$ for $n \geq 3$ over \mathbb{Q}_p imply that the class $d_2(P^{2^{n-1}k}v_n)$ must map to $d_2(P^{2^{n-1}k}v_n) = P^{2^{n-1}k}v_{n-1}^2$ in $\text{Ext}(\mathbb{Q}_p)[h_1^{-1}]$. Comparison to \mathbb{R} also shows that the differential $d_2(P^{2^{n-1}k}v_n)$ maps to $P^{2^{n-1}k}v_{n-1}^2$ in $\text{Ext}(\mathbb{R})[h_1^{-1}]$ for $n \geq 3$, as determined by Guillou and Isaksen [\[8, Lemma 5.2\]](#). The differential d_2 over \mathbb{Q} vanishes on the classes $[\ell]P^k$ by comparison to \mathbb{Q}_p for all p . Finally, d_2 vanishes on all elements of $k_*^M(\mathbb{Q})$ for degree reasons. This accounts for all of irreducible classes of $\text{Ext}(\mathbb{Q})[h_1^{-1}]$; the generators for $\mathfrak{M}(\mathbb{Q})_3[h_1^{-1}]$ are given in [Table 3](#). Note that the classes $P^{2(2j+1)}v_2^2$ also survive but decompose as the product $P^{2(2j+1)}v_2 \cdot v_2$.




Figure 6: The E_∞ page of $\mathfrak{M}(\mathbb{Q})[h_1^{-1}]$ up to Milnor–Witt stem 15. The notational conventions of Figure 4 apply here.

The Hasse map

$$\mathfrak{M}(\mathbb{Q})_3[h_1^{-1}] \rightarrow \prod_v \mathfrak{M}(\mathbb{Q}_v)_3[h_1^{-1}]$$

is still injective. Over the p -adic fields, all further differentials vanish, and over \mathbb{R} the differentials are determined by Guillou and Isaksen [8, Lemma 5.8]; these comparisons determine the remaining differentials. \square

Proposition 17 *The E_∞ page of $\mathfrak{M}(\mathbb{Q})[h_1^{-1}]$ is generated over $k_*^M(\mathbb{Q})$ by the classes $\rho^{2^n-n-2} P^{2^{n-1}k} v_n$ for $k \geq 0$ and $n \geq 2$. Such a class has degree*

$$(2^{n+1}k + 1, 2^{n+1}k + 2^n - 1, 2^{n+1}k + 2^n - n - 1),$$

and its ρ -torsion is $n + 1$. (See Table 4 for some low-degree generators and Figure 6 for a chart of the E_∞ page up to Milnor–Witt stem 15).

class	(f, t, c)	ρ -torsion	conditions
$h_1^{\pm 1}$	$(\pm 1, 0, 0)$	∞	
ρ	$(0, 0, 1)$	∞	
$[2] + \rho$	$(0, 0, 1)$	1	
$[\ell]P^k$	$(0, 0, 1) + k(4, 4, 4)$	1	ℓ prime, $\ell \equiv 1 \pmod{4}$, $k \geq 0$
$[\ell]P^k$	$(0, 0, 1) + k(4, 4, 4)$	2	ℓ prime, $\ell \equiv 3 \pmod{4}$, $k \geq 0$
$[2]P^k$	$(0, 0, 1) + k(4, 4, 4)$	1	$k \geq 1$
$P^{2k}v_2$	$(1, 3, 1) + k(8, 8, 8)$	3	$k \geq 0$
$\rho^3 P^{4k}v_3$	$(1, 7, 4) + k(16, 16, 16)$	4	$k \geq 0$
$\rho^{10} P^{8k}v_4$	$(1, 15, 11) + k(32, 32, 32)$	5	$k \geq 0$
\vdots	\vdots	\vdots	\vdots

Table 4: Generators of $\mathfrak{M}(\mathbb{Q})[h_1^{-1}]_\infty$

Proof This is a consequence of the differential analysis of Proposition 16 and the result of Guillou and Isaksen [8, Proposition 5.9]. □

Definition 18 Let M be the submodule of $W(\mathbb{Q})_2^\wedge[\eta^{\pm 1}]$ generated by the rank-one forms $\ell \cdot X^2$ for ℓ a prime. As an abelian group, M is isomorphic to

$$\mathbb{Z}/2 \oplus \bigoplus_{p \equiv 3 \pmod{4}} \mathbb{Z}/4 \oplus \bigoplus_{p \equiv 1 \pmod{4}} (\mathbb{Z}/2)^2[\eta^{\pm 1}].$$

Following the notational convention of Guillou and Isaksen [8, Section 7], write $P^{2^{n-1}k}\lambda_n$ for a class in $\widehat{\Pi}_t(\mathbb{Q})[\eta^{-1}]$ detected by $\rho^{2^n-n-2}P^{2^{n-1}k}v_n$ in $\mathfrak{M}(\mathbb{Q})[h_1^{-1}]$, where $n \geq 2$, $k \geq 0$ and $t = 2^{n+1}k + 2^n - 1$. Also, we abuse notation and write $[\ell]P^k$ for a class in $\widehat{\Pi}_{4k}(\mathbb{Q})[\eta^{-1}]$ detected by the class of the same name in $\mathfrak{M}(\mathbb{Q})[h_1^{-1}]$.

Theorem 19 The η -inverted Milnor–Witt 0-stem of $\mathbb{1}_2^\wedge$ over \mathbb{Q} is

$$\widehat{\Pi}_0(\mathbb{Q})[\eta^{-1}] \cong W(\mathbb{Q})_2^\wedge[\eta^{\pm 1}].$$

The t^{th} η -inverted Milnor–Witt stem of $\mathbb{1}_2^\wedge$ over \mathbb{Q} is as follows:

$$\widehat{\Pi}_t(\mathbb{Q})[\eta^{-1}] \cong \begin{cases} \widehat{\Pi}_0(\mathbb{Q})[\eta^{-1}]/2^{n+1} & \text{if } t \geq 0, t \equiv 3 \pmod{4}, n = v_2(t + 1), \\ M & \text{if } t \equiv 0 \pmod{4}, t \geq 4, \\ 0 & \text{otherwise.} \end{cases}$$

Here $v_2(x)$ is the 2-adic valuation of an integer x , and M is the $\widehat{\Pi}_0(\mathbb{Q})[\eta^{-1}]$ -module of Definition 18.

The remaining product structure of $\widehat{\Pi}_*(\mathbb{Q})[\eta^{-1}]$ is determined by the following relations: the product of any two generators with Milnor–Witt stem congruent to 3 mod 4 is zero; $[q]P^j \cdot [\ell]P^k = [q] \cdot [\ell]P^{j+k}$ for all primes ℓ and q and $k, j \geq 0$; $[q] \cdot [\ell]P^k = 0$ if q is a prime or -1 and $[q][\ell] = 0$ in $k_*^M(\mathbb{Q})$.

Proof The zero stem was calculated in Proposition 1 and [16, Proof of Theorem 1.5] shows the one stem vanishes. Proposition 17 identifies the structure of the E_∞ page of the h_1 -inverted motivic Adams spectral sequence over \mathbb{Q} and Corollary 5 shows that $\mathfrak{M}(\mathbb{Q})[h_1^{-1}]$ strongly converges to $\widehat{\Pi}_*(\mathbb{Q})[\eta^{-1}]$ in Milnor–Witt stem at least 2. The 2-extensions are resolved because ρh_1 detects multiplication by 2, from which the additive structure of the η -inverted stems follows.

The product structure in the E_∞ page of the h_1 -inverted motivic Adams spectral sequence determines the $\widehat{\Pi}_0(\mathbb{Q})[\eta^{-1}]$ -module structure of the stems $\widehat{\Pi}_t(\mathbb{Q})[\eta^{-1}]$ for $t \not\equiv 3 \pmod 4$ and $t \equiv 3 \pmod 8$. It only remains to identify the hidden product of $[\ell]$ for ℓ a prime with a class $P^{2^{n-1}k}\lambda_n$ of $\widehat{\Pi}_t(\mathbb{Q})[\eta^{-1}]$ for $u \geq 3, k \geq 0$; note that $n = v_2(t + 1)$. Lemma 20 shows that the products $[\ell] \cdot P^{2^{n-1}k}\lambda_n$ and $a_\ell \cdot P^{2^{n-1}k}\lambda_n$ are always nonzero; hence the canonical map $\widehat{\Pi}_0(\mathbb{Q})[\eta^{-1}]/2^{n+1} \rightarrow \widehat{\Pi}_n(\mathbb{Q})[\eta^{-1}]$ is an isomorphism.

The product of any two generators with Milnor–Witt stem congruent to 3 mod 4 is zero for degree reasons. The remaining products are detected in the motivic Adams spectral sequence. □

Lemma 20 For $n \geq 3$ and $k \geq 0$, the products $[\ell] \cdot P^{2^{n-1}k}\lambda_n$ with ℓ a prime and $a_\ell \cdot P^{2^{n-1}k}\lambda_n$ with ℓ an odd prime in $\widehat{\Pi}_*(\mathbb{Q})[\eta^{-1}]$ are nonzero.

Proof For $m \geq 0$ and ℓ a prime, the Massey product $\langle \rho P^{2^m}v_2, \rho^2, [\ell] \rangle$ in $\text{Ext}(\mathbb{Q})[h_1^{-1}]$ contains $[\ell]P^{2^{m+1}}$ by Lemma 21 and this has no indeterminacy. The hypotheses of Moss’s convergence theorem [11, Theorem 3.1.1] hold here;¹ hence $[\ell]P^{2^{m+1}}$ detects a class of $\langle 2P^{2^m}\lambda_2, 2^2, [\ell] \rangle$. The indeterminacy of this Toda bracket is $[\ell]\widehat{\Pi}_{8m+4}$, which is in higher filtration than $[\ell]P^{2^{m+1}}$. We conclude $\langle 2P^{2^m}\lambda_2, 2^2, [\ell] \rangle$ does not contain zero.

The Massey product $\langle v_2, \rho P^{2^m}v_2, \rho^2 \rangle$ can be shown to contain $\rho^{2^n-n-2}P^{2^{n-1}k}v_n$ using the Adams differential

$$d_r(\rho^{2^n-n-4}P^{2^{n-1}k}v_n) = \rho P^{2^m}v_2^2,$$

¹Observe that for $r \geq 2, c > t$ and $t \equiv 3 \pmod 4$, the groups $E_2^{t,c}$ are trivial in the h_1 -inverted motivic Adams spectral sequence over \mathbb{Q} .

where $n = v_2(m + 1) + 3$ and $r = n - 1$. This Massey product has trivial indeterminacy. Moss's convergence theorem shows that $\rho^{2^n - n - 2} P^{2^{n-1}k} v_n$ detects a class of $\langle \lambda_2, 2P^{2^m} \lambda_2, 2^2 \rangle$; ² hence $P^{2^{n-1}k} \lambda_n$ is in the Toda bracket $\langle \lambda_2, 2P^{2^m} \lambda_2, 2^2 \rangle$.

We now use the shuffle relation

$$\lambda_2 \langle 2P^{2^m} \lambda_2, 2^2, [\ell] \rangle = \langle \lambda_2, 2P^{2^m} \lambda_2, 2^2 \rangle [\ell].$$

Multiplication by λ_2 is an injection on the stems $\widehat{\Pi}_{4j}(\mathbb{Q})[\eta^{-1}] \rightarrow \widehat{\Pi}_{4j+3}(\mathbb{Q})[\eta^{-1}]$ by the product structure in the motivic Adams spectral sequence, hence the left-hand side of the shuffle relation does not contain zero. As $[\ell] \cdot P^{2^{n-1}k} \lambda_n$ is in the right-hand side of the shuffle relation, we conclude that $[\ell] \cdot P^{2^{n-1}k} \lambda_n$ is nonzero.

A similar argument using the shuffle relation

$$\lambda_2 \langle 2P^{2^k} v_2, 2^2, a_\ell \rangle = \langle \lambda_2, 2P^{2^k} v_2, 2^2 \rangle a_\ell$$

establishes the claim that $a_\ell \cdot P^{2^u - 1} k$ is nonzero. □

Lemma 21 *Let $m \geq 0$ and ℓ a prime. The Massey product $\langle \rho P^{2^m} v_2, \rho^2, [\ell] \rangle$ in $\text{Ext}(\mathbb{Q})[h_1^{-1}]$ contains $[\ell] P^{2^{m+1}}$ and has trivial indeterminacy.*

Proof The ρ -Bockstein spectral sequence differential $d_3(P^{2^{m+1}}) = \rho^3 P^{2^m} v_2$ shows that $\langle \rho P^{2^m} v_2, \rho^2, [\ell] \rangle$ contains $[\ell] P^{2^{m+1}}$; this Massey product has trivial indeterminacy. To verify the hypotheses of May's convergence theorem [11, Theorem 2.2.1], first note the degree of $\rho^2[\ell]$ is $\epsilon = 2$, $t = 0$, and $c = 1$. All ρ -Bockstein spectral sequence differentials vanish in this graded component. It remains to check d_R differentials on the graded piece with $\epsilon' \geq 3$, $t = 8m + 3$, $c = 8m + 4$ and $R > \epsilon'$ corresponding to $\rho^3 P^{2^m} v_2$.

We now look for elements of the E_4 page of the ρ -Bockstein spectral sequence which land in degrees (t, c) for which $t + c \equiv 7 \pmod{8}$. Given the description of

²The condition to check for $\rho P^{2^m} v_2^2$ in Moss's theorem is that $d_{r'}: E_{r'}^{8m+7, c'} \rightarrow E_{r'}^{8m+6, c''}$ must be zero when $c' \leq 8m + 1 - v_2(m + 1)$, $c'' \geq 8m + 4$ and $r' = c'' - c'$. Nonzero differentials are only possible in these Milnor-Witt stems on classes $\rho^2 P^2 v_n$; it follows that $v_2(m + 1) = n - 3$, so $r' \geq n$. But by Proposition 16, nonzero differentials on such classes occur only when $n \geq r' + 1$. The condition to check for the element $\rho^3 P^{2^m} v_2$ is that $d_{r'}: E_{r'}^{8m+4, c'} \rightarrow E_{r'}^{8m+3, c''}$ must be zero when $c' \leq 8m + 2 - v_2(m + 1)$, $c'' \geq 8m + 5$ and $r' = c'' - c'$. The classes in Milnor-Witt stems $4 \pmod{8}$ are generated by the classes $\rho, h_1^{\pm 1}$ and $[\ell] P^k$. The Adams differentials vanish on these classes, so the hypotheses of Moss's theorem are true.

the generators of the E_4 page, it suffices to consider products of just v_n for $n \geq 3$, $[\ell]P$, v_2 , and ρ . The sum $t + c \pmod 8$ for each of these generators is 0, 1, 4, and 1, respectively. As $([\ell]P)^2 = 0$, $\rho^3 v_2 = 0$, and $\rho^2 [\ell]P v_2 = 0$ in the E_4 page, the only nonzero product of these generators in degree (t, c) with $t + c \equiv 7 \pmod 8$ must be of the form $\rho^\epsilon v_3^{a_3} \cdots v_i^{a_i}$.

Suppose now that $\rho^\epsilon v_3^{a_3} \cdots v_i^{a_i}$ is in degree $t = 8m + 3$, $c = 8m + 4$ for some m . Let $A = \sum a_i$ and $j = \min\{x \mid a_x \neq 0\}$. Under these assumptions it follows that

$$8m + 3 = \sum a_i (2^i - 1) \geq A(2^j - 1).$$

Hence $\epsilon \geq A(2^j - 2) + 1$. Note that A must be at least 2 in order for the Milnor–Witt stem t to be congruent to 3 modulo 8. It follows that if $R > \epsilon$, then $R > 2^j - 1$ and so the class $\rho^\epsilon v_3^{a_3} \cdots v_i^{a_i}$ is zero in the E_R page, by the relation $\rho^{2^j - 1} v_j = 0$, which arises from a $d_{2^j - 1}$ differential. We conclude May’s convergence theorem applies in this situation. It is straightforward to check that the indeterminacy is trivial. \square

References

- [1] **A Ananyevskiy, M Levine, I Panin**, *Witt sheaves and the η -inverted sphere spectrum*, J. Topol. 10 (2017) 370–385 [MR](#)
- [2] **M Andrews, H Miller**, *Inverting the Hopf map*, preprint (2017) [arXiv](#)
- [3] **JM Boardman**, *Conditionally convergent spectral sequences*, from “Homotopy invariant algebraic structures” (J-P Meyer, J Morava, W S Wilson, editors), Contemp. Math. 239, Amer. Math. Soc., Providence, RI (1999) 49–84 [MR](#)
- [4] **A K Bousfield**, *The localization of spectra with respect to homology*, Topology 18 (1979) 257–281 [MR](#)
- [5] **A K Bousfield, DM Kan**, *Homotopy limits, completions and localizations*, Lecture Notes in Math. 304, Springer (1972) [MR](#)
- [6] **D Dugger, DC Isaksen**, *Motivic Hopf elements and relations*, New York J. Math. 19 (2013) 823–871 [MR](#)
- [7] **B J Guillou, DC Isaksen**, *The η -local motivic sphere*, J. Pure Appl. Algebra 219 (2015) 4728–4756 [MR](#)
- [8] **B J Guillou, DC Isaksen**, *The η -inverted \mathbb{R} -motivic sphere*, Algebr. Geom. Topol. 16 (2016) 3005–3027 [MR](#)
- [9] **P Hu, I Kriz, K Ormsby**, *Convergence of the motivic Adams spectral sequence*, J. K-Theory 7 (2011) 573–596 [MR](#)
- [10] **DC Isaksen**, *Classical and motivic Adams charts*, preprint (2014) [arXiv](#)

- [11] **D C Isaksen**, *Stable stems*, preprint (2014) [arXiv](#)
- [12] **M Knebusch**, *Symmetric bilinear forms over algebraic varieties*, from “Conference on Quadratic Forms — 1976” (G Orzech, editor), Queen’s Papers in Pure and Appl. Math. 46, Queen’s Univ., Kingston, Ont. (1977) 103–283 [MR](#)
- [13] **J Milnor**, *Algebraic K-theory and quadratic forms*, Invent. Math. 9 (1969/1970) 318–344 [MR](#)
- [14] **F Morel**, \mathbb{A}^1 -*algebraic topology over a field*, Lecture Notes in Math. 2052, Springer (2012) [MR](#)
- [15] **K M Ormsby, P A Østvær**, *Motivic Brown–Peterson invariants of the rationals*, Geom. Topol. 17 (2013) 1671–1706 [MR](#)
- [16] **K Ormsby, O Röndigs, P A Østvær**, *Vanishing in stable motivic homotopy sheaves*, preprint (2017) [arXiv](#)
- [17] **W Scharlau**, *Quadratic and Hermitian forms*, Grundle. Math. Wissen. 270, Springer (1985) [MR](#)
- [18] **J-P Serre**, *A course in arithmetic*, Graduate Texts in Math. 7, Springer (1973) [MR](#)
- [19] **C A Weibel**, *An introduction to homological algebra*, Cambridge Studies in Adv. Math. 38, Cambridge Univ. Press (1994) [MR](#)
- [20] **G M Wilson, P A Østvær**, *Two-complete stable motivic stems over finite fields*, Algebr. Geom. Topol. 17 (2017) 1059–1104 [MR](#)

Department of Mathematics, University of Oslo
Oslo, Norway

glenw@math.uio.no

Received: 30 August 2017 Revised: 26 October 2017

